

Effects of dietary arginine in ameliorating the deleterious effects induced by mycotoxins on growth, immune system, body organs in growing pigs

Joseph Moturi^{1#}, SeYoung Yoon^{1#}, Abdolreza Hosseindoust¹, SangHun Ha¹, Habeeb Tajudeen¹, JunYoung Mun¹, SooJin Sa^{2*} and JinSoo Kim^{1*}

¹Department of Animal Industry Convergence, Kangwon National University, Chuncheon 24341, Korea

²Swine Division, National Institute of Animal Science, Rural Development Administration, Cheonan 31000, Korea



Received: Apr 18, 2022

Revised: Jun 18, 2022

Accepted: Jun 20, 2022

#These authors contributed equally to this work.

*Corresponding author

SooJin Sa

Swine Division, National Institute of Animal Science, Rural Development Administration, Cheonan 31000, Korea.

Tel: +82-41-580-3450

E-mail: soojinsa@korea.kr

JinSoo Kim

Department of Animal Industry Convergence, Kangwon National University, Chuncheon 24341, Korea.

Tel: +82-33-250-8616

E-mail: kjs896@kangwon.ac.kr

Copyright © 2022 Korean Society of Animal Sciences and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Joseph Moturi

<https://orcid.org/0000-0002-0626-0255>

SeYoung Yoon

<https://orcid.org/0000-0001-6601-8201>

Abstract

Mycotoxin contamination in pig feeds has a negative impact on growth performance, the immune system, and major body organs. Arginine (Arg) plays an important role in animals' body biochemistry and physiology. This study aimed to determine the effect of dietary Arg supplementation on mitigating the negative effects of mycotoxins in growing pigs. A total of 72 growing pigs (Landrace × Large white) with initial mean body weight (BW) = 55 ± 2.5 kg were allotted to four treatment groups with three replicates per group of six pigs per replicate in a completely randomized design. The treatments included a non-toxin diet with 1.2% Arg (NT1.2) and mycotoxin-challenged treatments supplemented with 1.2% Arg (TX1.2), 1.3% Arg (TX1.3), and 1.4% Arg (TX1.4). Statistical analysis of data included the effects of dietary level of Arg. The results indicated a significantly higher BW ($p < 0.05$), average daily gain ($p < 0.05$), and gain-to-feed ratio ($p < 0.05$) in the NT1.2 group than in the TX1.2, TX1.3, and TX1.4 groups. The relative weight of the liver was higher ($p < 0.05$) in the TX1.2 compared to that of the NT1.2 group, although it was not different from that of TX1.3 and TX1.4. The level of tumor necrosis factor-alpha was significantly up-regulated ($p < 0.05$) in the liver tissue of the TX1.2 group compared to that of the other treatments. Overall, dietary Arg supplementation remedied liver injury and alleviated the compromised immune system caused by mycotoxin toxicity.

Keywords: Mycotoxin, Aflatoxin, Cytokines, Villus height, Immunity, Toll like receptor

INTRODUCTION

Mycotoxins are toxic secondary metabolites formed by several molds, especially *aspergillus*, *penicillium*, and *fusarium* [1] that could contaminate food and feed at various stages in the animal feed chain. Despite numerous modern and improved management practices implemented in the agricultural and

Abdolreza Hosseindoust
<https://orcid.org/0000-0001-9191-0613>
SangHun Ha
<https://orcid.org/0000-0003-3779-1144>
Habeeb Tajudeen
<https://orcid.org/0000-0002-5623-3175>
JunYoung Mun
<https://orcid.org/0000-0002-3075-7157>
SooJin Sa
<https://orcid.org/0000-0002-2634-5109>
JinSoo Kim
<https://orcid.org/0000-0002-9518-7917>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was carried out with the support of Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ01504401 Rural Development Administration, Korea).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Moturi J, Hosseindoust A, Kim JS.
Data curation: Moturi J, Yoon SY, Ha SH, Sa SJ.
Formal analysis: Moturi J, Hosseindoust A, Tajudeen H.
Methodology: Moturi J, Hosseindoust A.
Software: Yoon SY, Hosseindoust A, Sa SJ.
Validation: Ha SH, Mun JY.
Writing - original draft: Moturi J, Hosseindoust A.
Writing - review & editing: Moturi J, Yoon SY, Hosseindoust A, Ha SH, Tajudeen H, Mun JY, Sa SJ, Kim JS.

Ethics approval and consent to participate

The project underwent proper ethical standards and the experiments were approved by the Institutional Animal Care and Use Committee of Kangwon National University (210503-6), Chuncheon, Korea.

feed manufacturing industries, mycotoxin contamination of animal diets remains a challenge [2]. Pigs are sensitive to mycotoxins ingestion [3]. Furthermore, pig diet formulations are dependent on cereals, a high-risk factor for mycotoxin exposure [4]. The major mycotoxins that contaminate animal feeds include aflatoxins, ochratoxin, zearalenone, fumonisins, trichothecenes, deoxynivalenol (DON), trichothecene-2 toxin, and hydroxytrichothecene-2 toxin [3]. Mycotoxin contamination decreases growth efficiency, lowers feed conversion rate, reproductive rates, and impairs resistance to diseases in domestic animals such as dairy cattle, sheep, swine, and poultry [5,6]. They increase the vulnerability of animals to infectious diseases, decrease vaccine effectiveness, and cause pathological harm to body tissues, kidneys, liver, and the reproductive, nervous, and immune systems [7,8]. Some mycotoxins are carcinogenic, genotoxic, and teratogenic [9]. Consequently, it negatively affects the health of animals. Overall, apart from the negative impacts on production imposed by mycotoxin contamination in swine production, farmers equally incur huge costs to combat the harmful effects caused by mycotoxins in animals. This increases the cost of production hence diminishing pork producers' profits. This calls for cost-effective strategies to overcome the challenges of mycotoxins in livestock production.

Most studies on mycotoxicosis alleviation have focused on methods of physical and chemical mycotoxin degradation [10] and the use of adsorbents, including aluminosilicates [11] and esterified glucomannan [12]. However, *in vivo* studies have shown that these approaches are not always effective against chemically diverse mycotoxins. [11]. Currently, research on nutritional therapy to treat mycotoxin-induced effects in swine has focused on dietary supplementation of amino acids (AA) in mycotoxin compromised animals' diets. AA are nutritional components that have been found to enhance the immune function of animals. Some are essential for regulating major metabolic pathways; hence, they are known as functional AA [13]. This class of AA includes arginine (Arg) and methionine. Previous studies [14,15] conducted on pigs supplemented with higher amounts of Arg or methionine showed an improvement in growth performance and antioxidant status during heat stress. L-Arg is a conditionally essential AA in swine [16]. It stimulates intestinal development and maturation [17], further, it increases phagocytic activity, and accelerates the elimination of endotoxins in the gut and other organs [18]. Furthermore, mucosal microcirculation and absorption are enhanced by Arg [19]. It promotes the expression of anti-oxidative genes [20] through the nitric oxide (NO) synthesis mechanism. Moreover, Arg plays a vital role in swine nutrition regulatory functions, especially in immune response and nutrient metabolism [21]. Therefore, the present study aimed to evaluate the beneficial effects of L-Arg supplementation in growing pigs fed a mycotoxin-contaminated diet on growth performance, immune system, and major body organs

MATERIALS AND METHODS

Animal care

The experiment was conducted with proper observation of ethical standards and institutional protocols approved by the Kangwon National University Animal Welfare and Use Committee in the Republic (210503-6).

Animals, experimental design, and diets

This study was performed in Kangwon National University Animal Farm, Chuncheon, Korea. Seventy-two cross-bred (Landrace × Large white) 21-day-old piglets of mixed sex, 1/2 of the entire males and 1/2 females of initial body weight (BW) 55.2 ± 2.5 kg were randomly allotted to one of the four dietary treatments; NT1.2 (non- toxin basal diet with 1.2% Arg), TX1.2 (mycotoxin-

contaminated basal diet with 1.2% Arg), TX1.3 (mycotoxin-contaminated basal diet with 1.3% Arg), and TX1.4 (mycotoxin-contaminated basal diet with 1.4% Arg). The contaminated feed was mildewed under ambient conditions (temperature, 23 °C–28 °C; humidity, 68%–85%), as described by Liu et al. [11]. Liquid chromatography (Beijing Taileqi, Beijing, China) was used to detect the mycotoxins (Table 1). Basal diets were composed of a corn-soybean meal formulated as recommended by the NRC [22] to meet or exceed the nutritional requirements of growing pigs (Table 2). Arg was supplemented at the rate of 1.2%, 1.3%, and 1.4% per kg of the basal diet throughout the 20 days experimental period. Every treatment had three replicate pens composed of six pigs per pen (3 male; 3 female) group-housed. The experimental room contained 12 pens (2.28 × 1.85 = 4.2 m²) per pen with slatted floors. The room was equipped with automated fan-assisted central heating, forced ventilation and artificial lighting to ensure a minimum of 12 h of light/day. Every pen was provided with; one hopper-type feeder with two feeding points and a nipple drinker. Throughout the experiment, piglets had unlimited access to water and feed. Routine analysis of the diet and drinking water was performed for any contaminant that could affect the integrity of the study. On day 20 of the study, two pigs from each treatment group were euthanized by approved anesthesia and exsanguination, and tissue samples were harvested.

Sample collection and analyzes

Blood samples

Blood samples were collected from two pigs per group from the jugular vein and centrifuged at 3,000 rpm for 10 min at 4 °C. The serum was transferred to Eppendorf tubes and stored at –20 °C until analysis. Interleukin (IL)-4, tumor necrosis factor α (TNF- α), IL-6, toll-like receptor-4 (TLR-4), and interferon-gamma (IFN γ) levels were measured using ELISA test kits (Beijing Laboratory Biotech, Beijing, China).

Organ weights

The kidneys, spleen, heart, and liver were removed and weighed, and the organ weight and total BW percentage were recorded.

Intestinal histomorphology

Gut samples were collected from the duodenum, jejunum, and ileum, frozen in liquid nitrogen at –80 °C, and stored for analysis of intestinal histomorphology. The duodenal, jejunal, and ileal samples were cut into 5 cm lengths and fixed for 24 h in 10% neutral buffered formalin, transferred to a solution of 70% ethanol, embedded in wax, sectioned, and stained with eosin and hematoxylin. Slices were mounted on slides for analysis as described by Hosseindoust et al. [23].

Table 1. Mycotoxin volume

Item	Volume
Aflatoxins B1 (µg/kg) (ppb)	392
Aflatoxins B2 (µg /kg) (ppb)	136
Aflatoxins G1 (µg /kg) (ppb)	290
Aflatoxins G2 (µg /kg) (ppb)	ND
Ochratoxin A (µg /kg) (ppb)	158
DON (mg/kg)	0.73
FB1	ND
T-2	ND

ND, not detected; DON, deoxynivalenol; FB1, fumonisin B (1); T-2, tricothecene-2.

Table 2. Formula and chemical composition of basal diets (as-fed basis)

Items	Basal diet			
	1.2	1.2	1.3	1.4
Ingredient (%)	100	100	100	100
Corn	72.44	72.34	72.24	72.14
SBM dehulled	20.20	20.20	20.20	20.20
Blood meal	3.00	3.00	3.00	3.00
Soy oil	2.22	2.22	2.22	2.22
L-Lys (78%)	0.02	0.02	0.02	0.02
Arg 100%	0.00	0.10	0.20	0.30
Limestone	1.30	1.30	1.30	1.30
MCP	0.52	0.52	0.52	0.52
Vitamin premix ¹⁾	0.10	0.10	0.10	0.10
Mineral premix ²⁾	0.20	0.20	0.20	0.20
Chemical composition (%)				
Metabolizable energy	12.46	12.46	12.46	12.46
Crude protein	18.00	18.00	18.00	18.00
Crude fat	4.59	4.59	4.59	4.59
Crude fiber	2.49	2.49	2.49	2.49
Ash	3.72	3.72	3.72	3.72
Calcium	0.74	0.74	0.74	0.74
Available phosphorus	0.54	0.54	0.54	0.54
Lys	1.00	1.00	1.00	1.00
Met	0.28	0.28	0.28	0.28
Met_Cys	0.57	0.57	0.57	0.57
Thr	0.68	0.68	0.68	0.68
Trp	0.20	0.20	0.20	0.20
Arg	1.10	1.20	1.30	1.40

¹⁾Supplied per kilogram of diet: 16,000 IU vitamin A, 3,000 IU vitamin D₃, 5.0 mg vitamin B₁, 20 mg vitamin B₂, 4 mg vitamin B₆, 0.08 mg vitamin B₁₂, 40 IU vitamin E, 5.0 mg vitamin K₃, 75 mg niacin, 40 mg pantothenic acid, 0.15 mg biotin, 0.65 mg folic acid.

²⁾Supplied per kilogram of diet: 0.25 mg Co, 50 mg Cu, 15 mg Mn, 25 mg Zn, 45 mg Fe, 0.35 mg I, 0.13 mg Se.

SBM, soybean meal; MCP, monocalcium phosphate; Lys, lysine; Cys, cysteine; Met, methionine; Thr, threonine; Trp, tryptophan; Arg, arginine.

Intestinal morphology was evaluated by measuring the height and crypt depths (CD) of the most prominent villi from each of the three sections using an Olympus Vanox-S Microscope (Olympus Corporation, Lake Success, NY, USA) and analyzed using Spot basic imaging software (Diagnostic Instruments, Sterling Heights, MI, USA).

Growth performance

All the experimental pigs were weighed individually on the first and last day of the experiment. Feed consumption was also recorded to calculate the average daily feed intake (ADFI), average daily gain (ADG), and gain: feed ratio.

RNA extraction and cDNA synthesis

Complete RNA was extracted from liquid nitrogen-pulverized tissues using TRIzol reagent (Invitrogen, Waltham, MA, USA) and treated with DNase I (Invitrogen) as directed by the manufacturer and described by Hosseindoust et al. [24]. After staining with 10 g/mL ethidium

bromide, RNA purity was evaluated using 1 percent agarose gel electrophoresis. The optical density (OD)₂₆₀ : OD₂₈₀ ratio of the RNA was between 1.8 and 2.0. First-strand cDNA was synthesized using oligo (dT)₂₀ and Superscript II reverse transcriptase (Invitrogen).

mRNA quantification by real-time polymerase chain reaction analysis

Primer 5.0 was used to design primers based on the pig's cDNA sequence (Table 3). To standardize the target gene transcript levels, β -actin was used as the housekeeping gene. SYBR Green PCR Mix, comprising MgCl₂, dNTP, and Hotstar Taq polymerase, was used for real-time PCR. cDNA template (2 μ L) was added to a 25 μ L mixture including 12.5 μ L of SYBR Green solution and 1 mol/L forward and reverse primers. The following protocol was used: (a) pre-denaturation (10 s at 95 °C); (b) amplification and quantification, repeated 40 times (5 s at 95 °C, 20 s at 60 °C); and (c) melting curve (60 °C–99 °C with a heating rate of 0.1 °C s⁻¹ and fluorescence measurement).

Statistical analysis

All the results were expressed as mean \pm standards error of the mean, statistical analyses were done using the GLM procedure of SAS (SAS Institute, Cary, NC, USA) based on a completely randomized design. Significant differences of ($p < 0.05$) among the treatment were considered statistically significant differences using Turkey's Honestly Significant Difference procedure. Pig was considered as an experimental unit for all parameters.

RESULTS

Growth performance

Growth performance, as measured and evaluated by BW, ADG, ADFI, and feed conversion efficiency, is presented in Table 4. Overall, final BW ($p < 0.05$), ADG ($p < 0.05$), and gain: feed ratio ($p < 0.05$) was significantly higher in the NT1.2 treatment group as compared to all three toxin treatment groups (TX1.2, TX1.3, and TX1.4). No significant differences were observed in the ADFI ($p > 0.05$) among the groups.

Table 3. Primers

Gene	Primer sequence (5'__>3')
β actin_F	CAACACAGTGCTGTCTGGTGGTA
β actin_R	ATCGTACTCCTGCTTGCTGATCC
IL4_F	TGTGCCACGCTGTGCTTACA
IL4_R	CTTGTGGCAGTGCTGGCTCTCC
IL6_F	AGAAATCCCTCCTCGCCAAT
IL6_R	AAATAGCGAACGGCCCTCA
IFN γ _F	CTGAAGAAGTGGACAGAGAG
IFN γ _R	CACCAGCTTCTGTAAGATGC
HSP27_F	GGAGATCACCGGCAAACACG)
HSP27_R	CCTCCACTGTCCAGCATCCCA
TLR4_F	GTCTCTCCTTCTTACCTGCTGTTCC
TLR4_R	AGGAGGAGAAAGACAGGGTAGGTG
TNF- α	CCCCAGAAGGAAGAGTTTC

IL, interleukin; IFN, interferon; HSP, heat shock protein; TLR, toll-like receptor; TNF, tumor necrosis factor.

Table 4. Effects of dietary L-arginine supplementation on growth performance of growing pigs fed a mold-contaminated diet¹⁾

Item	Non-toxin (Arg 1.2%)	Toxin with Arg			SEM	p-value
		1.2	1.3	1.4		
Initial weight (kg)	24.14	24.07	23.92	24.05	0.270	0.952
Final weight (kg)	43.92 ^a	39.82 ^b	40.34 ^b	40.04 ^b	0.590	< 0.001
ADG (g)	707 ^a	563 ^b	586 ^b	571 ^b	25.4	0.002
ADFI (g)	989	995	1,005	994	21.7	0.836
Gain:feed ratio	0.71 ^a	0.56 ^b	0.58 ^b	0.57 ^b	0.030	0.007

¹⁾Data represents means based on six replicates per treatment (n = 6).

^{a,b}Means with different superscripts within rows are significantly different at ($p < 0.05$).

Arg, arginine; ADG, average daily gain; ADFI, average daily feed intake.

Organ relative weight

There was no significant difference ($p > 0.05$) in the relative weights of the kidney, heart, and spleen among the groups. However, the relative weight of the liver was significantly higher ($p < 0.05$) in the TX1.2 group compared with the NT1.2 and both the TX1.3 and TX1.4 groups, with the NT1.2 group having the lowest liver weight (Table 5).

Intestinal histomorphology

The villous height (VH) in the duodenal section of the gut in the NT1.2 group was statistically higher, although not significant ($p = 0.074$; Table 6). No significant differences ($p > 0.05$) were observed in the jejunum and ileum of the intestine in terms of VH, CD, and VH:CD ratio among the groups.

Cytokines and toll-like receptor-4 gene expression in the liver, muscle, and jejunum

There were no significant differences ($p > 0.05$) in the liver IL-4, IL-6, IFN γ , and TLR-4 levels. The mRNA level of *TNF α* in the liver tissue was significantly down-regulated in the non-toxin, TX1.3, and TX1.4 treatment groups as compared with the TX1.2 at ($p < 0.05$; Figs. 1A–1E). In contrast, significant differences were observed in mRNA level changes in muscle *IL-6* ($p < 0.05$) and *TNF α* ($p < 0.05$) between the NT1.2 and the TX1.2 treatment groups, although not different from TX1.3 and TX1.4. The mRNA levels of the other cytokines and *TLR-4* were not significantly different ($p > 0.05$) among the groups (Figs. 2A–2E). There was no statistical difference ($p > 0.05$) in the changes in the mRNA levels of all cytokines and *TLR-4* in the jejunum among the groups (Figs. 3A–3E).

Table 5. Effects of dietary L-arginine supplementation on relative organ weight (g/kg BW) of growing pigs fed a mold-contaminated diet¹⁾

Item	Non-toxin (Arg1.2%)	Toxin with Arg			SEM	p-value
		1.2	1.3	1.4		
Heart	0.54	0.50	0.51	0.55	0.030	0.538
Liver	2.46 ^b	2.65 ^a	2.54 ^{ab}	2.55 ^{ab}	0.030	0.006
Spleen	0.21	0.18	0.19	0.20	0.020	0.778
Kidney	0.48	0.53	0.50	0.52	0.040	0.808

¹⁾Data represents means based on six replicates per treatment (n = 6).

^{a,b}Means with different superscripts within rows are significantly different at ($p < 0.05$).

BW, body weight; Arg, arginine.

Table 6. Effects of dietary L-arginine supplementation on intestinal morphology in growing pigs fed a mold-contaminated diet¹⁾

Item	Non-toxin (Arg1.2%)	Toxin with Arg			SEM	p-value
		1.2	1.3	1.4		
Villus height (µm)						
Duodenum	549	428	478	492	30.5	0.074
Jejunum	448	414	442	439	19.7	0.638
Ileum	348	330	318	342	30.6	0.897
Crypt depth (µm)						
Duodenum	275	262	233	228	27.5	0.574
Jejunum	263	240	244	238	8.9	0.206
Ileum	226	248	239	231	29.4	0.953
VH/CD						
Duodenum	2.06	1.73	2.10	2.19	0.180	0.392
Jejunum	1.73	1.74	1.83	1.84	0.110	0.862
Ileum	1.65	1.59	1.35	1.55	0.250	0.844

¹⁾Data represent means based on six replicates per treatment (n=6).

Arg, arginine; VH/CD, villous height: crypt depth.




Fig. 1. Relative expression of IL-4, IL-6, IFN γ , TLR4 and TNF α in liver of growing pigs. Error bars represent standard error of means. Bars with different letters (a,b) differ significantly across all 4 treatment groups ($p < 0.05$). IL, interleukin; Arg, arginine; IFN, interferon; TLR, toll like receptor; TNF, tumor necrosis factor.




Fig. 2. Relative expression of IL-4, IL-6, IFN γ , TLR4 and TNF α in muscle of growing pigs. Error bars represent standard error of means. Bars with different letters (a,b) differ significantly across all 4 treatment groups ($p < 0.05$). IL, interleukin; Arg, arginine; IFN, interferon; TLR, toll like receptor; TNF, tumor necrosis factor.

DISCUSSION

L-Arg is an essential AA that is a precursor for synthesizing urea, NO, polyamines, creatine, and proteins and modulates the intestinal gene expression of cationic AA transporters and antioxidants [16,20]. Arg is vital for growth performance improvement and the morphological development of the small intestine in growing pigs fed on a mold-contaminated diet [15,16]. Results from a previous study showed that the consumption of mycotoxin-contaminated feeds led to decreased ADG and ADFI, thereby lowering growth performance in pigs [13]. In this study, mycotoxin contamination negatively affected BW, ADG, and feed conversion efficiency. These results are consistent with a previous study that reported a significant decrease in pig performance by the ingestion of DON-contaminated feeds [24]. Previous studies have shown great damage to intestinal function through an impaired gut barrier, damage to the intestinal structure, and reduced immunity [25,26]. Disruption of intestinal function interferes with the normal absorption of the end products of digestion into the bloodstream; hence, fewer nutrients are available for growth [27]. In our study, supplementation with various levels of Arg did not mitigate growth suppression caused by mycotoxins. Similar to our results, a previous study on supplementation with glutamine and Arg




Fig. 3. Relative expression of IL-4, IL-6, IFN γ , TLR4 and TNF α in jejunum of growing pigs. Error bars represent standard error of means. Bars with different letters (a,b) differ significantly across all 4 treatment groups ($p < 0.05$). IL, interleukin; Arg, arginine; IFN, interferon; TLR, toll like receptor; TNF, tumor necrosis factor.

failed to overcome growth suppression induced by mycotoxins in grower pigs [13]. However, in contrast to our findings, previous studies have shown positive results in terms of a higher growth rate on supplementation with peptides and AA preparations in alleviating mycotoxicosis in pigs and mice [28,29]. This may have been due to the combined effects of various peptides, each with an additive effect contrary to our study with only Arg as the only AA supplemented hence lacking the complementary advantage when many peptides and AA are supplemented.

Because of its detoxification function, the liver is a target organ for mycotoxins, leading to edema and liver enlargement [3]. Similarly, Yarru et al. [30] reported an increase in liver weight and accumulation of fat in the liver, resulting in disruption of hepatic functions. In our current study, the weight of the liver was significantly higher in the TX1.2 group compared to that of the NT1.2, TX1.3, and TX1.4 groups. The lower liver weights in the TX1.3 and TX1.4 groups could have resulted from attenuation of the toxic effect of mycotoxins with higher levels of Arg supplementation, which could have improved the hepatic function of the liver, preventing edema. These findings imply that Arg protect the immune system and minimize liver harm in pigs exposed to mycotoxin. We hypothesized that Arg could increase NO production in the small intestine and other tissues, which is important for regulating the antioxidant defense system and scavenging

excess reactive oxygen species induced by moldy feeds, thus improving the balance between reactive oxygen species production (e.g., superoxide anion, hydrogen peroxide, and hydroxyl radical) and biological defense against their toxicity. Similar to our results, a previous study reported a 10.7% increase in liver weight in a DON-stressed toxin control group compared to a non-toxin control in growing pigs [13]. Furthermore, Trenholm et al. [31] reported an increase in kidney and liver weights in growing pigs fed diets with higher DON concentrations and prolonged ingestion of feed contaminated with mycotoxins. However, a previous study [32] found no significant weight change in the liver, spleen, or kidneys of growing pigs fed a diet contaminated with aflatoxins and DON. In our current study, there was no significant difference in the relative weights of the kidney, heart, and the spleen among the treatments, we attributed this to be due to increased protein synthesis as a result of Arg supplementation and organ expansion induced by severe mycotoxin systemic toxicity interacting competitively to influence relative organ size

IL-6, IFN γ , and TNF- α are pro-inflammatory cytokines associated with porcine immune status. TNF- α is involved in systemic inflammation [33]. It is one of the first response cytokines produced following mycotoxin exposure, especially ochratoxin [34], when the immune system is stimulated. Therefore, its gene expression is up-regulated in the liver and muscle following mycotoxin toxicity. The liver is the first organ to offer a protective barrier against mycotoxin absorption in the gut because it prevents the systemic distribution of toxins throughout the body [35]. In our current study, the expression of both IL-6 and TNF- α were significantly lower in the NT1.2, TX1.3, and TX1.4 treatments compared to the TX1.2 treatment. This may be due to the protective effects of the increased Arg level on the immune system, thereby down-regulating TNF- α and IL-6 production and reducing inflammatory responses in both the muscle and liver. We hypothesized that this may have been due to NO production as a result of Arg metabolism. NO is a vital signaling molecule that plays a role in, vascular homeostasis, neurotransmission, immunological modulation, and host defense [18,21,36] hence counteracting the suppressive effects and abnormal regulation of metabolic pathways and immunological responses [37]. Similar to our findings, Zhou et al. [38] reported an increased expression of TNF- α and IL-2 mRNA in pigs fed a diet contaminated with DON. TLR-4 is a pattern-recognition receptor subtype expressed at high levels along the intestinal tract. It acts as the first immune barrier in the gastrointestinal innate immune system. TLR-4 plays a key role in translating and transcription of inflammatory mediators [39]. Our research found that the level of TLR-4 expression was significantly higher in the TX1.2 treatment compared to that of the NT1.2 group. This could result from the toxicity eliciting the up-regulation in the level of TLR-4. Notably, the expression level was not significantly different in the TX1.3 and TX1.4 groups compared to that of both the negative and positive controls, which could indicate the ability of higher levels of Arg to counteract the effects of mycotoxin toxicity, hence modulating immune responses. As a result, we hypothesized that exogenous Arg would improve animals' metabolism engaged in mitogen-activated protein kinase and nuclear factor-kappaB pathways, reducing the impairment generated by mycotoxins stress and improving immune important cytokines such as IL-4 which is an anti-inflammatory cytokine even though in this study no significant difference was observed across the groups in all the tissues under study. This may have been due to the concentration of the mycotoxins used and the relatively shorter period (20 days) which the animals were exposed to the treatment diets.

In conclusion, high dietary Arg supplementation in mycotoxin-contaminated growing pigs significantly reduced morphological liver injury induced by toxicity from ingesting mycotoxin-contaminated feed. Furthermore, higher levels of Arg significantly down-regulated the pro-inflammatory cytokine TNF- α , protecting the immune system and even the liver. This improved the health of mycotoxin-compromised animals.

REFERENCES

1. Smith RF. Cast (Council for Agricultural Science and Technology). *Bull Entomol Soc Am*. 1976;22:31-6. <https://doi.org/10.1093/besa/22.1.31>
2. Bryden WL. Mycotoxins in the food chain: human health implications. *Asia Pac J Clin Nutr*. 2007;16:95-101.
3. Bennett JW, Klich M. Mycotoxins. *Clin Microbiol Rev*. 2003;3:497-516. <https://doi.org/10.1128/cmr.16.3.497-516.2003>
4. Pierron A, Alassane-Kpembi I, Oswald IP. Impact of mycotoxin on immune response and consequences for pig health. *Anim Nutr*. 2016;2:63-8. <https://doi.org/10.1016/j.aninu.2016.03.001>
5. Fokunang CN, Tembe-Fokunang EA, Tomkins P, Barkwan S. Global impact of mycotoxins on human and animal health management. *Outlook Agric*. 2006;35:247-53. <https://doi.org/10.5367/000000006779398263>
6. Jones FT, Genter MB, Hagler WM, Hansen JA, Mowrey BA, Poore MH, et al. Understanding and coping with effects of mycotoxins in livestock feed and forage. Raleigh, NC: North Carolina Cooperative Extension Service; 1994. Report No.: AG523.
7. Oswald IP, Marin DE, Bouhet S, Pinton P, Taranu I, Accensi F. Immunotoxicological risk of mycotoxins for domestic animals. *Food Addit Contam*. 2005;22:354-60. <https://doi.org/10.1080/02652030500058320>
8. Wild CP, Gong YY. Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis*. 2010;31:71-82. <https://doi.org/10.1093/carcin/bgp264>
9. Maresca M, Fantini J. Some food-associated mycotoxins as potential risk factors in humans predisposed to chronic intestinal inflammatory diseases. *Toxicon*. 2010;56:282-94. <https://doi.org/10.1016/j.toxicon.2010.04.016>
10. Young JC, Subryan LM, Potts D, McLaren ME, Gobran FH. Reduction in levels of deoxynivalenol in contaminated wheat by chemical and physical treatment. *J Agric Food Chem*. 1986;34:461-5. <https://doi.org/10.1021/jf00069a021>
11. Liu YL, Meng GQ, Wang HR, Zhu HL, Hou YQ, Wang WJ, et al. Effect of three mycotoxin adsorbents on growth performance, nutrient retention and meat quality in broilers fed on mould-contaminated feed. *Br Poult Sci*. 2011;52:255-63. <https://doi.org/10.1080/00071668.2011.559453>
12. Aravind KL, Patil VS, Devegowda G, Umakantha B, Ganpule SP. Efficacy of esterified glucomannan to counteract mycotoxicosis in naturally contaminated feed on performance and serum biochemical and hematological parameters in broilers. *Poult Sci*. 2003;82:571-6. <https://doi.org/10.1093/ps/82.4.571>
13. Wu L, Wang W, Yao K, Zhou T, Yin J, Li T, et al. Effects of dietary arginine and glutamine on alleviating the impairment induced by deoxynivalenol stress and immune relevant cytokines in growing pigs. *PLOS ONE*. 2013;8:e69502. <https://doi.org/10.1371/journal.pone.0069502>
14. Hu S, Li X, Rezaei R, Meininger CJ, McNeal CJ, Wu G. Safety of long-term dietary supplementation with L-arginine in pigs. *Amino Acids*. 2015;47:925-36. <https://doi.org/10.1007/s00726-015-1921-5>
15. Yun W, Song M, Lee J, Oh H, An J, Kim G, et al. Arginine addition in a diet for weaning pigs can improve the growth performance under heat stress. *J Anim Sci Technol*. 2020;62:460-7. <https://doi.org/10.5187/jast.2020.62.4.460>
16. Yin J, Ren W, Duan J, Wu L, Chen S, Li T, et al. Dietary arginine supplementation enhances intestinal expression of SLC7A7 and SLC7A1 and ameliorates growth depression in

- mycotoxin-challenged pigs. *Amino Acids*. 2014;46:883-92. <https://doi.org/10.1007/s00726-013-1643-5>
17. Wu G, Knabe DA, Kim SW. Arginine nutrition in neonatal pigs. *J Nutr*. 2004;134:2783S-90S. <https://doi.org/10.1093/jn/134.10.2783S>
 18. Wu L, Liao P, He L, Feng Z, Ren W, Yin J, et al. Dietary L-arginine supplementation protects weanling pigs from deoxynivalenol-induced toxicity. *Toxins*. 2015;7:1341-54. <https://doi.org/10.3390/toxins7041341>
 19. Yao K, Guan S, Li T, Huang R, Wu G, Ruan Z, et al. Dietary L-arginine supplementation enhances intestinal development and expression of vascular endothelial growth factor in weanling piglets. *Br J Nutr*. 2011;105:703-9. <https://doi.org/10.1017/S000711451000365X>
 20. Jobgen W, Meininger CJ, Jobgen SC, Li P, Lee MJ, Smith SB, et al. Dietary l-arginine supplementation reduces white fat gain and enhances skeletal muscle and brown fat masses in diet-induced obese rats. *J Nutr*. 2009;139:230-7. <https://doi.org/10.3945/jn.108.096362>
 21. Wu G, Bazer FW, Davis TA, Kim SW, Li P, Marc Rhoads J, et al. Arginine metabolism and nutrition in growth, health and disease. *Amino Acids*. 2009;37:153-68. <https://doi.org/10.1007/s00726-008-0210-y>
 22. National Research Council. Nutrient requirements of swine. Washington, DC: The National Academies Press; 2012.
 23. Hosseindoust AR, Lee SH, Kim JS, Choi YH, Noh HS, Lee JH, et al. Dietary bacteriophages as an alternative for zinc oxide or organic acids to control diarrhoea and improve the performance of weanling piglets. *Vet Med*. 2017;13:62:53-61. <https://doi.org/10.17221/7/2016-VETMED>
 24. Hosseindoust A, Oh SM, Ko HS, Jeon SM, Ha SH, Jang A, et al. Muscle antioxidant activity and meat quality are altered by supplementation of astaxanthin in broilers exposed to high temperature. *Antioxidants*. 2020;9:1032. <https://doi.org/10.3390/antiox9111032>
 25. Burkhardt B, Pfeiffer E, Metzler M. Absorption and metabolism of the mycotoxins alternariol and alternariol-9-methyl ether in Caco-2 cells in vitro. *Mycotoxin Res*. 2009;25:149-57. <https://doi.org/10.1007/s12550-009-0022-2>
 26. Becker C, Reiter M, Pfaffl MW, Meyer HHD, Bauer J, Meyer KHD. Expression of immune relevant genes in pigs under the influence of low doses of deoxynivalenol (DON). *Mycotoxin Res*. 2011;27:287-93. <https://doi.org/10.1007/s12550-011-0106-7>
 27. Lorenzoni G. Poultry diseases influenced by gastrointestinal health: traditional treatments and innovative solutions. Traditional treatments and innovative solutions. Nottingham: Nottingham University Press; 2010.
 28. Belokrylov GA, Popova OY, Sorochinskaya EI. Detoxication of benzene and aflatoxin B1 with amino acid and peptide preparations in mice and chicken. *Bull Exp Biol Med*. 1999;128:1020-2. <https://doi.org/10.1007/BF02433195>
 29. Mézes M, Balogh K, Tóth K. Preventive and therapeutic methods against the toxic effects of mycotoxins — a review. *Acta Vet Hung*. 2010;58:1-17. <https://doi.org/10.1556/avet.58.2010.1.1>
 30. Yarru LP, Settivari RS, Antoniou E, Ledoux DR, Rottinghaus GE. Toxicological and gene expression analysis of the impact of aflatoxin B1 on hepatic function of male broiler chicks. *Poult Sci*. 2009;88:360-71. <https://doi.org/10.3382/ps.2008-00258>
 31. Trenholm HL, Thompson BK, Foster BC, Charmley LL, Hartin KE, Coppock RW, et al. Effects of feeding diets containing Fusarium (naturally) contaminated wheat or pure deoxynivalenol (DON) in growing pigs. *Can J Anim Sci*. 1994;74:361-9. <https://doi.org/10.4141/cjas94-049>

32. Chaytor AC, Hansen JA, van Heugten E, See MT, Kim SW. Occurrence and decontamination of mycotoxins in swine feed. *Asian-Australas J Anim Sci.* 2011;24:723-38. <https://doi.org/10.5713/ajas.2011.10358>
33. Zang JM, An J. Cytokines, inflammation, and pain. *Int Anesthesiol Clin.* 2007;45:27-37. <https://doi.org/10.1097/AIA.0b013e318034194e>
34. Mehrzad J, Hosseinkhani S, Malvandi AM. Human microglial cells undergo proapoptotic induction and inflammatory activation upon in vitro exposure to a naturally occurring level of aflatoxin B1. *Neuroimmunomodulation.* 2018;25:176-83. <https://doi.org/10.1159/000493528>
35. Escrivá L, Font G, Manyes L, Berrada H. Studies on the presence of mycotoxins in biological samples: an overview. *Toxins.* 2017;9:251. <https://doi.org/10.3390/toxins9080251>
36. Tan B, Yin Y, Kong X, Li P, Li X, Gao H, et al. L-Arginine stimulates proliferation and prevents endotoxin-induced death of intestinal cells. *Amino Acids.* 2010;38:1227-35. <https://doi.org/10.1007/s00726-009-0334-8>
37. Peranzoni E, Marigo I, Dolcetti L, Ugel S, Sonda N, Taschin E, et al. Role of arginine metabolism in immunity and immunopathology. *Immunobiology.* 2008;212:795-812. <https://doi.org/10.1016/j.imbio.2007.09.008>
38. Zhou HR, Yan D, Pestka JJ. Induction of cytokine gene expression in mice after repeated and subchronic oral exposure to vomitoxin (deoxynivalenol): differential toxin-induced hyporesponsiveness and recovery. *Toxicol Appl Pharmacol.* 1998;151:347-58. <https://doi.org/10.1006/taap.1998.8469>
39. Spehlmann ME, Eckmann L. Nuclear factor-kappa B in intestinal protection and destruction. *Curr Opin Gastroenterol.* 2009;25:92-9. <https://doi.org/10.1097/MOG.0b013e328324f857>