

1 **JAST (Journal of Animal Science and Technology)**2 **Upload this completed form to website with submission**

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	The effects of synbiotics-glyconutrients on growth performance, nutrient digestibility, gas emission, meat quality, and fatty acid profile of finishing pigs
Running Title (within 10 words)	Synbiotics-glyconutrients effect on pigs
Author	Olivier Munezero [†] , Sungbo Cho [†] , and In Ho Kim* [†] Olivier Munezero and Sungbo Cho contributed equally
Affiliation	Department of Animal Resource and Science, Dankook University, Cheonan, 31116, Republic of Korea Tel: 82-41-550-3652; Fax: 82-41-550-3652; E-mail: inhokim@dankook.ac.kr
ORCID (for more information, please visit https://orcid.org)	Olivier Munezero: https://orcid.org/0000-0003-4574-0494 In Ho Kim: https://orcid.org/0000-0001-6652-2504 Sungbo Cho: https://orcid.org/0000-0002-2593-2758
Competing interests	There was no conflict of interest disclosed by the authors.
Funding sources State funding sources (grants, funding sources, equipment, and supplies). Include the name and the number of grant if available.	Not applicable.
Acknowledgments	This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. NRF 2022R111A3063332). This work also received support from the Department of Animal Resource & Science through the Research-Focused Department Promotion & Interdisciplinary Convergence Research Projects as a part of the University Innovation Support Program for Dankook University in 2022.
Availability of data and material	If you contact the corresponding author with a rational request, he can provide you with the datasets from this study.

<p>Authors' contributions</p> <p>Please specify the authors' role using this form.</p>	<p>Conceptualization: Olivier Munezero, Sungbo Cho, In Ho Kim</p> <p>Data curation: Olivier M, Sungbo C</p> <p>Formal analysis: Olivier M, Sungbo C</p> <p>Methodology: Olivier M, Sungbo C</p> <p>Software: Olivier M, Sungbo C</p> <p>Validation: Kim IH.</p> <p>Investigation: Olivier M, Sungbo C, Kim IH.</p> <p>Writing - original draft: Olivier M, Sungbo C</p> <p>Writing - review & editing: Olivier M, Sungbo C, Kim IH.</p>
<p>Ethics approval and consent to participate</p>	<p>The experimental procedure was reviewed and accepted by Dankook University's Institutional Animal Care and Use Committee (IACUC) with IACUC #DK-2-2128.</p>

3 CORRESPONDING AUTHOR CONTACT INFORMATION

<p>For the corresponding author (responsible for correspondence, proofreading, and reprints)</p>	<p>Fill in the information in each box below</p>
<p>First name, middle initial, last name</p>	<p>In Ho Kim</p>
<p>Email address – this is where your proofs will be sent</p>	<p>inhokim@dankook.ac.kr</p>
<p>Secondary Email address</p>	
<p>Address</p>	<p>Department of Animal Resource and Science, Dankook University, Cheonan, 31116, Republic of Korea</p>
<p>Cell phone number</p>	<p>82-41-550-3652</p>
<p>Office phone number</p>	
<p>Fax number</p>	<p>82-41-550-3652</p>

4

5

6

7 **Abstract**

8 Glyconutrients help in the body's cell communication. Glyconutrients and synbiotics are promising options for
9 improving immune function. Therefore, we hypothesized that combining synbiotics and glyconutrients will enhance
10 pig nutrient utilization. 150 pigs (Landrace × Yorkshire × Duroc), initially weighing 58.85 ± 3.30 kg of live body weight
11 (BW) were utilized to determine the effects of synbiotics-glyconutrients (SGN) on the pigs' performance, feed
12 efficiency, gas emission, pork traits, and composition of fatty acids. The pigs were matched by BW and sex and chosen
13 at random to 1 of 3 diet treatments: control = Basal diet; TRT1 = Basal diet + SGN 0.15%; TRT2 = Basal diet + SGN
14 0.30%. The trials were conducted in two phases (weeks 1 - 5 and weeks 5 - 10). The average daily gain was increased
15 in pigs fed a basal diet with SGN ($p = 0.036$) in weeks 5-10. However, the apparent total tract digestibility of dry
16 matter, nitrogen, and gross energy did not differ among the treatments ($p > 0.05$). Dietary treatments had no effect on
17 NH_3 , H_2S , methyl mercaptans, acetic acids, and CO_2 emissions ($p > 0.05$). Improvement in drip loss on day 7 ($p = 0.053$)
18 and tendency in the cooking loss were observed ($p = 0.070$) in a group fed basal diets and SGN at 0.30% inclusion
19 level. The group supplemented with 0.30% of SGN had higher levels of palmitoleic acid (C16:1), margaric acid
20 (C17:0), omega-3 fatty acid, omega-6 fatty acid, and ω -6: ω -3 ratio ($p = 0.034, 0.020, 0.025, 0.007, \text{ and } 0.003$,
21 respectively) in the fat of finishing pigs. Furthermore, group supplemented with 0.30% of SGN improved margaric
22 acid (C17:0), linoleic acid (C18:2n6c), arachidic acid (C20:0), omega 6 fatty acid, omega-6 to omega-3 ratio,
23 unsaturated fatty acid, and monounsaturated fatty acid ($p = 0.037, 0.05, 0.0142, 0.036, 0.033, 0.020, \text{ and } 0.045$,
24 respectively) in the lean tissues of finishing pigs compared to pigs fed with the control diets. In conclusion, the
25 combination of probiotics, prebiotics, and glyconutrients led to higher average daily gain, improved the quality of
26 pork, and more favorable fatty acid composition. Therefore, these results contributed to a better understanding of the
27 potential of synbiotic-glyconutrient combinations as a feed additive for pigs.

28

29 **Keywords:** finishing pigs, glyconutrient, performance, prebiotic, probiotic

30

31

32

33 **Introduction**

34 The pig industry faces the challenge of meeting global demand for pork, which is expected to grow by 19% from
35 2019 to 2029 [1]. This must be achieved while also ensuring the production of affordable and high-quality pork that
36 meets consumer preferences and expectations. Pig meat quality and nutritional value are significantly influenced by
37 fatty acid composition in both adipose tissue and muscle [2]. There are numerous ways that fatty acids affect pig's
38 meat quality and nutritional value, including melting point, firmness, flavor, oxidative stability, and shelf life. A
39 high level of saturated fatty acids increases fat melting point and firmness, and a high level of polyunsaturated fatty
40 acids (PUFAs) decreases them [3]. PUFAs also contribute to meat flavor, but they are more prone to oxidation,
41 which can cause rancidity and off-flavors [2]. Nutrient intake is one of the factors that affect fatty acid levels in pig
42 tissues, along with genetics, diet composition, and management practices [4], and this leads to the improvement of
43 pork quality and nutritional value. Moreover, since the feed cost represents 65-75% of the overall production cost in
44 swine production [5, 6], efforts to lower feed costs are a primary concern for boosting the pig industry's
45 competitiveness. Therefore, improving feed efficiency is essential for the profitability of pig production [7, 8]. By
46 improving feed conversion into body weight gain, less feed is required per unit of meat produced, which lowers
47 production costs and increases profits for pig farmers. There are many factors contributing to optimal feed
48 efficiency, including genetics, diet, feed, management, housing, and environment [9]. One of the key factors in
49 improving feed efficiency is maintaining a healthy gastrointestinal tract for optimal metabolic utilization of dietary
50 nutrients. Thus, a healthy gut facilitates or enhances feed digestion and nutrient absorption [7, 10]. Gut health affects
51 nutrient absorption and digestion by influencing the metabolic activity and stability of the gut microbiome, the
52 production and secretion of digestive enzymes, and the function and integrity of the gut immune system [7, 11].

53 The use of synbiotic-glyconutrient combinations as a feed additive has gained popularity in recent years due to their
54 positive effects on gut health and growth performance in other livestock species [12-15]. Glyconutrients and
55 synbiotics are substances that can improve the meat quality of pigs by influencing their gut microbiota and fatty acid
56 composition. According to a study by Núñez-Benítez et al. [12], a standardized mixture of synbiotic-glyconutrients
57 as a feed additive in steers fed a finishing diet improved ruminal fermentation, microbial protein synthesis, and
58 carcass traits. Additionally, a study by Chang et al. [16] found that probiotic-friendly pig production improved meat

59 quality and physicochemical characteristics of pigs by reducing drip loss, cooking loss, shear force, and pH value of
60 pork. Moreover, in lambs, the addition of synbiotics and glyconutrients combination enhanced growth, energy, and
61 carcass weight [13].

62 In our study, we used a synbiotic-glyconutrient mixture composed of probiotics (*L. plantanum*, *B. subtilis*, and *S.*
63 *cerevisiae*), prebiotics (yeast cell wall β -Glucans), and glyconutrients (simple sugars) [12, 17]. *L. plantarum* is a
64 probiotic bacteria known for its beneficial effects on gut health and immunity [18]. *B. subtilis* is another probiotic that
65 has been shown to improve digestive function and reduce pathogenic bacteria in the gut [19]. While *S. cerevisiae* is a
66 yeast commonly used as a probiotic in animal feeds due to its ability to improve nutrient utilization and gut health
67 [20]. Yeast cell wall β -Glucans, a type of prebiotic, act as a source of soluble fiber that feeds the beneficial bacteria in
68 the gut [21]. The glyconutrients in this combination include simple sugars with anti-inflammatory and antimicrobial
69 properties, and are believed to improve cell function and overall health [14]. However, there is inadequate
70 investigation on the effects of this combination on finishing pigs.

71 We expect that the pigs fed the synbiotic-glyconutrient combination will have improved feed efficiency, with higher
72 average daily gain, lower feed conversion ratio, and higher nutrient digestibility compared to the control group. We
73 also anticipate that the treatment will result in a reduction in gas emissions and improvement in meat quality, with a
74 more favorable fatty acid composition. Therefore, the aim of the study was to examine the effects of a standardized
75 synbiotic-glyconutrient combination on growth performance, nutrient digestibility, gas emission, meat quality, and
76 fatty acid profile of finishing pigs.

77 **Materials and Methods**

78 **Animals and ethics**

79 The experimental procedure was reviewed and accepted by the Dankook University's Institutional Animal Care and
80 Use Committee (IACUC) with IACUC #DK-2-2128. The study was carried out at the Dankook University's pig
81 research farm (Gongju, Republic of Korea).

82

83

84 **Animals, diets, and sampling**

85 During 10 weeks of the trial, 150 finishing pigs in total (Landrace × Yorkshire × Duroc) initially weighing 58.85±3.30
86 kg of live body weight (BW) were selected based on BW and sex. Then they were assigned into 30 experimental pens
87 in a completely randomized block design (ten pens for each treatment; with five pigs for each pen; three females and
88 two males per pen). Slatted floors and environmentally controlled rooms were used to house the pigs. Diets based on
89 the corn-soybean meal (Table 1), were created in order to meet or exceed NRC [22] guidelines. The combination of
90 synbiotic and glyconutrient was sourced from Maxcell Global Co. LTD, Seoul, Rep. Korea, containing *L. plantarum*,
91 *B. subtilis*, and *S. cerevisiae* of 1×10^7 CFU/g, with yeast cell wall β -Glucans of 5% from *S. cerevisiae*, and 7% of
92 glyconutrients composed of N-acetylglucosamine, D-xylose, and Fucose. A synbiotic is a combination of probiotics
93 and prebiotics [17]. Probiotics are beneficial bacteria that contribute to a host's health and it's well-being when given
94 sufficient amounts [23]. Prebiotics are non-digestible feed supplements that help promote the multiplication of
95 beneficial microbiota in the gut [24]. Glyconutrients are simple sugars which act as cell integrity promoters, improving
96 health and energy efficiency [14]. Pigs were fed to three different diets (CON, basal diets; TRT1, CON + SGN 0.15%;
97 and TRT2, CON + SGN 0.30%) in phase I and phase II (weeks 1 to 5 and weeks 5 to 10, respectively). During the
98 trial, feed and water were freely available to the pigs.

99 **Samples collection, processing, and calculations**

100 **Growth performance**

101 In order to calculate the average daily gain (ADG), body weight (BW) of every individual pig was recorded on days
102 1, 35, and 70. In addition, the consumed and remained feeds (per pen) were noted for the calculation of average daily
103 feed intake (ADFI), and ultimately the amount of ADG and ADFI were taken to calculate the feed conversion ratio
104 (FCR).

105 **Apparent total tract digestibility (ATTD)**

106 The 2 g/kg of chromium oxide (Cr_2O_3) as indigestible marker were mixed with diets seven days before fecal sample
107 collection in order to determine the ATTD of nitrogen (N), dry matter (DM), and gross energy (GE) of pigs. At the
108 completion of the trial, we have selected one gilt and one barrow per each pen for fresh fecal sample collection. The
109 samples were obtained by massaging a pig's rectum. The samples were directly put into a chilled box. Later on, we
110 transported the samples to the lab, and kept at a temperature of -20°C until they were examined by trained personnel.

111 Following 72 hours of drying at 70°C, samples were finely powdered and sieved through a 1-mm screen. The AOAC
112 [25] methods were applied to assess the digestibility of DM, N, and GE. The analysis of ATTD was conducted using
113 the method utilized in the previous research of Munezero and Kim [26]. A UV absorption spectrophotometric
114 measurement was performed to measure chromium levels (UV-1201, Shimadzu, Kyoto, Japan). A sample of 2 grams
115 of faecal and feed was analyzed using an oxygen bomb calorimeter (Parr 6400 Instrument Company, Moline, Illinois,
116 USA). Moreover, in order to calculate the protein, the N was assessed by using Kjeltec 8600 (Foss Tecator AB,
117 Hoeganaes, Sweden).

118 **Gas emission**

119 At the end of weeks 5 and 10 of the study, fecal samples were collected from one gilt and one barrow per pen to
120 measure the concentration of methyl mercaptans, acetic acid, H₂S, NH₃, and CO₂. A plastic box (2.6 liters) was filled
121 with 300g of collected faeces and covered with adhesive plaster to allow fermentation for 24 hours at 25 degree Celsius.
122 Before measurement, each box with a sample was agitated around 30 seconds to homogenize the sample. The level of
123 gas emission was determined by using a Gastec (Model GV100).

124 **Meat quality**

125 When pigs attained 110 kg of an average weight, they were sacrificed at a nearby abattoir. In order to ensure that the
126 samples were chilled, the carcasses were kept at 2°C for 24 hours. The sample was taken around the ribs positioned at
127 10th to 11th. Before testing, meat samples were put at the temperature of 26 ° C. The color, marbling, and firmness
128 scores were performed at room temperature in accordance with NPPC [27] Standards. The panel had 10 trained
129 personnel who had all been briefed to estimate the sensory features of color, marbling, and firmness. Using a Model
130 CR410 Chromameter, the values of lightness (L*), redness (a*), and yellowness (b*) of each sample were examined
131 immediately after it was cut (Konica Minolta Sensing Inc., Osaka, Japan). Simultaneously, using the pH meter, the
132 pH of each sample were recorded (Model77p, Istek, Seoul, Korea). The water-holding capacity (WHC) was
133 determined using the method described by Kauffman [28]. A meat sample was compressed and the region with
134 moisture was drawn and assessed by the use of digitizing area line sensor (MT10S; M.T. Precision). Then we
135 calculated the WHC values, a ratio with lower value indicates a higher WHC). The area of longissimus muscle (LM)
136 was measured by tracing the LM surface at the 10th rib with the previously mentioned digitizing area line sensor. 3g
137 of meat was sampled to determine the drip loss utilizing the plastic bag technique described by Honikel [29], and cook
138 loss was determined as outlined by Sullivan [30].

139 **Fatty acid profile**

140 Samples were collected to examine the fatty acid composition at the end of the experiment (week 10). In brief, a crude
141 fat extractor was used to collect the sample, which was then placed in a cellulose cup. The sample was then mixed to
142 a 5 ml of n-hexane. After that, BF-3 Methanol (3 ml) and 1 ml of extraction sample were added, homogenized, and
143 reacted at 100°C for 1 hour. Following the reaction, 2 ml of saturated saline and 2 ml of Hexane were added,
144 homogenized, and then purified. A Gas Chromatography-FID (Agilent, Santa Clara, USA) was used to analyze the
145 hexane layer (upper layer) in the distribution solution after 30 minutes.

146 **Statistical analysis**

147 Statistical analysis was performed with General Linear Model procedure of the SAS software (SAS Inst. Inc., Cary,
148 NC, USA) using the Tukey's honest significance test. The pen was taken as the experimental unit. The outcomes from
149 the analysis were illustrated as mean and the standard error of the mean (SEM) values. A p -value < 0.05 was considered
150 significant.

151 **Results**

152 **Growth Performance**

153 Data and results recorded from week 1 to week 10 are shown in Table 2. In weeks 5 to 10, ADG increased ($p=0.036$)
154 as SGN supplementation increased in finishing diets. Finishing pigs supplemented with 0.3% SGN demonstrated
155 higher ADG than other groups. However, the inclusion of SGN in the pigs' diet did not affect BW gain, ADFI, and
156 FCR ($p>0.05$) throughout the experiment.

157 **Apparent total tract digestibility (ATTD)**

158 Table 3 summarizes effects of SGN on ATTD of finishing pigs. SGN supplementation had no effects on ATTD
159 parameters (DM, N, or GE; $p>0.05$).

160 **Gas emissions**

161 Table 4 presents the gas emission results from finishing pigs. The inclusion of SGN to the finishing pig's diets had no
162 significant change on NH_3 , H_2S , methyl mercaptans, acetic acids, and CO_2 emission throughout the trial ($p>0.05$).

163 **Meat quality**

164 Table 5 presents the results of the meat color (lightness, redness, and yellowness), WHC, longissimus muscle area,
165 cooking loss, drip loss, and sensory features. Pigs fed a diet containing 0.3% SGN has indicated a significant lower
166 drip loss than that of other diet-fed pigs on d 7 ($p=0.053$). Moreover, due a diet containing 0.3% SGN, the cooking
167 loss tended to improve ($p=0.070$). Nevertheless, the significant effects of WHC, longissimus muscle area, meat color,
168 and sensory features were not found ($p>0.05$).

169 **Fatty acid profiles in fat of finishing pigs**

170 The effects of SGN supplementation on fatty acid profiles in the fat of finishing pigs are listed in Table 6. The increased
171 levels of palmitoleic acid (C16:1), margaric acid (C17:0), omega-3 fatty acid (ω -3 FA), omega-6 fatty acid (ω -6 FA),
172 and ω -6: ω -3 ratio ($p= 0.034, 0.020, 0.025, 0.007, \text{ and } 0.003$, respectively) were observed in the fat of finishing pigs
173 fed on 0.3% of SGN compared with pigs fed the control diets. Moreover, the SGN tended to increase the concentration
174 of heneicosylic acid (C21:0) and polyunsaturated fatty acid /saturated fatty acid (PUFA/SFA) in the fat of finishing
175 pigs ($p= 0.0813 \text{ and } 0.0877$, respectively).

176 **Fatty acid profiles in the lean tissues of finishing pigs**

177 The impact of SGN addition on the fatty acid profiles in finishing pig lean is illustrated in table 7. SGN tended to
178 increase the concentration of lauric acid (C12:0), palmitic acid (C16:0), and omega 3 fatty acid in the lean of finishing
179 pig ($p=0.094, 0.091, \text{ and } 0.094$ respectively). Furthermore, increased levels of margaric acid (C17:0), linoleic acid
180 (C18:2n6c, LA), arachidic acid (C20:0), omega 6 fatty acid, omega-6 to omega-3 ratio, unsaturated fatty acid, and
181 monounsaturated fatty acid ($p=0.037, 0.052, 0.014, 0.036, 0.033, 0.020, \text{ and } 0.045$, respectively) were observed in the
182 lean of finishing pigs fed diets containing 0.3% of SGN.

183 **Discussion**

184 The actions of this combination are recognized to improve cell communication, which enhances immune responses,
185 mediates inflammation, and reduces cellular stress in general [14]. Our findings indicated that finishing pigs fed SGN-
186 supplemented diets improved ADG in the period of 5-10 weeks. Similarly, an increased growth performance has been
187 observed in synbiotics-glyconutrient supplemented group in poultry and nursery pigs [31, 32]. Possibly, synbiotics

188 and glyconutrients may have improved finishing pig growth by modulating the gut microbiota, enhancing intestinal
189 barrier function, and regulating immune responses. Moreover, glyconutrients can increase energy efficiency and health
190 by aiding in the reduction of inflammation and microbial growth [14, 33]. In addition, Lee et al. [34] and Chu et al.
191 [35] have demonstrated that pigs fed synbiotic-supplemented diets achieved comparable growth rates as pigs fed
192 antibiotic-supplemented diets. In a disease challenge model, Guerra-Ordaz et al. [36] discovered that synbiotics can
193 improve ADG, but not ADFI and Gain: Feed. There is a possibility that the higher ADG observed in a SGN group is
194 due to a numerically higher ADFI in comparison to the control group.

195 A lot of attention has been paid to how probiotics and prebiotics help digest nutrients [37, 38]. This study did not find
196 any significant effect on nutrient digestibility of DM, N, and GE. It could be possible that synbiotics-glyconutrients
197 have a differential effect on the digestibility of different nutrients, such as protein, fat, and fiber, which was not
198 detected by the overall digestibility measurement used in this study. Currently, there is no research on the impact of
199 glyconutrients on nutrient digestibility in pigs. Similarly, a diet supplemented with *Enterococcus faecium* and inulin
200 for growing pigs did not impact the digestibility of DM, GE, and N [39]. Moreover, Weiss et al [40] concluded that
201 *Pediococcus acidilactici* and oligofructose together had no effect on the ileal DM and crude protein digestibility in
202 piglets. In contrast, combining probiotics from bacteria with prebiotics showed higher digestibility of DM and N in
203 weaning pigs [34]. Increased organic matter digestion with probiotic and prebiotic supplementation is largely due to
204 increases in neutral detergent fiber digestion [41]. The differences in our results compared to other research findings
205 could be due to the different breeds and stages of pigs used in these studies.

206 It is thought that changes in gut microbiota affect the composition of feces as well as the amount of gases released
207 from manure. Excreta noxious gas emissions are dependent on intestinal microbial communities and nutrient
208 metabolism [42]. NH₃, H₂S, methyl mercaptans, and acetic acid have not been affected with probiotic and prebiotic
209 inclusion between treatments in the study conducted by Yun et al. [43]. Although Haggins et al. [44] reported that
210 probiotic supplementation could reduce the NH₃ content of broiler excreta, the combination of synbiotic and
211 glyconutrient had no effect on NH₃, H₂S, methyl mercaptans, acetic acids, or CO₂. Perhaps, the microbiome of pigs
212 in the synbiotic-glyconutrient group has not been able to be positively manipulated in a way that can affect noxious
213 gas emissions. Moreover, this could be due to the low sensitivity of the gas measurement method or the high variability

214 of gas production among individual pigs. Furthermore, more accurate and reliable methods of gas emission
215 measurement should be employed to evaluate the environmental impact of synbiotics-glyconutrients in pig diets.

216 The quality of meat and carcass traits determines taste, tenderness, juiciness, and overall consumer acceptance. Our
217 study showed positive outcomes for cooking and drip loss parameters. Synbiotics-glyconutrients may have
218 improved finishing pig pork quality by altering their fatty acid profile and increasing the content of beneficial
219 omega-3 fatty acids, which have positive effects on human health. Consistent with our findings, Zhu et al. [45]
220 found that adding maternal probiotics increased cooking yield and reduced drip loss. Following slaughter, the
221 production of lactic acid through the glycolysis process reduces pH, which is related to drip loss and shear force
222 [46]. According to earlier research, adding *L. plantarum* ZJ316 to piglets' diets increased meat quality by raising the
223 pH_{45min} value [47], and adding *B. coagulans* had a positive impact on meat quality by reducing drip loss [48]. In the
224 study conducted by Zhu et al. [45], maternal synbiotics increased redness while decreasing lightness in the
225 *longissimus thoracis* muscle. Dietary probiotic consumption has been observed to increase redness but not whiteness
226 [49]. These variations may be related to the feeding stage, the type, and the dose of probiotics or synbiotics and
227 components of glyconutrients applied. Therefore, it would be beneficial to conduct further studies to determine the
228 optimum dosage and absorption mechanism of this combination to gain a deeper understanding of its effects on pig
229 meat quality.

230 The fatty acid composition of pork is found to be slightly different from the meat of other animals, such as beef and
231 lamb [50]. The fatty acid profile of pork influences its nutritional value, organoleptic properties, and eating quality
232 [2, 45, 51]. There are several essential fatty acids in pork, including myristic acid, palmitic acid, palmitoleic acid,
233 stearic acid, oleic acid, and linoleic acid [52]. However, fatty acid imbalances can even be harmful to consumers
234 [53]. Pigs store dietary fatty acids in tissues without further modification [54]. Pigs are classified as homolipoid
235 organisms [55], which means that their fatty acid composition closely matches that of their diet. Currently, animal
236 husbandry relies on grain-fed systems, leading to a high intake of omega-6 fatty acids which causes an imbalance
237 between omega-6 and omega-3 [56]. People with this imbalance are more likely to develop cardiovascular disease,
238 inflammation, diabetes, and autoimmune diseases [57]. Furthermore, the increase in linoleic acid could stimulate
239 lipid oxidation in pork, and have a hypocholesterolemic effect and thus slowing the development of atherosclerosis

240 for the consumer [58]. Also, in the body, linoleic acid elongates and desaturates to form C20:4n6, a precursor to pro-
241 inflammatory compounds that can be harmful to health [59]. This prompted a call to rebalance their ratio in the
242 feeds supplied to the animals. Employing nutritional strategies to improve meat fatty acid composition is a beneficial
243 approach. Our study showed that supplementing finishing pigs' diets with the combination of probiotics, prebiotics,
244 and glyconutrients increased the amount of palmitoleic acid (C16:1), margaric acid (C17:0), omega-3 fatty acid,
245 omega-6 fatty acid, and ω -6: ω -3 ratio in fat of the pig meat. In addition, the higher levels of margaric acid (C17:0),
246 linoleic acid (C18:2n6c), arachidic acid (C20:0), omega 6 fatty acid, omega-6 to omega-3 ratio, unsaturated fatty
247 acid, and monounsaturated fatty acid were observed in the lean of finishing pigs. Synbiotics-glyconutrients may
248 have improved fatty acid profiles in pigs due to their prebiotic and probiotic effects, which induce the production of
249 short-chain fatty acids and other metabolites that regulate lipid metabolism. Similarly, probiotics such as
250 *Lactobacillus amylovorus* and *Enterococcus faecium* have also been found to boost the C18:2n6c, monounsaturated
251 fatty acids, and polyunsaturated fatty acids (PUFA) content in pork [54]. Furthermore, our results are constant with
252 a study conducted by Chang et al. [16] who found that omega 3 and 6 fatty acids were significantly higher in the
253 supplemented probiotic group. The addition of probiotics may improve the primary fatty acids content in offspring
254 muscle, resulting in favorable changes in the gut microbiome [60]. Moreover, meat flavor is positively correlated
255 with C16:1 content [61]. The addition of synbiotic-glyconutrient increased the C16:1 level, which suggests that these
256 additives might improve pork flavor. There is insufficient evidence to confirm that dietary prebiotics and probiotic
257 supplementation effectively alter tissue fatty acid profiles. For example, longissimus dorsi muscle fatty acid profile
258 did not change following the addition of inulin and horse chestnuts [62]. However, the inclusion of inulin into rabbit
259 diets has resulted in an increase in linoleic acid and omega-3-PUFA levels, as well as a decrease in the indices of
260 atherogenicity and thrombogenicity [63]. The probiotics, prebiotics, and simple sugars used in this study likely
261 altered the gut microbiota, which could explain the positive results found. Consequently, this led to an increase in
262 the content of beneficial fatty acids in pork.

263 **Conclusions**

264 Results of this experiment indicated that supplementing a diet containing 0.3% SGN improved growth performance,
265 meat quality, and fatty acid profile in both lean and fat tissues. However, the synbiotics-glyconutrients addition had

266 no effect on ATTD and gas emissions of finishing pigs as we expected. The use of synbiotic-glyconutrient
267 combinations as feed additives could lead to improved feed efficiency, higher average daily gain, and improved meat
268 quality. Nevertheless, this study failed to demonstrate an interaction between synbiotics and glyconutrients when this
269 combination was mixed in the pigs' diet. Therefore, our team is developing a robust approach to elucidate more deeply
270 the interaction between synbiotics and glyconutrients and their roles in the health and productivity of livestock.

271

ACCEPTED

272 **References**

- 273 1. Oecd/Fao. OECD-FAO agricultural outlook 2021–2030. OECD Paris, France; 2021.163-77.
- 274 2. Wood J, Enser M, Fisher A, Nute G, Sheard P, Richardson R, et al. Fat deposition, fatty acid composition and
275 meat quality: A review. *Meat Sci.* 2008;78:343-58. <https://doi.org/10.1016/j.meatsci.2007.07.019>.
- 276 3. Wood J, Scollan N. Fatty acids in meat: effects on nutritional value and meat quality. *New Aspects of Meat
277 Quality*: Elsevier; 2022. p. 609-47.
- 278 4. Wood J. Fat deposition and the quality of fat tissue in meat animals. *Fats in animal nutrition.* 1984;407:435.
- 279 5. Singh NM, Singh LA, Kumari LV, Kadirvel G, Patir M. Effect of supplementation of molasses (*Saccharum
280 officinarum*) on growth performance and cortisol profile of growing pig in north eastern hill ecosystem of India.
281 *J. Entomol. Zool. Stud.* 2020;8:302-5.
- 282 6. Munezero O, Kim I. Effect of condensed molasses fermentation solubles (CMS) to replace molasses on the
283 growth performance, nutrient digestibility, and backfat thickness in growing pigs. *Korean J. Agric. Sci.*
284 2022;49:185-92. [10.7744/kjoas.20220015](https://doi.org/10.7744/kjoas.20220015).
- 285 7. Liao SF, Nyachoti M. Using probiotics to improve swine gut health and nutrient utilization. *Anim Nutr.*
286 2017;3:331-43. [10.1016/j.aninu.2017.06.007](https://doi.org/10.1016/j.aninu.2017.06.007).
- 287 8. Munezero O, Muhizi S, Kim IH. Influence of protected organic acids on growth performance, fecal microbial
288 composition, gas emission, and apparent total tract digestibility in growing pigs. *Can J Anim Sci.* 2022;102:554-
289 60. <https://doi.org/10.1139/cjas-2022-0005>.
- 290 9. Renaudeau D, Gilbert H, Noblet J. Effect of climatic environment on feed efficiency in swine. *Feed efficiency
291 in swine*: Springer; 2012. p. 183-210.
- 292 10. Willing BP, Malik G, Van Kessel AG. Nutrition and gut health in swine. *Sustainable swine nutrition.* 2012:197-
293 213. [10.1002/9781118491454](https://doi.org/10.1002/9781118491454).
- 294 11. Zheng L, Duarte ME, Sevarolli Loftus A, Kim SW. Intestinal health of pigs upon weaning: Challenges and
295 nutritional intervention. *Front. Vet. Sci.* 2021;8:628258. <https://doi.org/10.3389/fvets.2021.628258>.
- 296 12. Núñez-Benítez V, Barreras A, Estrada-Angulo A, Castro-Pérez B, Urías-Estrada J, Zinn R, et al. Evaluation of a
297 standardized mixture of synbiotic-glyconutrients as a feed additive in steers fed a finishing diet: Site and extent
298 of digestion, ruminal fermentation, and microbial protein synthesis. *Livest. Sci.* 2021;243:104373.
299 [10.1016/j.livsci.2020.104373](https://doi.org/10.1016/j.livsci.2020.104373).

- 300 13. Castro-Pérez BI, Núñez-Benítez VH, Estrada-Angulo A, Urías-Estrada JD, Gaxiola-Camacho SM, Rodríguez-
301 Gaxiola MA, et al. Evaluation of standardized mixture of synbiotic-glyconutrients supplemented in lambs
302 finished during summer season in tropical environment: growth performance, dietary energetics, and carcass
303 characteristics. *Can J Anim Sci.* 2021;102:155-64. 10.1139/cjas-2020-0202.
- 304 14. Sierpina VS, Murray RK. Glyconutrients: the state of the science and the impact of glycomics. *Explore.*
305 2006;2:488-94. 10.1016/j.explore.2006.08.016.
- 306 15. Valencia GL, Zapata-Ramirez O, Nunez-Gonzalez L, Nunez-Benitez V, LÓPEZ HL, Lopez-Soto M, et al.
307 Effective use of probiotic-glyconutrient combination as an adjuvant to antibiotic therapy for diarrhea in rearing
308 dairy calves. *Turk J Vet Anim Sci.* 2017;41:578-81. 10.3906/vet-1701-54.
- 309 16. Chang SY, Belal SA, Da Rae Kang YIC, Kim YH, Choe HS, Heo JY, et al. Influence of probiotics-friendly pig
310 production on meat quality and physicochemical characteristics. *Korean J Food Sci Anim Resour.* 2018;38:403.
311 10.5851/kosfa.2018.38.2.403.
- 312 17. Markowiak P, Ślizewska K. The role of probiotics, prebiotics and synbiotics in animal nutrition. *Gut Pathog.*
313 2018;10:1-20. 10.1186/s13099-018-0250-0.
- 314 18. Yang S, Deng C, Li Y, Li W, Wu Q, Sun Z, et al. Complete genome sequence of *Lactiplantibacillus plantarum*
315 ST, a potential probiotic strain with antibacterial properties. *J. Anim. Sci. Technol.* 2022;64:183.
316 <https://doi.org/10.5187/jast.2022.e7>.
- 317 19. Kim YJ, Cho SB, Song MH, Lee SI, Hong SM, Yun W, et al. Effects of different *Bacillus licheniformis* and
318 *Bacillus subtilis* ratios on nutrient digestibility, fecal microflora, and gas emissions of growing pigs. *J. Anim. Sci.*
319 *Technol.* 2022;64:291. 10.5187/jast.2022.e12.
- 320 20. Sun H, De Laguna FB, Wang S, Liu F, Shi L, Jiang H, et al. Effect of *Saccharomyces cerevisiae boulardii* on
321 sows' farrowing duration and reproductive performance, and weanling piglets' performance and IgG
322 concentration. *J. Anim. Sci. Technol.* 2022;64:10. <https://doi.org/10.5187/jast.2021.e106>.
- 323 21. De Vries H, Geervliet M, Jansen CA, Rutten VP, Van Hees H, Groothuis N, et al. Impact of yeast-derived β -
324 glucans on the porcine gut microbiota and immune system in early life. *Microorganisms.* 2020;8:1573.
325 <https://doi.org/10.3390/microorganisms8101573>.
- 326 22. NRC. Nutrient requirements of swine. 2012.
- 327 23. Reid G, Jass J, Sebulsky MT, McCormick JK. Potential uses of probiotics in clinical practice. *Clin. Microbiol.*
328 *Rev.* 2003;16:658-72. 10.1128/CMR.16.4.658-672.2003.
- 329 24. Gibson GR, Probert HM, Van Loo J, Rastall RA, Roberfroid MB. Dietary modulation of the human colonic
330 microbiota: updating the concept of prebiotics. *Nutr. Res. Rev.* 2004;17:259-75.
331 <https://doi.org/10.1079/NRR200479>.

- 332 25. AOAC A. Official methods of analysis 16th Ed. Association of official analytical chemists. Washington DC,
333 USA. Sci. Educ. 1995.
- 334 26. Munezero O, Kim I. Effects of protease enzyme supplementation in weanling pigs' diet with different crude
335 protein levels on growth performance and nutrient digestibility. *J. Anim. Sci. Technol.* 2022;64:854-62.
336 10.5187/jast.2022.e51.
- 337 27. NPPC, editor Procedures to evaluate market hogs. Natl.; 1991: Pork Producers Council Des Moines, IA.
- 338 28. Kauffman R, Eikelenboom G, Van der Wal P, Engel B, Zaar M. A comparison of methods to estimate water-
339 holding capacity in post-rigor porcine muscle. *Meat Sci.* 1986;18:307-22. 10.1016/0309-1740(86)90020-3.
- 340 29. Honikel KO. Reference methods for the assessment of physical characteristics of meat. *Meat Sci.* 1998;49:447-
341 57. 10.1016/S0309-1740(98)00034-5.
- 342 30. Sullivan ZM, Honeyman MS, Gibson LR, Prusa KJ. Effects of triticale-based diets on finishing pig performance
343 and pork quality in deep-bedded hoop barns. *Meat Sci.* 2007;76:428-37. 10.1016/j.meatsci.2006.12.002.
- 344 31. Zheng L, Duarte M, Park I, Kim S. Supplemental effects of fermented rice bran extracts on gut health and growth
345 of nursery pigs. *J. Anim. Sci.* 2017;95:109. 10.2527/asasmw.2017.12.228.
- 346 32. Zheng L, Duarte M, Park I, Kim S. Supplemental effects of fermented rice bran extracts on growth performance,
347 bone characteristics, and immune response of broiler chickens. *J. Anim. Sci.* 2017;95:75.
348 10.2527/asasmw.2017.12.158.
- 349 33. Timmerman HM, Mulder L, Everts H, Van Espen D, Van Der Wal E, Klaassen G, et al. Health and growth of
350 veal calves fed milk replacers with or without probiotics. *J. Dairy Sci.* 2005;88:2154-65. 10.3168/jds.S0022-
351 0302(05)72891-5.
- 352 34. Lee SJ, Shin NH, Ok JU, Jung HS, Chu GM, Kim JD, et al. Effects of dietary synbiotics from anaerobic
353 microflora on growth performance, noxious gas emission and fecal pathogenic bacteria population in weaning
354 pigs. *Asian-Australas J Anim Sci.* 2009;22:1202-8. 10.5713/ajas.2009.90045.
- 355 35. Chu GM, Lee SJ, Jeong HS, Lee SS. Efficacy of probiotics from anaerobic microflora with prebiotics on growth
356 performance and noxious gas emission in growing pigs. *Anim Sci J.* 2011;82:282-90. 10.1111/j.1740-
357 0929.2010.00828.x.
- 358 36. Guerra-Ordaz A, González-Ortiz G, La Ragione R, Woodward M, Collins J, Pérez J, et al. Lactulose and
359 *Lactobacillus plantarum*, a potential complementary synbiotic to control postweaning colibacillosis in piglets.
360 *Appl. Environ. Microbiol.* 2014;80:4879-86. 10.1128/AEM.00770-14.

- 361 37. Arowolo MA, He J. Use of probiotics and botanical extracts to improve ruminant production in the tropics: A
362 review. *Anim Nutr.* 2018;4:241-9. 10.1016/j.aninu.2018.04.010.
- 363 38. Lei XJ, Zhang WL, Cheong JY, Lee SI, Kim IH. Effect of antibiotics and synbiotic on growth performance,
364 nutrient digestibility, and faecal microbial shedding in growing-finishing pigs. *J Appl Anim Res.* 2018;46:1202-
365 6. 10.1080/09712119.2018.1484359.
- 366 39. Böhmer B, Branner G, Roth-Maier D. Precaecal and faecal digestibility of inulin (DP 10–12) or an
367 inulin/*Enterococcus faecium* mix and effects on nutrient digestibility and microbial gut flora. *J. Anim. Physiol.*
368 *Anim. Nutr.* 2005;89:388-96. 10.1111/j.1439-0396.2005.00530.x.
- 369 40. Weiss E, Eklund M, Semaskaite A, Urbaityte R, Metzler-Zebeli B, Sauer N, et al. Combinations of feed additives
370 affect ileal fibre digestibility and bacterial numbers in ileal digesta of piglets. *Czech J Anim Sci.* 2013;58:351-9.
371 10.17221/6901-CJAS.
- 372 41. Chaucheyras-Durand F, Walker N, Bach A. Effects of active dry yeasts on the rumen microbial ecosystem: Past,
373 present and future. *Anim. Feed Sci. Technol.* 2008;145:5-26. 10.1016/j.anifeedsci.2007.04.019.
- 374 42. Hakkinen M, Schneitz C. Efficacy of a commercial competitive exclusion product against a chicken pathogenic
375 *Escherichia coli* and *E coli* 0157: H7. *Vet. Rec.* 1996;139:139-41. 10.1136/vr.139.6.139.
- 376 43. Yun W, Lee D, Choi Y, Kim I, Cho J. Effects of supplementation of probiotics and prebiotics on growth
377 performance, nutrient digestibility, organ weight, fecal microbiota, blood profile, and excreta noxious gas
378 emissions in broilers. *J Appl Poult Res.* 2017;26:584-92. 10.3382/japr/pfx033.
- 379 44. Higgins S, Higgins J, Wolfenden A, Henderson S, Torres-Rodriguez A, Tellez G, et al. Evaluation of a
380 *Lactobacillus*-based probiotic culture for the reduction of *Salmonella enteritidis* in neonatal broiler chicks. *Poult.*
381 *Sci.* 2008;87:27-31. 10.3382/ps.2007-00210.
- 382 45. Zhu Q, Song M, Azad MAK, Ma C, Yin Y, Kong X. Probiotics and Synbiotics Addition to Bama Mini-Pigs' Diet
383 Improve Carcass Traits and Meat Quality by Altering Plasma Metabolites and Related Gene Expression of
384 Offspring. *Front. Vet. Sci.* 2022;9. 10.3389/fvets.2022.779745.
- 385 46. Jiang S, Liu Y, Shen Z, Zhou B, Shen QW. Acetylome profiling reveals extensive involvement of lysine
386 acetylation in the conversion of muscle to meat. *J. Proteomics.* 2019;205:103412. 10.1016/j.jprot.2019.103412.
- 387 47. Suo C, Yin Y, Wang X, Lou X, Song D, Wang X, et al. Effects of *Lactobacillus plantarum*ZJ316 on pig growth
388 and pork quality. *BMC Vet. Res.* 2012;8:1-12. 10.1186/1746-6148-8-89.
- 389 48. Zhou X, Wang Y, Li W. Effect of *Lactobacillus* cultures on growth performance, chemical composition, and
390 meal Guangxi of Guangxi yellow chicken. *Poult Sci.* 2010;89:588-93. 10.3382/ps.2009-00319.

- 391 49. Meng Q, Yan L, Ao X, Zhou T, Wang J, Lee J, et al. Influence of probiotics in different energy and nutrient
392 density diets on growth performance, nutrient digestibility, meat quality, and blood characteristics in growing-
393 finishing pigs. *J. Anim. Sci.* 2010;88:3320-6. [10.2527/jas.2009-2308](https://doi.org/10.2527/jas.2009-2308).
- 394 50. Wood J, Enser M. Factors influencing fatty acids in meat and the role of antioxidants in improving meat quality.
395 *Br. J. Nutr.* 1997;78:S49-S60. <https://doi.org/10.1079/BJN19970134>.
- 396 51. Aboagye G, Zappaterra M, Pasini F, Dall'Olio S, Davoli R, Costa LN. Fatty acid composition of the intramuscular
397 fat in the longissimus thoracis muscle of Apulo-Calabrese and crossbreed pigs. *Livest. Sci.* 2020;232:103878.
398 [10.1016/j.livsci.2019.103878](https://doi.org/10.1016/j.livsci.2019.103878).
- 399 52. Zhang Y, Zhang J, Gong H, Cui L, Zhang W, Ma J, et al. Genetic correlation of fatty acid composition with
400 growth, carcass, fat deposition and meat quality traits based on GWAS data in six pig populations. *Meat Sci.*
401 2019;150:47-55. <https://doi.org/10.1016/j.meatsci.2018.12.008>.
- 402 53. Mehta N, Ahlawat S, Sharma D, Dabur R. Novel trends in development of dietary fiber rich meat products—a
403 critical review. *J. Food Sci. Technol.* 2015;52:633-47. [10.1007/s13197-013-1010-2](https://doi.org/10.1007/s13197-013-1010-2).
- 404 54. Ross GR, Van Nieuwenhove CP, González SN. Fatty acid profile of pig meat after probiotic administration. *J.*
405 *Agric. Food Chem.* 2012;60:5974-8. [10.1021/jf205360h](https://doi.org/10.1021/jf205360h).
- 406 55. Shorland F. Effect of the dietary fat on the composition of the depot fats of animals. *Nature.* 1950;165:766-
407 [10.1038/165766a0](https://doi.org/10.1038/165766a0).
- 408 56. Beak S-H, Lee Y, Lee EB, Kim KH, Kim JG, Bok JD, et al. Study on the fatty acid profile of phospholipid and
409 neutral lipid in Hanwoo beef and their relationship to genetic variation. *J. Anim. Sci. Technol.* 2019;61:69.
410 <https://doi.org/10.5187/jast.2019.61.2.69>.
- 411 57. Simopoulos AP. Importance of the omega-6/omega-3 balance in health and disease: evolutionary aspects of diet.
412 *Healthy agriculture, healthy nutrition, healthy people.* 102: Karger Publishers; 2011. p. 10-21.
- 413 58. Wu JH, Lemaitre RN, King IB, Song X, Psaty BM, Siscovick DS, et al. Circulating omega-6 polyunsaturated
414 fatty acids and total and cause-specific mortality: the Cardiovascular Health Study. *Circulation.* 2014;130:1245-
415 [53. 10.1161/CIRCULATIONAHA.114.011590](https://doi.org/10.1161/CIRCULATIONAHA.114.011590).
- 416 59. Naughton SS, Mathai ML, Hryciw DH, McAinch AJ. Linoleic acid and the pathogenesis of obesity.
417 *Prostaglandins Other Lipid Mediat.* 2016;125:90-9. [10.1016/j.prostaglandins.2016.06.003](https://doi.org/10.1016/j.prostaglandins.2016.06.003).
- 418 60. Li X, Cao Z, Yang Y, Chen L, Liu J, Lin Q, et al. Correlation between jejunal microbial diversity and muscle
419 fatty acids deposition in broilers reared at different ambient temperatures. *Sci. Rep.* 2019;9:1-12.
420 [10.1038/s41598-019-47323-0](https://doi.org/10.1038/s41598-019-47323-0).

- 421 61. Cameron N, Enser M, Nute G, Whittington F, Penman J, Fisker A, et al. Genotype with nutrition interaction on
422 fatty acid composition of intramuscular fat and the relationship with flavour of pig meat. *Meat Sci.* 2000;55:187-
423 95. 10.1016/S0309-1740(99)00142-4.
- 424 62. Grela ER, Świątkiewicz M, Florek M, Bąkowski M, Skiba G. Effect of inulin source and a probiotic supplement
425 in pig diets on carcass traits, meat quality and fatty acid composition in finishing pigs. *Animals.* 2021;11:2438.
426 10.3390/ani11082438.
- 427 63. Juárez-Silva ME, Cuchillo-Hilario M, Villarreal-Delgado E. Dietary supplementation of inulin or flavomycin
428 and type of cut of rabbit meat: changes on fatty acid profile and sensorial characteristics. *Rev. Mex. Cienc.*
429 2019;10:552-70. 10.22319/rmcp.v10i3.4714

430

431

432

433

ACCEPTED

434 **Table 1.** Experimental diet ingredient composition (as-fed basis)

Raw Material	phase1	phase2
	%	%
Corn	63.71	68.91
Soybean meal	19.84	11.90
Rapeseed meal	3.00	4.00
DDGS (corn)	5.00	7.00
Tallow	3.40	3.10
Molasses	2.00	2.00
Limestone	1.24	1.27
MDCP	0.53	0.37
Salt	0.30	0.30
DL-Methonine	0.04	-
L-Lysine H ₂ SO ₄	0.41	0.45
L-Threonine	0.06	0.07
L-Tryptophan (10%)	0.17	0.33
Vit/Min premix ¹	0.20	0.20
Phytase	0.05	0.05
Carbohydrase	0.05	0.05
Total	100.00	100.00
Analyzed values		
Moisture	12.90	12.98
CP	16.74	14.41
EE	5.71	5.64
Fiber	2.95	2.89
Ash	5.07	4.72
NSP	120.55	116.40

NDF	10.17	10.80
ADF	2.98	3.09
Ca	0.69	0.66
P	0.42	0.38
Na	0.15	0.16
Cl	0.28	0.28
K	0.83	0.71
Lysine	1.0164	0.8560
Methionine	0.3241	0.2629
Threonine	0.6729	0.5864
Tryptophan	0.1961	0.1771
Met+Cys	0.6204	0.5329

435 ¹Provided per kg diet: Fe, 100 mg as ferrous sulfate; Cu, 17 mg as copper sulfate; Mn, 17 mg as manganese oxide; Zn,
436 100 mg as zinc oxide; I, 0.5 mg as potassium iodide; and Se, 0.3 mg as sodium selenite. Provided per kilograms of
437 diet: vitamin A, 10,800 IU; vitamin D3, 4,000 IU; vitamin E, 40 IU; vitamin K3, 4 mg; vitamin B1, 6 mg; vitamin B2,
438 12 mg; vitamin B6, 6 mg; vitamin B12, 0.05 mg; biotin, 0.2 mg; folic acid, 2 mg; niacin, 50 mg; D-calcium
439 pantothenate, 25 mg;
440 MDCP, monocalcium phosphate; DDGS, dried distillers grains solubles; CP, crude protein; EE, ether extract; NSP,
441 non-starch polysaccharides; ADF, acid-detergent fiber; NDF, neutral-detergent fiber; Met, methionine; Cyst, cystine.

442

443

444

445

446

447

448 **Table 2.** The effect of synbiotic-glyconutrient supplementation on growth performance in finishing pigs¹

Items	CON	TRT1	TRT2	SEM ²	P-Value
BW, kg					
Week 0	53.85	53.85	53.84	0.01	0.807
Week 5	81.12	81.68	81.78	0.43	0.520
Week 10	111.14	113.50	113.95	0.98	0.124
Initial - Week 5					
ADG, g	779	795	798	12	0.512
ADFI, g	2377	2403	2400	22	0.664
FCR	3.054	3.024	3.009	0.030	0.564
Week 5 - Week 10					
ADG, g	858 ^b	909 ^{ab}	919 ^a	16	0.036
ADFI, g	2907	3005	3027	42	0.128
FCR	3.391	3.308	3.295	0.035	0.138
Overall					
ADG, g	819	852	859	14	0.126
ADFI, g	2642	2704	2713	29	0.205
FCR	3.231	3.175	3.162	0.028	0.224

449 ¹Abbreviation: CON, Basal diet; TRT1, Basal diet + SGN 0.15%; TRT2, Basal diet + SGN 0.30%; SGN: synbiotics-
 450 glyconutrients.

451 ²Standard error of means.

452 ^{a,b}Means in the same row with different superscripts differ significantly (P<0.05).

453

454

455

456 **Table 3.** The effect of synbiotic-glyconutrient supplementation on apparent total tract digestibility in finishing pigs¹

Items, %	CON	TRT1	TRT2	SEM ²	P-Value
Week 10					
Dry matter	68.69	69.17	69.54	1.58	0.929
Nitrogen	64.11	65.41	65.88	1.58	0.722
Gross energy	70.41	71.57	71.64	1.51	0.813

457 ¹Abbreviation: CON, Basal diet; TRT1, Basal diet + SGN 0.15%; TRT2, Basal diet + SGN 0.30%. SGN: synbiotics-
458 glyconutrients.

459 ²Standard error of means.

460

ACCEPTED

461 **Table 4.** The effect of synbiotic-glyconutrient supplementation on gas emission in finishing pigs¹

Items, ppm	CON	TRT1	TRT2	SEM ²	P-Value
Week 5					
NH ₃	6.0	5.8	5.8	0.4	0.857
H ₂ S	4.68	5.08	4.65	0.31	0.582
Methyl mercaptans	7.5	7.8	7.5	1.1	0.976
Acetic acid	12.0	12.8	12.0	1.4	0.913
CO ₂	14775.0	14675.0	14525.0	349.6	0.881
Week 10					
NH ₃	7.3	7.3	6.3	0.8	0.564
H ₂ S	6.60	6.53	6.40	0.40	0.940
Methyl mercaptans	7.8	7.5	7.3	0.8	0.901
Acetic acid	12.5	12.3	12.3	1.1	0.983
CO ₂	16675.0	16475.0	16225.0	379.3	0.716

462 ¹Abbreviation: CON, Basal diet; TRT1, Basal diet + SGN 0.15%; TRT2, Basal diet + SGN 0.30%. SGN: synbiotics-
 463 glyconutrients. NH₃, ammonia; H₂S, Hydrogen sulfide; CO₂, carbon dioxide.

464 ²Standard error of means.

465

466 **Table 5.** The effect of synbiotic-glyconutrient supplementation on meat quality in finishing pigs¹

Items	CON	TRT1	TRT2	SEM ²	P-Value
Water holding capacity, %	48.40	49.67	50.94	3.54	0.881
Longissimus muscle area, mm ²	6552.39	6853.80	6861.49	377.80	0.809
Meat color					
L*	51.79	51.13	51.71	0.33	0.345
a*	32.97	32.81	32.68	0.25	0.720
b*	6.16	6.02	6.05	0.13	0.751
Cooking loss, %	32.25	31.86	30.74	0.39	0.070
Drip loss,%					
d1	7.86	7.76	7.50	0.43	0.837
d3	14.04	13.89	12.82	0.50	0.233
d5	19.66	19.51	19.31	0.19	0.463
d7	24.44 ^a	23.76 ^{ab}	24.04 ^b	0.14	0.053
Sensory evaluation					
Color	3.16	3.13	3.31	0.20	0.778
Marbling	3.34	3.34	3.28	0.09	0.845
Firmness	3.31	3.28	3.25	0.10	0.901

467 ¹Abbreviation: CON, Basal diet; TRT1, Basal diet + SGN 0.15%; TRT2, Basal diet + SGN 0.30%. SGN: synbiotics-
 468 glyconutrients. L*, lightness; a*, redness; b*, yellowness; d1, day one; d3, day three; d5, day five; d7, day seven.

469 ²Standard error of means.

470 ^{a,b}Means in the same row with different superscripts differ significantly (P<0.05).

471

472

473

474 **Table 6.** The effect of synbiotic-glyconutrient supplementation on fatty acid profile in finishing pig's fat¹

Items, %	CON	TRT1	TRT2	SEM ²	P-Value
C16:0	20.64	22.97	24.17	1.02	0.238
C16:1	1.75 ^b	2.20 ^{ab}	2.25 ^a	0.11	0.034
C17:0	0.14 ^b	0.47 ^{ab}	0.48 ^a	0.07	0.020
C17:1	0.10	0.18	0.24	0.06	0.406
C18:0	10.85	11.10	14.48	1.12	0.112
C18:1,T	19.00	20.33	20.29	5.81	0.725
C18:1,C	22.96	33.43	44.13	7.85	0.241
C18:2N6T	11.91	16.56	17.46	1.82	0.147
C18:3N3	0.53	0.77	0.91	0.11	0.125
C20:0	0.24	0.25	0.28	0.05	0.834
C20:1	0.86 ^b	0.99 ^b	1.20 ^a	0.10	0.109
C20:2	0.60	0.61	0.61	0.04	0.945
C20:3N6	0.06	0.08	0.09	0.01	0.259
C21:0	0.11	0.17	0.18	0.02	0.087
C20:3N3	0.08	0.09	0.10	0.01	0.454
C22:1N9	0.01	0.03	0.03	0.01	0.298
C23:0	0.05	0.08	0.08	0.01	0.316
ω-3 FA	0.53 ^b	0.85 ^{ab}	0.91 ^a	0.07	0.025
ω-6 FA	11.9 ^b	16.65 ^{ab}	17.57 ^a	0.84	0.007
ω-6: ω-3	18.2 ^b	22.28 ^{ab}	22.74 ^a	0.61	0.003
ΣSFA	40.48	34.42	33.58	3.35	0.347
ΣUSFA	59.51	65.57	66.41	2.53	0.191
ΣMUFA	46.34	47.29	47.40	2.64	0.953
ΣPUFA	13.17	18.27	19.00	1.87	0.133
MUFA/SFA	1.14	1.37	1.41	0.08	0.107

PUFA/SFA	0.32	0.53	0.56	0.07	0.088
TOTAL FATTY ACIDS	100.0	100.00	100.00	0.00	-

475 ¹Abbreviation: CON, Basal diet; TRT1, Basal diet + SGN 0.15%; TRT2, Basal diet + SGN 0.30%. SGN: synbiotics-
476 glyconutrients. Σ SFA: Sum of saturated fatty acids, Σ USFA: Sum of unsaturated fatty acids, Σ MUFA: Sum of
477 monounsaturated fatty acids, Σ PUFA: Sum of polyunsaturated fatty acids, MUFA/SFA: Ratio of monounsaturated
478 fatty acids and saturated fatty acids, PUFA/SFA: Ratio of polyunsaturated fatty acid and saturated fatty acids.

479 ²Standard error of means.

480 ^{a,b}Means in the same row with different superscripts differ significantly ($P < 0.05$).

481

ACCEPTED

482 **Table 7.** The effect of synbiotic-glyconutrient supplementation on fatty acid profile in finishing pig's lean¹

Items, %	CON	TRT1	TRT2	SEM ²	P-Value
C10:0	0.01	0.03	0.03	0.01	0.388
C12:0	0.02	0.04	0.07	0.01	0.094
C14:0	1.41	1.43	1.53	0.04	0.106
C16:0	22.07	22.96	23.05	0.28	0.091
C16:1	3.12	3.13	3.25	0.06	0.303
C17:0	0.01 ^b	0.03 ^{ab}	0.04 ^a	0.01	0.037
C17:1	0.01	0.03	0.03	0.01	0.351
C18:0	10.96	11.28	11.36	0.14	0.189
C18:1,t	41.84	42.05	42.39	0.28	0.435
C18:1,c	3.80	4.18	4.48	0.23	0.198
C18:2N6C, LA	12.44 ^b	13.29 ^{ab}	13.54 ^a	0.26	0.052
C18:3N3, ALA	0.45	0.51	0.60	0.06	0.237
C20:0	0.01 ^b	0.03 ^{ab}	0.04 ^a	0.004	0.014
C20:1	0.69	0.75	0.80	0.06	0.424
C20:2	0.44	0.49	0.53	0.04	0.356
C21:0	0.27	0.37	0.40	0.04	0.158
ω-3 FA	0.45	0.51	0.60	0.04	0.094
ω-6 FA	12.44 ^b	12.96 ^{ab}	13.62 ^a	0.24	0.036
ω-6: ω-3	22.44 ^b	25.59 ^{ab}	28.39 ^a	0.95	0.033
ΣSFA	36.37	35.30	34.95	0.76	0.439
ΣUSFA	63.62 ^b	65.04 ^{ab}	66.69 ^a	0.54	0.020
ΣMUFA	50.16 ^b	51.28 ^{ab}	53.83 ^a	0.81	0.045
ΣPUFA	13.46	14.86	15.86	0.73	0.132
MUPA/SFA	1.37	1.43	1.57	0.09	0.325

PUFA/SFA	0.40	0.42	0.44	0.02	0.306
TOTAL FATTY ACIDS	100.00	100.00	100.00	0.00	-

483 ¹Abbreviation: CON, Basal diet; TRT1, Basal diet + SGN 0.15%; TRT2, Basal diet + SGN 0.30%. SGN: synbiotics-
484 glyconutrients. FA, fatty acid; LA, linoleic acid; ALA, alpha-linolenic acid Σ SFA: Sum of saturated fatty acids,
485 Σ USFA: Sum of unsaturated fatty acids, Σ MUFA: Sum of monounsaturated fatty acids, Σ PUFA: Sum of
486 polyunsaturated fatty acids, MUFA/SFA: Ratio of monounsaturated fatty acids and saturated fatty acids,
487 PUFA/SFA: Ratio of polyunsaturated fatty acid and saturated fatty acids.

488 ²Standard error of means.

489 ^{a,b}Means in the same row with different superscripts differ significantly ($P < 0.05$).

ACCEPTED