

1 TITLE PAGE

ARTICLE INFORMATION	
Article Type	Research article.
Article Title	Effect of <i>Achyranthes japonica</i> extract on growth, digestibility, microbiota, gas emission, and meat quality in broilers fed different protein diets.
Running Title	AJE in different levels of protein diet
Author	Sharif Uddin Khan ¹ , Sung Bo Cho ¹ , In Ho Kim ¹ *
Affiliation	1 Department of Animal Resource and Science, Dankook University, No. 29 Anseodong, Cheonan, Choongnam 330-714, South Korea.
ORCID (for more information, please visit https://orcid.org)	Sharif Uddin khan (https://orcid.org/0000-0001-7664-5057)
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources	Not applicable.
Acknowledgements	Not applicable.
Availability of data and material	Upon reasonable request, the datasets of this study can be available from the corresponding author.
Authors' contributions	<p>Conceptualization: Sharif Uddin khan, Sung Bo Cho, In ho Kim.</p> <p>Data curation: Sharif Uddin khan.</p> <p>Formal analysis: Sharif Uddin khan.</p> <p>Methodology: Sharif Uddin khan, Sung Bo Cho.</p> <p>Software: Sharif Uddin khan.</p> <p>Validation: In ho Kim.</p> <p>Investigation: Sharif Uddin khan, Sung Bo Cho, In ho Kim</p> <p>Writing - original draft: Sharif Uddin khan.</p> <p>Writing - review & editing: Sharif Uddin khan, Sung Bo Cho, In ho Kim.</p>
Ethics approval	This article does not require IRB/IACUC approval because there are no human and animal participants.

2

3 CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	In Ho Kim
Email address – this is where your proofs will be sent	inhokim@dankook.ac.kr
Secondary Email address	
Address	Department of Animal Resource and Science, Dankook University, No. 29 Anseodong, Cheonan, Choongnam 330-714, South Korea.
Cell phone number	82-41-550-3652 / 82-10-8803-9598
Office phone number	
Fax number	+82-41-559-7881

4

5

6

7

8

9

10

11

12

Effect of *Achyranthes japonica* extract on growth, digestibility, microbiota, gas emission, and meat quality in broilers fed different protein diets.

Sharif Uddin Khan^{1†}, Sung Bo Cho^{1†}, In Ho Kim^{1 *}

1 Department of Animal Resource and Science, Dankook University, No. 29 Anseodong, Cheonan, Choongnam 330-714, South Korea.

[†]: These authors are equally contributed.

* Corresponding author: In Ho Kim

Email: inhokim@dankook.ac.kr

Phone: 82-41-550-3652 / 82-10-8803-9598

Fax: +82-41-559-7881

Running head: AJE in different levels of protein diet

Abstract

Achyranthes japonica extract (AJE) is a multifunctional products that express anti-inflammatory, antioxidant and anti-microbial properties. This study was aimed to evaluate the effects of AJE addition to standard and low crude protein diet on growth performance, nutrient digestibility, excreta bacterial count, excreta noxious gas emissions, breast meat quality, and organ weight of broiler chicken. A total of 340 one-day-old Ross 308 broilers [initial body weight of 43.10 ± 1.46 g, 5 replicate cages per treatment, and 17 birds per cage] were randomly distributed into 1 of 4 dietary treatment groups for a 35 day trial. The diets were provided based on three age stage of the broiler. In the starter stage broiler were fed basal diet. Experimental diet were fed to broiler from day 8 to 35. In growing (days 8-21) and finishing (days 22-35) stage broiler were fed: Standard crude protein (SCP) diet and low crude protein (LCP) diet with 0.025% and 0.05% of AJE supplementation respectively. Here, the SCP and LCP diets were 21.50% and 20.86% CP during days 8–21 and 20.00% and 19.40% CP during days 22–35, respectively. The SCP diets with 0.025% AJE supplementation resulted in higher ($P<0.5$) body weight gain (BWG) at finishing stage and a tendency to lower feed conversion ratio (FCR) and body weight gain in the overall period compared to LCP diets with or without AJE supplementation. Moreover, dry matter and nitrogen digestibility were increased with SCP diet along with 0.025% of AJE. No significant difference was found in meat quality parameters except for pH. Interestingly, the NH_3 gas emission to the environment was found to be less with different levels of CP and AJE supplementation. Therefore, we concluded that the addition of 0.025% AJE to the SCP diet improved broiler growth performance and nutrient digestibility with low fecal NH_3 emissions.

Keywords: *Achyranthes japonica*, broiler, growth performance, low and high protein diet

Introduction

The population of the world is rapidly increasing, especially in developing countries. For their better growth and development, proteins are very essential elements. The broiler is a good source of high-quality protein. A significant portion of global protein demand is met by the broiler industry [1]. Nutritionally balanced feed is more efficient for better broiler production. Feed cost accounts for 70% of the total production expenditure in the poultry industry [2]. Protein is the second-most vital and expensive nutrient component in ration formulation whose price are increasing day by day. Soybean meals are widely used in poultry's diet [3] because they are a preferred source of protein, which contains well-balanced amino acids that are highly digestible. Reducing dietary protein while maintaining essential amino acid balance can be an effective technique to minimize feed price and nitrogen release into the environment [4].

Since 2006, the use of antibiotics in livestock industry was banned by european union due to residual effect on food [5]. After that, researchers are paying attention to alternative growth promoters like probiotics, prebiotics, herbal products, natural marine products, and organic acids [6]. Phytogetic feed additives are one of them. *Achyranthes japonica* extract (AJE) is a phytogetic feed additive belongs to the family *Amaranthaceae*. AJE is mostly available in Asian countries, especially in Japan, China, and South Korea [7]. AJE contains a wide range of bioactive compounds, including polyphenols, flavonoids, and saponins. These compounds have a variety of health benefits, such as anti-inflammatory, antioxidant, and anti-microbial effects [8,9]. In broilers, supplementation of AJE with low crude protein (LCP) diet showed higher growth performance by improving nitrogen digestibility and controlling the bacterial count in the gut [2,10]. Previous research on AJE addition with different crude protein levels is quite limited. Therefore, we hypothesized that the synergistic effect of AJE on nutrient digestibility and microbial population in broilers could help recover their production performance when supplemented with LCP diet. Our objective was to evaluate the impact of AJE supplementation on growth performance, nutrient digestibility, excreta bacterial count, and gas emission in broilers fed different protein diets.

Materials and methods

Experimental site and Ethical considerations

Experimental research was conducted at the experimental farm of Dankook University, Jeonui (Sejong, Republic of Korea). All the research procedures and animal care activities were followed in accordance with the care and use of experimental animals. The experiment was approved by the Animal Care Committee of Dankook University in South Korea (No. DK-1-1939).

Additives information with processing method

For this study, we commercially purchased AJE from Synergen, Inc. In brief, the roots of *Achyranthes Japonica* were washed more than three times in clean water and dried at 80 °C. The dried roots were then milled using an IKA M20 machine (Gemini BV company). Milled specimens were extracted with distilled water for approximately 4 hours to obtain an initial extract, and then it was filtered by a high-velocity centrifugal machine. Then the useful parts were taken and eluted with an ethanol solution, and the extracts were let cool completely before being dried in a spray dryer. Finally, 1.15 mg/g of flavonoids, 4.26 mg/g of total polyphenols, and 0.47 mg/g of saponin was obtained from the AJE.

Animals, diets, and experimental design

A total of 340 one-day-old Ross 308 broiler chicks (43.10 ± 1.46 g) were randomly assigned to one of four experimental treatment groups. There were five replicate cages per treatment, including 17 birds per cage. The diets were provided based on three age stage of the broiler. In starter stage (days 1-8), broiler were fed basal (Control) diet. Experimental diet were provided to broiler from day 8 to 35 of age. In growing (days 8-21) and finishing (days 22-35) stage broiler were fed: Standard crude protein (SCP) diet and LCP diet with/without 0.025% and 0.050% AJE respectively. During the LCP diet formulation, 3% of the CP level was reduced based on the CP level of the SCP diet. CP levels in SCP and LCP were 21.5% and 20.86, respectively, during the growing stage. In the finishing stage, SCP and LCP diets contained 20% and 19.4%, respectively. A double amount of AJE was added to the LCP diet compared to the SCP diet as a recovery source for reduced CP levels. AJE was incorporated into the diet by replacing the same amount of corn. All the experimental diets (Table 1) were formulated following Aviagen's 2019 nutrient requirements for poultry. The broiler were housed in an environmentally controlled shed.

The battery cage system ($1.75 \times 1.55 \text{ m}^2$) is followed by three stages of a stainless-steel cage. Approximately 33 °C temperature was maintained during the brooding period, and 1 °C was reduced each week until 22 °C was reached. The cage was well arranged with attached feeders and a nipple drinker for ad libitum feed and water consumption. Fluorescent lights were used as a source of artificial light for 24 h per a day.

Growth performance

Body weight (BW) was recorded (340 birds) on days 0, 8, 22, and 35 in each pen, as along with feed intake (FI). Body weight gain (BWG) and Feed conversion ratio (FCR) was calculated by the using of recorded data during the experiment.

Apparent nutrient digestibility

In total, 0.2% indigestible marker chromium oxide (Cr_2O_3) was added to the broilers' diets at the 28th day of age to measure the apparent total tract digestibility of dry matter, nitrogen, and metabolic energy. After 5 days of the supplemented diet, an excreta sample (2 sample/cage, 10 samples/treatment) was collected at days 35. The plastic plates were placed on the pen's floors for two continuous days, and 400 grams of excreta were taken from each pen. Until further chemical analysis was performed, samples were stored at -20°C. The excreta samples were cooled to 0°C and de-moisturized at 70°C for three days to perform a chemical analysis. The samples were then finely crushed and put through a 1-mm screen to filter. Following the guidelines established by the Association of Official Analytical Chemists (AOAC), samples of both feed and excreta were examined. Dry matter and nitrogen were analyzed using the 934.01 (AOAC) and 968.0 (AOAC) methods, respectively. During nitrogen analysis, a Kjeltec 2300 nitrogen Analyzer (Foss Tecator AB, Hoeganaes, Sweden) was used. Through the use of an automated adiabatic oxygen bomb calorimeter, the gross energy (GE) has been determined (Parr 6300 Calorimeter, Moline, IL). To measure the concentration of chromium, an UV absorption spectrophotometer (UV-1201, Shimadzu Japan) was used. The apparent total tract digestibility (ATTD) of a nutrient was calculated using the following equation:

$$\text{ATTD (\%)} = (1 - ((\text{Nf} \times \text{Cd}) / (\text{Nd} \times \text{Cf})) \times 100$$
, where Nf = concentration of nutrient in feces (% DM), Nd = concentration of nutrient in diet (% DM), Cf = concentration of chromium in feces (% DM), and Cd = concentration of chromium in diet (% DM).

Excreta bacterial count

After taking the final body weight at day 35 of age, fresh excreta samples were collected from each pen (2 samples/cage). For feces collection, randomly selected five birds from each cage were placed in an empty separate cage with clean plastic plates for 2 hours. The 10 g of fresh mixed samples were collected from each plate. Immediately, the collected samples were sent to the laboratory for analysis. Five samples from treatment were taken, each of which contained 1 g of mixed excreta with 9 mL of 1% peptone broth (Becton, Dickinson and Company, USA) and was vortexed. These "homogenized specimens" were mixed to a tenfold serial dilution with a 1% peptone broth solution and put on MacConkey agar plates (Difco Laboratories, MI, USA), Lactobacilli medium agar plates (Medium 638; DSMZ, Braunschweig, Germany), and *Salmonella Shigella* agar plates (Becton, Dickinson and Company, USA) to incubate *E. coli*, *Salmonella*, and *Lactobacillus*, respectively. The *Lactobacillus* bacterial count was incubated for 48 hours at 39 °C under anaerobic conditions. *Salmonella Shigella* and MacConkey agar plates were incubated for 24 hours at 37 °C. Immediately, bacterial colonies were counted after being taken out of the incubator.

Excreta gas emission

Finally, on day 35, grinded excreta samples (2 samples/cage) were collected. The collected samples were kept in 2.6-L plastic boxes, where there was a hole in the center of the box side wall that was airtight with tape. The samples were fermented at 28 °C for five days. A multi-gas detector meter (Model PGM-6208, RAE system, USA) was used to measure NH₃, H₂S, and methyl mercaptans on the first, third, and fifth days of fermentation. The average values of three days of measurements were recorded.

Statistical analysis

All data were statistically analyzed as a randomized complete block design using GLM procedure of SAS software [11]. Preplanned contrast were performed to test the effects of different level of CP and AJE. Data was presented as the mean and standard errors of means. The Cage was considered as an experimental unit. Statistical significance was taken consideration when the *p*-value was remained under 0.05., and when it was less than 0.10, trends were taken into account.

Results

Growth performance

The effect of AJE supplementation with different protein diets on broiler growth performance is summarized in Table 2. The SCP diet with (0.025%) AJE supplementation resulted in significantly higher BWG ($P \leq 0.05$) at the finishing stage and a tendency (0.0911, 0.0980) to lower FCR and higher body weight gain at the overall period compared to the LCP diet with or without (0.050%) AJE supplementation. In addition, throughout the experimental period, FI remained unaffected among all treatment groups.

Apparent nutrient digestibility

The effects of different levels of protein diet with/without AJE supplementation on broiler apparent nutrient digestibility are shown in Table 3. The apparent total tract digestibility of dry matter and nitrogen shows significantly improved results with the SCP diet along with 0.025% AJE supplementation compared to the LCP diet with/without AJE supplementation. Interestingly, metabolizable energy digestibility remained unaffected ($p > 0.05$) throughout the experimental period in all the treatment groups.

Excreta bacterial count

The effect of different protein diets with/without AJE supplementation on the cecal bacterial count in broiler feces is shown in Table 4. The bacterial populations of *Lactobacillus* increased and *E. coli* decreased significantly ($p < 0.05$), respectively, as a result of 0.025% AJE supplementation along with SCP diet compared to LCP diet with/without AJE supplementation. Dietary crude protein and AJE supplementation levels both failed to have an effect on the *salmonella* population in broiler excreta samples throughout the experimental period.

Excreta gas emission

Table 5 shows the effect of different protein diets with/without AJE supplementation on broiler excreta gas emissions. However, all the mentioned gases were not influenced by different levels of protein in the diets with AJE supplementation, except NH_3 , throughout the experimental period.

Meat quality and visceral percentage

The meat quality and visceral percentage of broilers are summarized in Table 6. The dietary supplementation of AJE along with different levels of protein had no significant improvement ($p > 0.05$) on meat quality and visceral percentage. Surprisingly, the pH parameter shows a significant ($p < 0.05$) effect with different level of AJE compared to SCP diet without supplementation.

Discussion

AJE has several physiological and biological properties, like anti-oxidant, anti-inflammatory, and bioactive components, which have attracted the attention of researchers who are working on it as an alternative source of antibiotic growth promoters. There are limited studies performed in broilers with AJE supplementation, as it was recently introduced to the feed industry. Therefore, our main focus was to evaluate the effects of AJE with different levels of protein in diets as a natural growth promoter. Hoque and kim, [2] reported that the LCP and HCP diets with 0.025% and 0.05% of AJE significantly increased BWG compared to the LCP diet without AJE. Similarly, Park and kim, Sun et al. [12,13] in broilers, and [14] in ducks observed a linear increase in BWG and reduced FCR through the supplementation of *Achyranthes bidentata* extract. This improvement in BWG can be caused by several mechanisms since AJE supplementation has beneficial effects on the gut health of animals [15]. Hashemi and Davoodi [16] proposed that significant body weight gain was caused by increased digestibility, an improved intestinal microbial population, and enhanced digestive enzyme activities. This proposed mechanism was supported by our study as supplementation of AJE increased nitrogen and dry matter digestibility and created a favorable environment for the microorganism. As a result, BWG significantly increased at the finishing stage, though BWG and FCR tend to increase and decrease with (0.025%) AJE supplementation, respectively, at the overall period. On the other hand, reducing the CP level in diet had negative effects on broiler BWG and FCR in our study. Similarly, Liu and Kim, [17] in pig and [2] found the similar results in broiler. The negative effect of reducing the level of crude protein in the broiler diet can be minimized by the addition of essential amino acids [18]. Previous researchers explained that the LCP diet impairs the growth performance and gastrointestinal conditions in broilers [19]. However, in this experiment, supplementation of AJE in broiler diet failed to improve growth performance in LCP diet compared to SCP diet, although AJE showed improved growth performance with high protein levels. Regarding the negative effect of the LCP diet, [18] showed that reduced growth performance in broilers can be minimized by the addition of essential amino acids to the broiler diet. However, we request further study to find out the optimum AJE level in the LCP diet to show similar performance as a high protein diet.

218

219 In our study, the ATTD of DM and nitrogen digestibility were significantly improved with 0.025% of AJE
220 supplementation. Similar findings were stated by [2,12,13]. Previous study showed that, plant extract
221 supplementation has been shown to increase villi height and thus may improve nutrient absorption ability in
222 broilers and quail [17]. Furthermore, the results from the other studies suggest that feeding plant extract can
223 increase digestibility by stimulating bile secretion and enhancing the digestive enzyme of the small intestine [12].

224

225 The intestinal tract has vital roles in digestion and the absorption of nutrients. It has some specific functions and
226 acts as a harbor for a wide range of microorganisms [20]. As a result, the increasing population of beneficial
227 bacteria can help in the utilization of unprocessed nutrients and maintain the gut microbiota [2]. In the present
228 study, AJE supplementation (0.025%) with a SCP diet increased *lactobacillus* and reduced *E. coli* populations in
229 the broiler's excreta sample. Similarly, Lan et al., Sub et al. [21,13], and [2] found similar results. AJE
230 supplementation creates a favorable condition in the digestive system of broilers that increases *lactobacillus*
231 counts, although the mechanism is not yet understood. A large number of such beneficial bacteria inhibit the
232 harmful bacteria in the animal gut [22] due to the competition for nutrients. A small amount of nutrients is left by
233 the *lactobacillus* for the utilization and multiplication of *E. coli* bacteria. The anti-inflammatory and antioxidant
234 properties of AJE could develop the normal gut microbiota balance by suppressing the pathogenic bacteria and
235 lead to healthy intestinal development [12]. It is expected that a lower level of dietary protein will reduce the
236 bacterial population in feces [23], but in this study, we got the reverse result. Similarly, Liu and Kim, [17] in pigs
237 and [2] in broilers did not find an effect of protein levels on the microbial population. The mechanism of the effect
238 of protein level on microbial count is yet not well known. The different sources of protein may be the possible
239 reason for this result. Further study is required to determine the effect of protein in the microbial population.

240

241 Noxious gas emissions from animals in the environment are alarming issues due to climate change and negative
242 effect on animal performance. Excreta noxious gas emissions of animals are related to the intestinal microflora,
243 particularly the harmful bacterial population [24]. In this study, supplementation with 0.025% and 0.05% of AJE
244 reduced the NH₃ emission in the excreta sample compare to SCP diet without supplementations. Similarly, Park
245 and Kim [12] and [2] stated similar results in the broiler. Additionally, [25] found a lower amount of H₂S and NH₃

gas in the environment through pig fecal. Mainly, ammonia gas is produced from the indigestible dietary protein released in feces and urine [2]. The AJE supplemented diet led the way to better microbiota balance and higher nutrient digestibility compared to the control diets in this study. Due to the increase in nitrogen digestibility and lactobacillus count, we conjectured that ammonia gas emissions may be reduced.

Good quality of meat takes the attention of consumers due to its texture, color, flavor, and long shelf life. The pH level of meat is measured by the amount of lactic acid present in the muscle tissue. The effects of plant extract on carcass quality are ambiguous. Our results suggested that AJE (0.025% and 0.050%) supplementation reduces the pH level compared to the control diet along with SCP and LCP, which has an overall effect on meat quality. Dang et al. [26] reported that supplementation of AJE to the diet had no positive effect on carcass quality. Interestingly, in our study, we found a lower pH level with AJE supplementation. We found lower pH may be due to a reduction in bacterial population caused by AJE supplementation. In addition, by using AJE supplementation, broiler meat texture, color, flavor, and shelf life increased compared to the control diet.

Conclusion

Supplementation of 0.025% AJE with SCP diet improved the growth performance at finishing and overall experimental period and increased the DM and nitrogen digestibility in broilers. Reduced ammonia gas emissions and improved meat quality by keeping pH at a moderate level. However, further study is required to determine the optimum dose of AJE to recover the negative effect of LCP diet.

- 268 1. Khan, S.U., Fouzder, S.K. and Sarkar, P.K. 2023. Productivity and profitability of commercial broiler
269 chickens under various farming conditions. *Journal of Experimental Biology and Agricultural Sciences*. 11,
270 1 (Feb. 2023), 209–215. DOI:[https://doi.org/10.18006/2023.11\(1\).209.215](https://doi.org/10.18006/2023.11(1).209.215).
- 271 2. Hoque M, Kim I. H. 2022. Effect of *Achyranthes japonica* Naki extract on growth performance, apparent
272 nutrition digestibility, excreta microbial count and gas emission in broiler fed different protein diets". *Journal*
273 *of animal and feed science*, 32(1),34-42. doi:10.22358/jafs/155126/2022.
- 274 3. Munezero, O., Z. F. Zhang, and I. H. Kim. "The Effects of *Yucca Schidigera* Extract and Multi-carbohydrase
275 in Different Crude Protein Diets on Growth Performance, Nutrient Digestibility, Carcass Parameters and
276 Excreta Noxious Gas Contents in Broilers." *Brazilian Journal of Poultry Science* 25 (2022).
- 277 4. Saleh A.A., Amber K.A., Soliman M.M., Soliman M.Y., Morsy W.A., Shukry M., Alzawqari M.H., 2021.
278 Effect of Low Protein Diets with Amino Acids Supplementation on Growth Performance, Carcass Traits,
279 Blood Parameters and Muscle Amino Acids Profile in Broiler Chickens under High Ambient Temperature.
280 *Agriculture*. 11, 185
- 281 5. Janardhana, V., Broadway, M. M., Bruce, M. P., Lowenthal, J. W., Geier, M. S., Hughes, R. J., Bean, A. G.
282 D. (2009). Prebiotics Modulate Immune Responses in the Gut-Associated Lymphoid Tissue of Chickens.
283 *The Journal of Nutrition*,139(7):1404-9.
- 284 6. Ayasan, T, 2013. Effects of dietary inclusion of protexin (probiotic) on hatchability of Japanese quails. *Indian*
285 *J. Anim.Sci.*83(78–81).
- 286 7. Jung, S. M., Choi, S. I., Park, S. M., & Heo, T. R. (2007). Antimicrobial effect of *Achyranthes japonica*
287 Nakai extracts against *Clostridium difficile*. *Korean Journal of Food Science and Technology*, 39, 564–568.
- 288 8. Choi, H. A., D. E. Cheong, H. D. Lim, W. H. Kim, M. H. Ham, M. H. Oh, Y. Wu, H. J. Shin, and G. J. Kim.
289 2017. Antimicrobial and anti-biofilm activities of the methanol extracts of medicinal plants against dental
290 pathogens *Streptococcus mutans* and *Candida albicans*. *J. Microbiol. Biotechnol.* 27:1242–1248
- 291 9. Upadhaya, S. D., and I. H. Kim. 2017. Efficacy of phytogetic feed additive on performance, production and
292 health status of mono-gastric animals—a review. *Ann. Anim. Sci.* 17:929–948.
- 293 10. Md Mortuza Hossain, Sung Bo Cho, and In Ho Kim. 2022. Effects of adding graded levels of *Achyranthes*
294 *japonica* root extract to low crude protein diet on growth performance, nutrient digestibility, fecal microbiota,
295 and meat quality parameters in broilers. *Canadian Journal of Animal Science*. 103(1): 26-32.
296 <https://doi.org/10.1139/cjas-2022-0092>
- 297 11. SAS [Statistical Analysis System]. SAS software for PC. Release 9.3. Cary, NC: SAS Institute; 2012.
- 298 12. J.H. Park, I.H. Kim, 2020. Effects of dietary *Achyranthes japonica* extract supplementation on the growth
299 performance, total tract digestibility, cecal microflora, excreta noxious gas emission, and meat quality of
300 broiler chickens, *Poultry Science*, Volume 99, Issue 1, 2020, Pages 463-470, ISSN 0032-5791,
301 <https://doi.org/10.3382/ps/pez533>.

- 302 13. Sun H.Y., Kim Y.M., Kim I.H., 2020. Evaluation of *Achyranthes japonica* Nakai extract on growth
303 performance, nutrient utilization, cecal microbiota, excreta noxious gas emission, and meat quality in broilers
304 fed corn-wheat-soybean meal diet. *Poult. Sci.* 99, 5728–5735, <https://doi.org/10.1016/j.psj.2020.07.023>
- 305 14. Ao X., Kim I.H., 2020. Effects of *Achyranthes bidentata* polysaccharides on performance, immunity,
306 antioxidant capacity, and meat quality in Pekin ducks. *Poult. Sci.* 99, 4884–4891,
307 <https://doi.org/10.1016/j.psj.2020.06.026>
- 308 15. Liu, Z., X. Wang, S. Ou, M. Arowolo, D. X. Hou, and J. He. 2018. Effects of *Achyranthes bidentata*
309 polysaccharides on intestinal morphology, immune response, and gut microbiome in yellow broiler chickens
310 challenged with *Escherichia coli* K88. *Polymers* 10:1233-1247.
- 311 16. Hashemi S.R., Davoodi H., 2010. Phytochemicals as new class of feed additive in poultry industry. *J. Anim. Vet.*
312 *Adv.* 9, 2295–2304, <https://doi.org/10.3923/javaa.2010.2295.2304>
- 313 17. Liu X., Kim I.H., 2021. Effects of long-term feeding of *Achyranthes japonica* Nakai extract as a supplement
314 to diets with different protein levels diets on the growth performance and meat quality characteristics of
315 growing-fattening pigs. *Anim. Feed Sci. Technol.* 279, 115030, [https://doi.org/10.1016/j.anifeed-](https://doi.org/10.1016/j.anifeed-sci.2021.115030)
316 [sci.2021.115030](https://doi.org/10.1016/j.anifeed-sci.2021.115030)
- 317 18. Peng X., Hu L., Liu Y. et al., 2016. Effects of low - protein diets supplemented with indispensable amino
318 acids on growth performance, intestinal morphology and immunological parameters in 13 to 35 kg pigs.
319 *Animal* 10, 1812–1820, <https://doi.org/10.1017/S1751731116000999>
- 320 19. Dozier, W.A., Moran, E.T., Jr., and Kidd, W.T. 2000. Response of fast- and slow-feathering male broilers to
321 dietary threonine during 42 to 56 days of age. *J. Appl. Poult. Res.* 9: 460–467. doi:10.1093/japr/9.4.460
- 322 20. Yan L., Meng Q.W., Kim I.H., 2011. The effect of an herb extract mixture on growth performance, nutrient
323 digestibility, blood characteristics and fecal noxious gas content in growing pigs. *Livest. Sci.* 141, 143–147,
324 <https://doi.org/10.1016/j.livsci.2011.05.011>
- 325 21. Lan R.X., Lee S.I., Kim I.H., 2016. Effects of multistrain probiotics on growth performance, nutrient
326 digestibility, blood profiles, faecal microbial shedding, faecal score and noxious gas emission in weaning
327 pigs. *J. Anim. Physiol. Anim. Nutr.* 100, 1130–1138, <https://doi.org/10.1111/jpn.12501>
328 <https://doi.org/10.1111/jpn.12501>
- 329 22. Jang G.Y., Kim H.Y., Lee S.H., Kang Y., Hwang I.G., Woo K.S., Kang T.S., Lee J.S., Jeong H.S., 2012.
330 Effects of heat treatment and extraction method on antioxidant activity of several medicinal plants. *J. Korean*
331 *Soc. Food Sci. Nutr.* 41, 914–920, <https://doi.org/10.3746/jkfn.2012.41.7.914>
- 332 23. Zhou L., Fang L., Sun Y., Su Y., Zhu W., 2016. Effects of the dietary protein level on the microbial
333 composition and metabolomic profile in the hindgut of the pig. *Anaerobe* 38, 61–69,
334 <https://doi.org/10.1016/j.anaerobe.2015.12.009>
- 335 24. Ferket, P. R., E. van Heugten, T. A. T. G. van Kempen, and R. Angel. 2002. Nutritional strategies to reduce
336 environmental emissions from nonruminants. *J. Anim. Sci.* 80 (E. Suppl. 2): E168–E182.

- 337 25. Dang D.X., Kim Y.M., Kim I.H., 2020. Effects of a root extract from *Achyranthes japonica* Nakai on the
338 growth performance, blood profile, fecal microbial community, fecal gas emission, and meat quality of
339 finishing pigs. *Livest. Sci.* 239, 104160, <https://doi.org/10.1016/j.livsci.2020.104160>
- 340 26. Dang D.X., Yun K.S., Kim I.H. 2021. Effects of broiler chicks fed the low crude protein diet supplemented
341 with *Achyranthes Japonica* Nakai root extract on growth performance and carcass traits. *Can. J. Anim. Sci.*
- 342 27. Aviagen 2019. Ross Nutrition Specifications. [Accessed January 2020]

343

ACCEPTED

344 **Table**

345

346 Table 1. Ingredients and chemical composition of basal diets (as-fed basis)

Ingredients (%)	Starter	Grower		Finisher	
	SCP	SCP	LCP (-3% CP)	SCP	LCP (-3% CP)
Corn	43.63	47.45	49.07	53.78	55.31
Soybean meal	35.08	31.28	29.65	28.18	26.63
Corn gluten meal	13.00	13.00	13.00	10.00	10.00
Wheat bran	3.00	3.00	3.00	3.00	3.00
Soybean oil	1.76	1.74	1.74	1.51	1.51
TCP	1.81	1.81	1.85	1.81	1.86
Limestone	0.94	0.94	0.91	0.94	0.91
Salt	0.36	0.36	0.36	0.36	0.36
Methionine (99%)	0.19	0.19	0.17	0.19	0.17
Lysine	0.03	0.03	0.05	0.03	0.05
Mineral mix ¹	0.10	0.10	0.10	0.10	0.10
Vitamin mix ²	0.10	0.10	0.10	0.10	0.10
Total	100.00	100.00	100.00	100.00	100.00
Analyzed value					
Crude protein, %	23.00	21.50	20.86	20.00	19.40
Ca, %	1.10	1.08	1.08	1.07	1.07
P, %	0.83	0.82	0.82	0.79	0.79
Available P, %	0.54	0.53	0.53	0.52	0.52
Lys, %	1.26	1.15	1.12	1.06	1.03
Met, %	0.54	0.52	0.50	0.50	0.48
ME, kcal/kg	3200	3200	3200	3200	3200
FAT, %	4.45	4.51	4.54	4.32	4.35
Fiber, %	3.55	3.48	3.46	3.30	3.27
Ash, %	6.76	6.57	6.49	6.30	6.24

Note: SCP, Standard crude protein; LCP, low crude protein; TCP, tricalcium phosphate; CP, crude protein, ME, Metabolizable energy, ¹ Provided per kg of complete diet: 37.5 mg Zinc (as ZnSO₄); 37.5 mg Mn (as MnO₂); 37.5 mg Fe (as FeSO₄·7H₂O); 3.75 mg Copper (as CuSO₄·5H₂O); 0.83 mg I (as KI); and 0.23 mg Se (as Na₂SeO₃·5H₂O). ² Provided per kg of complete diet: 15 000 IU of vitamin A, 37.5 IU of vitamin E, 3750 IU of vitamin D3, 2.55 mg of vitamin K3, 24 µg of vitamin B12, 51 mg of niacin, 1.5 mg of folic acid, 3 mg of thiamin, 7.5 mg of riboflavin, 4.5 mg of vitamin B6, 0.2 mg of biotin, and 13.5 mg of calcium pantothenate.

Table 2. The effect of AJE supplementation on growth performance in broilers.

Items	SCP	LCP	TRT1	TRT2	SEM	SCP vs LCP	SCP vs TRT1	SCP vs TRT2	LCP vs TRT2
Day 8 to 21									
BWG, g	581	568	584	570	9	0.5176	0.8547	0.4553	0.9189
FI, g	926	914	918	923	24	0.6701	0.7709	0.4437	0.7299
FCR	1.595	1.614	1.571	1.62	0.048	0.7713	0.7137	0.9292	0.8399
Day 22 to 35									
BWG, g	938	902	972	920	14	0.1693	0.0338	0.1309	0.8805
FI, g	1681	1663	1668	1639	16	0.3455	0.1834	0.4816	0.1960
FCR	1.792	1.844	1.721	1.782	0.033	0.3049	0.1425	0.4429	0.7882
Overall									
BWG, g	1622	1574	1661	1599	12	0.1248	0.0911	0.2189	0.9060
FI, g	2734	2701	2713	2694	25	0.3950	0.5764	0.1147	0.4387
FCR	1.685	1.717	1.635	1.686	0.017	0.2036	0.0980	0.4760	0.5597

Note: SCP, standard crude protein; LCP, low crude protein; AJE, *Achyranthes Japonica* extract; TRT1, SCP + 0.025% AJE, TRT2, LCP + 0.050% AJE. SEM, Standard error of the mean. BWG, body weight gain; FI, Feed intake; FCR, Feed conversion ratio;

352 Table 3. The effect of AJE supplementation on nutrient digestibility in broilers.

Items	SCP	LCP	TRT1	TRT2	SEM	SCP vs LCP	SCP vs TRT1	SCP vs TRT2	LCP vs TRT2
Day 35									
Dry matter	70.71	69.52	75.16	71.30	1.40	0.2423	0.0283	0.2492	0.3964
Nitrogen	65.11	64.28	70.44	67.60	2.17	0.7088	0.0478	0.4982	0.4438
Energy	71.44	70.33	72.81	70.90	1.09	0.5238	0.4340	0.7536	0.7449

Note: SCP, standard crude protein, LCP, low crude protein; AJE, *Achyranthes Japonica* extract; TRT1, SCP + 0.025% AJE, TRT2, LCP + 0.050% AJE. SEM, standard error of the mean.

353

354

355

ACCEPTED

Table 4. The effect of AJE supplementation on microbial in broilers.

Items	SCP	LCP	TRT1	TRT2	SEM	SCP vs LCP	SCP vs TRT1	SCP vs TRT2	LCP vs TRT2
D 35									
<i>Lactobacillus</i>	7.82	7.67	8.02	7.69	0.08	0.1023	0.0274	0.1878	0.4513
<i>Escherichia coli</i>	5.82	5.92	5.56	5.85	0.09	0.2029	0.0457	0.2977	0.5440
<i>Salmonella</i>	3.73	3.87	3.68	3.71	0.11	0.5534	0.7614	0.9301	0.4967

Note: SCP, standard crude protein; LCP, low crude protein; AJE, *Achyranthes Japonica* extract; TRT1, SCP + 0.025% AJE; TRT2, LCP + 0.050% AJE; SEM, Standard error of the mean.

359

360 Table 5. The effect of AJE supplementation on gas emissions in broilers.

Items	SCP	LCP	TRT1	TRT2	SEM	SCP vs LCP	SCP vs TRT1	SCP vs TRT2	LCP vs TRT2
D 35									
NH ₃	16.75	13.50	12.00	9.00	1.14	0.1124	0.0589	0.0614	0.4845
H ₂ S	1.08	1.23	0.95	0.90	0.41	0.8713	0.8396	0.7771	0.6572
Methyl mercaptans	6.50	7.00	5.25	5.25	1.94	0.4902	0.6643	0.6643	0.2699
CO ₂	1450	1450	1425	1375	280	0.5878	0.9456	0.8381	0.4586
Acetic acid	2.75	3.25	2.75	2.50	0.77	0.6690	1.0000	0.8303	0.5234

Note: SCP, standard crude protein; LCP, low crude protein; AJE, *Achyranthes Japonica* extract; TRT1, SCP + 0.025% AJE, TRT2, LCP + 0.050% AJE; SEM, standard error of the mean.

361

362

363

364

365

366 Table 6. The effect of AJE supplementation on meat quality in broilers.

Items	SCP	LCP	TRT 1	TRT 2	SEM	SCP vs LCP	SCP vs TRT1	SCP vs TRT2	LCP vs TRT2
pH value	7.76	7.59	7.58	7.68	0.04	0.0010	0.0008	0.0035	0.5567
Breast muscle color									
Lightness(L*)	56.0 0	57.0 0	57.8 6	57.6 4	1.18	0.5496	0.2731	0.4570	0.8819
Redness(a*)	12.1 8	12.5 8	11.5 5	12.6 1	0.42	0.5828	0.3948	0.2940	0.6072
Yellowness(b*)	12.1 2	10.7 8	13.2 8	12.2 4	1.11	0.3609	0.4320	0.6516	0.1804
WHC, %	52.0 3	53.8 8	54.5 6	53.7 5	3.44	0.7154	0.6177	0.8959	0.8148
Cooking loss, %	20.0 3	18.5 6	20.5 8	20.5 0	1.55	0.5840	0.8386	0.8732	0.6968
Drip loss, %									
d 1	3.03	3.70	3.16	3.54	0.71	0.4169	0.5375	0.5660	0.1750
d 3	6.94	6.52	6.21	6.42	0.37	0.5496	0.2731	0.4570	0.8819
d 5	13.4 0	13.1 3	13.7 1	12.7 8	0.66	0.5828	0.3948	0.2940	0.6072
d 7	17.0 5	17.9 4	16.6 7	16.8 1	0.45	0.3609	0.4320	0.6516	0.1804
Relative organ weight, %									
Breast muscle	17.7 9	18.5 6	19.1 1	18.1 8	0.52	0.1990	0.1920	0.6871	0.0992
Liver	2.66	2.64	2.60	2.61	0.18	0.6474	0.7758	0.8002	0.8373
Bursa of Fabricius	0.16	0.15	0.14	0.16	0.02	0.9990	0.4821	0.8598	0.8608
Abdominal fat	0.86	0.93	1.18	1.1	0.24	0.3234	0.2798	0.4067	0.0797
Spleen	0.17	0.18	0.16	0.18	0.02	0.7269	0.8725	0.7220	0.4833
Gizzard	1.99	1.92	1.68	1.96	0.14	0.8825	0.1148	0.8489	0.7355

Note: SCP, standard crude protein; LCP, low crude protein; AJE, *Achyranthes Japonica* extract; TRT1, SCP + 0.025% AJE, TRT2, LCP + 0.050% AJE. SEM, standard error of the mean. WHC, water holding capacity.

367

368