

ARTICLE INFORMATION	Fill in information in each box below
<b>Article Type</b>	Research article
<b>Article Title (within 20 words without abbreviations)</b>	Impact of Substrate Type and Growth Stage on Nutrient Composition and Convergence Efficiency of <i>Hermetia illucens</i> Larvae
<b>Running Title (within 10 words)</b>	Optimizing Black Soldier Fly Larvae Growth and Nutrient Accumulation
<b>Author</b>	SangHun Ha <sup>1</sup> , Abdolreza Hosseindoust <sup>1</sup> , JunYoung Mun, SeRin Park, SoDam Choi, SeonAh Park, JinSoo Kim  <sup>1</sup> These authors contributed equally to this work.
<b>Affiliation</b>	Department of Animal Industry Convergence, Kangwon National University, Chuncheon 24341, Republic of Korea
<b>ORCID (for more information, please visit <a href="https://orcid.org">https://orcid.org</a>)</b>	Abdolreza Hosseindoust (0000-0001-9191-0613)  SangHun Ha (0000-0003-3779-1144)  JunYoung Mun (0000-0002-3075-7157)  SeRin Park (0009-0007-1413-5853)  SoDam Choi (0009-0009-2711-7106)  SeonAh Park (0009-0007-6584-2777)  JinSoo Kim (0000-0002-9518-7917)
<b>Competing interests</b>	No potential conflict of interest relevant to this article was reported.
<b>Funding sources</b>  State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry (IPET) through ((Companion Animal Life Cycle Industry Technology Development Program), funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA)(322094-3)
<b>Acknowledgements</b>	
<b>Availability of data and material</b>	Upon reasonable request, the datasets of this study can be available from the corresponding author.

<p><b>Authors' contributions</b></p> <p>Please specify the authors' role using this form.</p>	<p>Conceptualization: Hosseindoust A, Ha S.H, Mun J.Y.</p> <p>Data curation: Hosseindoust A, Kim J.S.</p> <p>Formal analysis: Mun J.Y, Park SA, Choi SD</p> <p>Validation: Hosseindoust A, Ha S.H, Park SR, Park SA</p> <p>Investigation: Ha S.H, Mun J.Y, Choi SD</p> <p>Writing - original draft: Hosseindoust A, Ha S.H, Mun J.Y, Kim J.S.</p> <p>Writing - review &amp; editing: Hosseindoust A, Ha S.H, Mun J.Y</p>
<p><b>Ethics approval and consent to participate</b></p>	<p>The animal care and experimental protocols used in this study received approval by the Institutional Animal Care and Use Committee of Kangwon National University (Ethical code: 210503-6).</p>

1

**CORRESPONDING AUTHOR CONTACT INFORMATION**

<p><b>For the corresponding author (responsible for correspondence, proofreading, and reprints)</b></p>	<p><b>Fill in information in each box below</b></p>
<p>First name, middle initial, last name</p>	<p>JinSoo Kim</p>
<p>Email address – this is where your proofs will be sent</p>	<p>Kjs896@kangwon.ac.kr</p>
<p>Secondary Email address</p>	
<p>Address</p>	<p>JinSoo Kim, Department of Animal Industry Convergence, Kangwon National University, Chuncheon 24341, Korea.</p>
<p>Cell phone number</p>	<p>+82-10-2566-5961</p>
<p>Office phone number</p>	<p>+82-33-250-8616</p>
<p>Fax number</p>	<p>+82-33-259-5572</p>

2

### 3 **Abstract**

4 This study investigates the bioconversion efficiency and nutrient accumulation in black soldier fly (BSF;  
5 *Hermetia illucens*), focusing on the effects of feeding on two different substrates (tofu by-products and  
6 food waste) and harvesting at two developmental stages (larvae and prepupae) within a 2×2 factorial  
7 arrangement. The growth performance, conversion efficiency, nutrient composition, amino acid profile,  
8 fatty acid composition, and nutrient composition of BSF meal were assessed. Results indicated that BSF  
9 reared on tofu by-products exhibited superior weight gain compared to those fed food waste, with  
10 significant enhancements observed in weight, length, and width upon harvesting at the prepupae stage.  
11 Moreover, tofu by-products promoted higher bioconversion rates, protein conversion efficiency, and  
12 lipid yield, while food waste favored lipid conversion. Analysis of nutrient composition revealed higher  
13 crude protein content in BSFs fed tofu by-products, with elevated levels of crude protein, ether extract,  
14 and chitin in prepupae-stage BSFs. Higher concentrations of isoleucine, leucine, and tryptophan were  
15 observed in tofu by-product-fed BSF. Conversely, BSFs harvested at the prepupae stage exhibited  
16 increased levels of threonine, alanine, and tyrosine, regardless of substrate. Higher proportions of  $\alpha$ -  
17 linolenic acid (C18:3n3) and docosahexaenoic acid (C22:6n3) were observed in tofu by-product-fed  
18 BSF. Conversely, BSF harvested at the larval stage displayed higher levels of saturated fatty acids,  
19 including lauric acid (C12:0) and myristic acid (C14:0). In conclusion, tofu by-products emerged as a  
20 promising substrate for enhancing essential amino acid and unsaturated fatty acid content in BSF, while  
21 harvesting at the prepupae stage offered advantages in nutrient density and storage stability of the  
22 harvested biomass.

23 **Keywords:** Food waste, tofu, bioconversion, larvae, prepupae, black soldier fly

24

## 25 INTRODUCTION

26 Sustainable agriculture strives to meet the world nutritional needs while minimizing environmental  
27 degradation and resource depletion [1,2]. Insects have emerged as promising candidates owing to their  
28 environmental sustainability and efficiency [3]. Among these alternatives, the black soldier fly (BSF;  
29 *Hermetia illucens*) has garnered considerable attention in recent years due to its potential to address  
30 sustainability challenges in agriculture [4,5]. The larvae of BSF offer numerous advantages as a protein  
31 source for animal nutrition, offering a viable solution to food security concerns. A primary advantage  
32 lies in BSF remarkable efficiency in converting organic waste into protein biomass [4,6,7]. These  
33 voracious larvae possess the ability to decompose a wide array of organic substrates such as food waste,  
34 agricultural residue, and animal manure into nutrient-rich biomass [2,3,8], presenting a sustainable  
35 solution to waste management while augmenting feed production. The larvae emit fewer greenhouse  
36 gases, rendering BSF cultivation a more eco-friendly alternative to traditional livestock farming  
37 practices [9].

38 Economically, BSF production offers cost-effective advantages over conventional livestock rearing [7].  
39 Notably, by-products from tofu processing, such as tofu whey, represent an underutilized feed resource  
40 for BSF rearing [3]. The abundance of tofu waste, estimated at millions of tons annually in select regions,  
41 underscores the potential for BSF cultivation to mitigate waste disposal challenges while generating  
42 valuable protein biomass [3].

43 Nevertheless, the nutritional composition and growth performance of BSF larvae are influenced by their  
44 diet, particularly botanical waste substrates [10–12]. Optimization of dietary inputs is pivotal to  
45 maximizing protein production efficiency per unit of organic waste, thereby enhancing the sustainability  
46 and cost-effectiveness of BSF-based feed production [8]. Moreover, the nutrient composition of BSF  
47 meal can be under the influence of the growth stage harvest including larvae and prepupae [13]. Thus,  
48 a systematic evaluation of BSF's growth dynamics, conversion efficiency, and larval quality across  
49 different botanical waste substrates is imperative to identify optimal feeding regimes.

50 This study endeavors to assess the growth performance, conversion efficiency, and larval quality of  
51 BSF reared on various botanical waste substrates, namely food waste and tofu by-products. By  
52 elucidating the effects of substrate type and developmental stage (larvae or prepupae) on nutrient  
53 composition, weight, length, development time, substrate consumption index, conversion efficiency,  
54 and fatty acid composition, this research aims to contribute to the optimization of BSF rearing practices  
55 for sustainable and efficient protein production in the context of waste valorization and animal feed  
56 supplementation.

57

## 58 MATERIAL AND METHODS

### 59 Larvae rearing control

60 The experiment was conducted at a commercial BSF farm located in Eui-sung, Gyeongsangbuk-do,  
61 Republic of Korea. Prior to hatching, the BSF insects used in this study were sourced from the farm.  
62 The trial colony was meticulously maintained under controlled environmental conditions, with a  
63 temperature set at  $27\pm 0.3^{\circ}\text{C}$ , relative humidity maintained at  $65\pm 8\%$ , and a photophase of 16 hours of  
64 natural light supplemented with LED lighting providing an intensity of 6000 lux. For the experimental  
65 setup, a total of 30 egg clutches were utilized, each containing approximately 50,000 larvae. The  
66 average weight of the egg clutches was measured at  $24.16\pm 0.91$  mg per clutch, with an average egg  
67 weight of 0.027 mg. The egg clutches were further divided into four treatment groups, comprising two  
68 different substrates (tofu by-product and food waste; Table 1) and two distinct growth stages (larvae  
69 and prepupae). Each treatment group consisted of 10 replicates, housed within experimental boxes  
70 measuring  $1,064\times 507\times 320$  mm (length $\times$ width $\times$ height). The trial continued until the end of the prepupal  
71 stage, just prior to the pupal stage transition. Egg counting within each clutch was performed according  
72 to the methodology outlined by Georgescu et al. c, employing an Alpha model binocular magnifier with  
73 7x to 45x magnification. To ensure accuracy, the eggs were dispersed in 70% ethanol and subsequently  
74 captured and enumerated through photographic means utilizing the ClickMaster2000 1.0 software,

75 accessible at <https://www.thregr.org/~wavexx/>. This meticulous process facilitated precise egg counting,  
76 essential for subsequent experimental procedures and data analysis.

### 77 **Assessment of larval growth performance**

78 To assess the growth performance of BSF larvae, measurements were taken at the time of harvest,  
79 including weight, length, width, and survivability. In order to standardize the comparison across harvest  
80 days, 100 larvae were randomly selected for measurement beginning on day 10, and continuing until  
81 the average larval weight reached 190 mg. The larval yield, indicative of survivability on the harvest  
82 day, was calculated using the following equation [3]:

$$\text{Larval yield \%} = \frac{\text{Larvae}_F}{\text{Larvae}_I} \times 100$$

83 Where:

84  $\text{Larvae}_F$  represents the number of larvae at the end of the trial.  $\text{Larvae}_I$  represents the number of larvae  
85 at the initiation of the trial.

86 This approach allowed for a comprehensive evaluation of larval growth and survival dynamics, essential  
87 for assessing the efficacy of different experimental treatments. Additionally, the utilization of  
88 standardized measurement protocols ensured accuracy and reliability in the assessment of larval  
89 performance across varying experimental conditions.

### 90 **Substrates preparation**

91 The substrates utilized in this study were procured from local farms and facilities to ensure relevance  
92 to regional waste streams. Prior to their use, dried substrates were moistened with water to achieve a  
93 target moisture content of 65%. Egg blocks were then positioned on two distinct substrates: food waste  
94 and tofu by-products. The food waste substrate consisted of organic waste collected from households,  
95 kitchens, and local dining establishments through a municipal waste collection facility located in Eui-  
96 sung, Gyeong-sangbuk-do, Republic of Korea. Tofu by-products, comprising soybean curd residue,  
97 were sourced from a local tofu manufacturing facility situated in Kang-reung, Kangwon-do, Republic

98 of Korea. To prevent fermentation and ensure uniformity across substrates, the materials were promptly  
99 dried using an industrial drying apparatus (Model: KAPD-A098D, Manufactured in Gwangju, Republic  
100 of Korea), operating at a temperature of 60°C for 48 hours, until the moisture content was reduced to  
101 less than 8%. Upon commencement of the experiment, water was applied to the substrates to reach the  
102 desired moisture level of 65%. Each experimental box was provisioned with a total of 20 kg (dry matter  
103 basis) of substrate. To create a suitable environment for larval development and prevent drowning, a  
104 layer of 1 kg of recycled paper pulp and 600 g of sawdust was spread atop the substrates. This bedding  
105 layer facilitated larval movement and provided an additional substrate for microbial activity, which may  
106 contribute to the decomposition process.

### 107 **Chemical composition analysis**

108 The chemical composition of the harvested larvae from each treatment was determined to assess their nutritional  
109 profile. The contents of dry matter (DM), crude protein (CP), ether extract (EE), crude ash (Ash), and chitin were  
110 analyzed using standardized methods outlined by the Association of Official Analytical Chemists [14]. Initially,  
111 fresh larvae were weighed to establish baseline measurements. Subsequently, the larvae were subjected to drying  
112 in an oven set at 85°C for a duration of three days to determine the DM content, following the procedure outlined  
113 in method 930.15 of the AOAC [14]. To quantify the protein content, AOAC official method 990.03 was  
114 employed, utilizing the Kjeldahl method. This method entails the digestion of the sample with sulfuric acid to  
115 release nitrogen, followed by conversion of the nitrogen to ammonia. The ammonia is then distilled and titrated  
116 to determine the nitrogen content, which is subsequently utilized to calculate the protein content [14]. Similarly,  
117 the ether extract content was determined using AOAC official method 990.15, involving the extraction of lipid  
118 content from the sample with a suitable solvent, followed by evaporation of the solvent to obtain the lipid content  
119 [14]. The gross energy of BSF larvae was measured using a bomb calorimeter (Model 1261, Parr Instrument Co.,  
120 Moline, IL, USA), providing insight into the calorific value of the larvae. Following analysis, any remaining  
121 samples were promptly stored at -20°C for future analysis and reference, ensuring preservation of sample integrity  
122 for subsequent investigations. The amino acid (AA) composition of BSF meal was assessed using high-  
123 performance liquid chromatography (HPLC) with a Waters 486 system (Waters, Milford, MA, USA).  
124 This analysis was conducted following acid hydrolysis, as described by Lee et al. [15], to release the  
125 amino acids from the protein matrix for quantification. To ensure comprehensive analysis, the

126 determination of methionine and cysteine was performed after oxidation with performic acid, following  
127 the methodology outlined by Kim et al. [16].

### 128 Larval convergence efficiency evaluation

129 To assess the convergence efficiency of larvae reared on different botanical wastes, several parameters  
130 including bioconversion rate (BCR), waste reduction rate (WR), protein conversion rate (PCR), lipid  
131 conversion rate (LCR), protein yield (PY), lipid yield (LY), and mass distribution patterns were  
132 evaluated using the following equations.

133 The BCR was calculated as:

$$134 \quad \text{BCR \%} = \frac{\text{DM}_F - \text{DM}_I}{\text{DM}_S} \times 100$$

135  
136 Where  $\text{DM}_F$  represents the dry matter of BSF larvae at harvest day,  $\text{DM}_I$  represents the dry matter of  
137 BSF larvae at the initiation of the trial, and  $\text{DM}_S$  represents the total dry matter of the substrate.

138 The WR was determined by:

$$\text{WR \%} = \frac{\text{DM}_S - \text{DM}_R}{\text{DM}_S} \times 100$$

139 Where  $\text{DM}_S$  represents the total dry matter of the substrate and  $\text{DM}_R$  represents the residual dry matter  
140 of the substrate after larval consumption.

141 The PCR was calculated as:

$$\text{PCR \%} = \frac{\text{DM}_F \times \text{Protein}_L \%}{\text{DM}_S} \times 100$$

142 Where  $\text{protein}_L$  % represents the protein content in the larvae,  $\text{DM}_F$  represents the dry matter of BSF  
143 larvae at harvest day, and  $\text{DM}_S$  represents the total dry matter of the substrate.

144 The LCR was determined by:

$$\text{LCR \%} = \frac{\text{DM}_F \times \text{EE}_L \%}{\text{DM}_S} \times 100$$

145 Where  $\text{EE}_L$  % represents the lipid content in the larvae,  $\text{DM}_F$  represents the dry matter of BSF larvae at  
 146 harvest day, and  $\text{DM}_S$  represents the total dry matter of the substrate.

147 The PY was calculated as:

$$\text{PY kg/c} = \text{g of DM}_F \text{ yield} \times \text{CP}_L \%$$

148 Where  $\text{DM}_F$  yield represents the dry matter of BSF larvae at harvest, and  $\text{CP}_L$  % represents the crude  
 149 protein content in the larvae.

150 The LY was determined as:

$$\text{LY kg/c} = \text{g of DM}_F \text{ yield} \times \text{EE}_L \%$$

151 Where  $\text{DM}_F$  yield represents the dry matter of BSF larvae at harvest,  $\text{EE}_L$  % represents the ether extract  
 152 content in the larvae, and 'c' represents the clutch size (50,000 larvae).

153

#### 154 **Fatty acid composition**

155 The determination of fatty acid content was conducted following the methodology outlined by  
 156 Georgescu et al. [17]. For analysis, 10 g of meal sample was collected and subsequently placed on a  
 157 clean substrate for a 24-hour period. Following this, the samples were washed with 70% ethanol to  
 158 remove any impurities and then stored at  $-80^\circ\text{C}$  to maintain sample integrity. Extraction and  
 159 identification of fatty acid methyl esters (FAME) from BSF larvae fats were performed using gas  
 160 chromatography with mass spectrometry detection, adhering to established standards including AOAC-  
 161 969.33 [14], ISO 3657: 2002, ISO 12966-2: 2011, and ISO 12966-2: 2017. The analytical process  
 162 involved saponification of the fat, achieved by subjecting the samples to methanolic sodium hydroxide  
 163 solution (0.5 M) at  $210^\circ\text{C}$  on a sand bath with a reflux rate of 1 drop/s. Subsequently, esterification with

164 a 15% vol boron trifluoride catalyst and cooling with hexane were carried out. Gas chromatographic  
165 analysis was conducted using a Perkin Elmer chromatographic system equipped with a mass  
166 spectrometer detector (GC-MS). This system featured an Elite-Wax chromatographic column with a  
167 stationary polar phase polyethylene glycol measuring 30 m in length, 0.25 mm in internal diameter, and  
168 a film thickness of 1.0  $\mu\text{m}$ . Operating parameters included an injection port temperature of 220°C, an  
169 injected sample volume of 1.0  $\mu\text{L}$ , a helium carrier gas flow rate of 1.5 mL/min, and a splitting ratio of  
170 40:1. The mass spectrometer settings comprised a transfer line temperature of 150°C, a source  
171 temperature of 150°C, a multiplier of 1500, and a solvent delay of 0-1.5 min. Quantification of fatty  
172 acid concentration was achieved by comparing the relative retention time of FAME to the certified  
173 standard Mix FAME Supelco. The individual fatty acid concentrations were expressed as a percentage  
174 of the total identified FAME. Moreover, key ratios between various fatty acid categories, including total  
175 saturated fatty acids (SFA) and total unsaturated fatty acids, were calculated. Additionally, the ratios  
176 between fatty acids from the n-6 and n-3 series were determined to further elucidate the nutritional  
177 profile of the BSF biomass.

### 178 **Statistical analysis**

179 The data were subjected to analysis employing a 2 $\times$ 2 factorial arrangement utilizing the General Linear  
180 Model procedure of SAS (version 9.4, 1996, SAS Inst. Inc., Cary, NC, USA). Differences in various  
181 parameters including growing performance, conversion efficiency, nutrient composition, amino acid,  
182 and fatty acid concentrations among the treatments were assessed using a two-way ANOVA test. To  
183 further elucidate the differences between treatments, the Tukey test was employed to compare the means  
184 of the treatments. Statistical significance was determined at a threshold of  $p < 0.05$ . Moreover, a p-value  
185 less than 0.1 was considered indicative of a trend. The results are expressed as the mean  $\pm$  standard  
186 deviation, providing insights into the central tendency and variability of the data.

187

## 188 **RESULTS**

189 **Growth performance**

190 The BSF larvae reared on tofu by-products exhibited significantly higher ( $p < 0.05$ ) weight gain  
191 compared to those fed with food waste (Table 2). Moreover, harvesting at the prepupae stage resulted  
192 in notable enhancements in weight, length, and width of BSF, indicating superior growth performance  
193 compared to larvae-harvested BSF, which displayed higher survival rates. The BSF larvae reared on  
194 tofu by-products exhibited significantly higher ( $p < 0.01$ ) BCR, PCR, PY, and LY compared to those  
195 fed with food waste (Table 2), however, LCR was lower in BSF fed tofu by-products compared with  
196 food waste. Moreover, harvesting at the prepupae stage resulted in lower BCR compared to larvae  
197 harvested BSF, however, harvesting at the prepupae stage showed higher PY, LCR, PY, and LY  
198 compared to larvae harvested BSF.

199

200 **Nutrient composition of meal**

201 The analysis revealed no significant differences in dry matter, gross energy, or ash content across  
202 different substrates and developmental stages (Table 3). However, BSFs fed on tofu by-products  
203 exhibited a higher crude protein content compared to those fed with food waste. Furthermore, harvesting  
204 at the prepupae stage led to elevated levels of crude protein, ether extract, and chitin, indicating  
205 enhanced nutrient composition in prepupae-stage BSF.

206 **Amino acid profile of the meal**

207 Amino acid profiling demonstrated distinct compositions in BSF fed with different substrates and  
208 harvested at different developmental stages (Table 4). BSFs reared on tofu by-products showed  
209 increased concentrations of isoleucine, leucine, tryptophan, alanine, and proline. In contrast, BSFs  
210 harvested at the prepupae stage displayed higher levels of threonine, tryptophan, alanine, serine, and  
211 tyrosine, indicating variations in amino acid profiles influenced by both substrate type and  
212 developmental stage.

213 **Fatty acid composition of meal**

214 Fatty acid analysis revealed variations in the composition of BSF lipids based on substrate and  
215 developmental stage (Table 5). BSFs fed on tofu by-products exhibited higher ratios of C10:0, C12:0,  
216 C14:0, C14:1, C18:3n3, and C20:1 fatty acids, while those consuming food waste showed a higher  
217 content of C18:4. Additionally, SFA such as C12:0, C15:0, C17:0, and C18:1 were more abundant in  
218 BSFs harvested at the prepupae stage.

## 219 **DISCUSSION**

220 The higher weight gain observed in BSF larvae reared on tofu by-products compared to those fed with  
221 food waste suggests that the nutrient composition and bioavailability in tofu by-products are more  
222 conducive to larval growth. The higher protein in tofu by-products may facilitate improved growth rates.  
223 The enhancements in weight, length, and width of BSF when harvested at the prepupae stage could be  
224 attributed to the larval maturation process. As larvae transition into prepupae, they are in a state of  
225 maximal nutrient accumulation and tissue development [7]. This stage is also characterized by efficient  
226 energy storage and nutrient conversion in preparation for metamorphosis [13], which may explain the  
227 observed improvements in growth performance at this developmental stage. In contrast, the higher  
228 survival rates in larvae harvested BSF as opposed to those harvested at the prepupae stage may be linked  
229 to the life cycle and physiological state of the larvae. Larval stages possess more resilience and  
230 adaptability to varying environmental conditions and nutrient availability compared to prepupae [18].  
231 This higher adaptability results in increased survival rates when harvesting at the larval stage [8].  
232 Overall, these results highlight the importance of substrate choice and timing of harvest in optimizing  
233 BSF growth performance. The mode of action related to nutrient availability, larval development, and  
234 life cycle dynamics are central to understanding the differences observed in growth performance.

235 The lack of significant differences in dry matter, gross energy, and ash content across different substrates  
236 and developmental stages suggests that the overall energy density and mineral content of BSF biomass  
237 remain relatively consistent regardless of dietary inputs or harvest timing. However, BSF fed tofu by-  
238 products demonstrated higher crude protein content compared to those fed food waste. This difference  
239 can be attributed to the higher protein levels in tofu by-products [3]. Harvesting BSF at the prepupae

240 stage led to increased levels of crude protein, ether extract, and chitin. This suggests that as BSF larvae  
241 mature and transition to the prepupae stage, they undergo biochemical and physiological changes that  
242 enhance their nutrient composition [19]. The increase in crude protein and ether extract may be due to  
243 the prepupae preparation for metamorphosis, which involves accumulating energy and nutrient reserves.  
244 Additionally, elevated chitin levels are associated with the development of the exoskeleton as the insects  
245 approach the pupal stage [20].

246 The amino acid profile of BSF is a crucial aspect of its nutritional value and potential applications as  
247 animal feed [3,13]. BSF reared on tofu by-products exhibited increased concentrations of isoleucine,  
248 leucine, tryptophan, alanine, and proline compared to those fed with food waste. This suggests that tofu  
249 by-products provide a rich source of essential and conditionally essential amino acids, contributing to  
250 a more balanced and nutritious profile for the larvae. The higher levels of these amino acids may result  
251 from the nutrient-rich nature of tofu by-products, which could provide an optimal blend of proteins and  
252 other macronutrients for larval development. In contrast, BSF harvested at the prepupae stage displayed  
253 elevated levels of threonine, tryptophan, alanine, serine, and tyrosine compared to those harvested at  
254 the larval stage. This shift in amino acid composition during the prepupae stage is linked to the  
255 physiological changes the insects undergo in preparation for metamorphosis [7,13,21]. The variation in  
256 amino acid profiles based on substrate type and developmental stage underscores the importance of  
257 optimizing these factors for specific applications of BSF biomass.

258 The fatty acid composition of BSF is a critical aspect of its nutritional quality and potential applications  
259 in various industries, including animal feed and biodiesel production [2,13]. BSF fed on tofu by-  
260 products exhibited higher ratios of specific fatty acids, including C10:0, C12:0, C14:0, C14:1, C18:3n3,  
261 and C20:1, compared to those consuming food waste. These differences in fatty acid composition may  
262 be due to the lipid profiles present in the respective substrates. The observed higher content of C18:4 in  
263 BSF consuming food waste suggests preferential utilization or accumulation of this fatty acid in  
264 response to the substrate lipid content. Additionally, SFA such as C12:0, C15:0, C17:0, and C18:1 were  
265 more abundant in BSF harvested at the prepupae stage. This is attributed to metabolic changes

266 associated with larval development and maturation into prepupae [19]. As the larvae transition to the  
267 prepupal stage, they undergo metabolic shifts to support pupation and subsequent adult emergence  
268 [2,19], leading to alterations in fatty acid synthesis and accumulation. The variations in the fatty acid  
269 composition of BSF biomass have significant implications for its nutritional value and suitability for  
270 various applications [22]. For instance, certain fatty acids, such as omega-3 and omega-6 fatty acids,  
271 are essential for animal health [1,2,11] and contribute to the nutritional quality of BSF-based feed  
272 formulations.

273 BSF meal derived from larvae reared on tofu by-products exhibited higher gross energy compared to  
274 meal from food waste-fed BSF. This difference could be attributed to the higher energy density of the  
275 tofu by-product substrate, which may have contributed to enhanced energy accumulation in the larvae.  
276 As a result, the meal from tofu by-product-fed BSF could provide a more energy-rich feed ingredient.  
277 Moreover, meals from prepupae-stage BSF demonstrated enhanced gross energy, crude protein, and  
278 chitin content. These changes can be due to the physiological and biochemical transformations that  
279 occur as the larvae mature into prepupae [2,6,19]. The increased gross energy may result from the  
280 accumulation of energy reserves to support metamorphosis, while the higher crude protein and chitin  
281 levels are associated with tissue development and exoskeleton formation during the pupal transition  
282 [2,20]. The distinct fatty acid profile favoring C12:0, C18:3n3, and C22:6n3 in meals from prepupae-  
283 stage BSFs suggests alterations in fatty acid metabolism as the larvae approach the pupal stage. These  
284 specific fatty acids may be selectively synthesized or accumulated in response to developmental cues,  
285 contributing to the observed fatty acid composition. The higher levels of SFA in prepupae-stage BSF  
286 meal could reflect shifts in metabolic pathways aimed at supporting pupation and adult emergence  
287 [9,10]. Saturated fatty acids play a role in providing stable energy sources during this critical  
288 developmental phase [2,10,17]. Understanding the mode of action underlying these variations in  
289 nutrient composition can help optimize BSF meal production for targeted applications. The higher gross  
290 energy and crude protein content in prepupae-stage BSF meal increases its value as a feed ingredient  
291 for high-performance livestock.

292 **Conclusion**

293 Our results demonstrate that substrate type significantly influences the growth performance, nutrient  
294 composition, and fatty acid profile of BSF larvae and prepupae. Larvae reared on tofu by-products  
295 exhibited superior growth performance and higher levels of crude protein, while those fed food waste  
296 displayed distinct fatty acid compositions. Additionally, harvesting BSF at the prepupae stage resulted  
297 in enhanced nutrient composition and fatty acid profiles, indicating the importance of the developmental  
298 stage in optimizing biomass quality. Furthermore, it was revealed that by understanding the mode of  
299 action underlying these variations, we can develop strategies to optimize BSF production systems for  
300 farm animal nutrition, sustainable agriculture, and waste management.

301

ACCEPTED

302 REFERENCES

- 303 1. Yamamoto FY, Suehs BA, Ellis M, Bowles PR, Older CE, Hume ME, et al. Dietary fishmeal  
304 replacement by black soldier fly larvae meals affected red drum (*Sciaenops ocellatus*) production  
305 performance and intestinal microbiota depending on what feed substrate the insect larvae were  
306 offered. *Anim Feed Sci Technol.* 2022;283:115179.  
307 <https://doi.org/10.1016/j.anifeedsci.2021.115179>
- 308 2. Hosseindoust A, Ha SH, Mun JY, Kim JS. A metanalysis to evaluate the effects of substrate sources  
309 on the nutritional performance of black soldier fly larvae: implications for sustainable poultry feed.  
310 *Poult Sci.* 2024;103. <https://doi.org/10.1016/j.psj.2023.103299>
- 311 3. Hosseindoust A, Ha SH, Mun JY, Kim JS. Quality Characteristics of Black Soldier Flies Produced  
312 by Different Substrates. *Insects.* 2023;14:500. <https://doi.org/10.3390/insects14060500>
- 313 4. Dzepe D, Magatsing O, Kuietche HM, Meutchieye F, Nana P, Tchuinkam T, et al. Recycling Organic  
314 Wastes Using Black Soldier Fly and House Fly Larvae as Broiler Feed. *Circ Econ Sustain*  
315 *[Internet]*. 2021;1:895–906. Available from: <https://doi.org/10.1007/s43615-021-00038-9>
- 316 5. Chang S, Song M, Lee J, Oh H, Song D, An J, et al. Effect of black soldier fly larvae as substitutes  
317 for fishmeal in broiler diet. *J Anim Sci Technol.* 2023;65. <https://doi.org/10.5187/jast.2023.e89>
- 318 6. Giannetto A, Oliva S, Riolo K, Savastano D, Parrino V, Cappello T, et al. Waste valorization via  
319 *hermetia illucens* to produce protein-rich biomass for feed: Insight into the critical nutrient taurine.  
320 *Animals.* 2020;10:1710. <https://doi.org/10.3390/ani10091710>
- 321 7. Fuso A, Barbi S, Macavei LI, Luparelli AV, Maistrello L, Montorsi M, et al. Effect of the rearing  
322 substrate on total protein and amino acid composition in black soldier fly. *Foods.* 2021;10:1773.  
323 <https://doi.org/10.3390/foods10081773>
- 324 8. Rehman K ur, Rehman A, Cai M, Zheng L, Xiao X, Somroo AA, et al. Conversion of mixtures of  
325 dairy manure and soybean curd residue by black soldier fly larvae (*Hermetia illucens* L.). *J Clean*  
326 *Prod.* 2017;154:366–73. <https://doi.org/10.1016/j.jclepro.2017.04.019>
- 327 9. Danieli PP, Lussiana C, Gasco L, Amici A, Ronchi B. The effects of diet formulation on the yield,  
328 proximate composition, and fatty acid profile of the black soldier fly (*Hermetia illucens* L.)  
329 prepupae intended for animal feed. *Animals.* 2019;9:178. <https://doi.org/10.3390/ani9040178>
- 330 10. Hoc B, Genva M, Fauconnier ML, Lognay G, Francis F, Caparros Megido R. About lipid  
331 metabolism in *Hermetia illucens* (L. 1758): on the origin of fatty acids in prepupae. *Sci Rep.*  
332 2020;10:11916. <https://doi.org/10.1038/s41598-020-68784-8>
- 333 11. Rodrigues DP, Calado R, Pinho M, Rosário Domingues M, Antonio Vázquez J, Ameixa OMCC.  
334 Bioconversion and performance of Black Soldier Fly (*Hermetia illucens*) in the recovery of

- 335 nutrients from expired fish feeds. *Waste Manag.* 2022;141:183–93.  
336 <https://doi.org/10.1016/j.wasman.2022.01.035>
- 337 12. Oh JH, Karadeniz F, Yang J, Lee H, Choi M-N, Jeon S, et al. Antioxidant, anti-inflammatory, anti-  
338 adipogenesis activities and proximate composition of *Hermetia illucens* larvae reared on food  
339 waste enriched with different wastes. *J Anim Sci Technol.* 2023;  
340 <https://doi.org/10.5187/jast.2023.e87>
- 341 13. Luparelli AV, Jasmine HS, Lolli V, Lazzi C, Sforza S, Caligiani A. Dynamic changes in molecular  
342 composition of black soldier fly prepupae and derived biomasses with microbial fermentation.  
343 *Food Chem X.* 2022;14:100327. <https://doi.org/10.1016/j.fochx.2022.100327>
- 344 14. Association of Official Analytical Chemists. *Official Methods of Analysis of AOAC International.*  
345 *Assoc Off Anal Chem Int.* 2007;1.
- 346 15. Lee SH, Hosseindoust A, Laxman Ingale S, Rathi PC, Yoon SY, Choi JW, et al. Thermostable  
347 xylanase derived from *Trichoderma citrinoviride* increases growth performance and non-starch  
348 polysaccharide degradation in broiler chickens. *Br Poult Sci.* 2020;61:57–62.  
349 <https://doi.org/10.1080/00071668.2019.1673316>
- 350 16. Kim JS, Hosseindoust AR, Shim YH, Lee SH, Choi YH, Kim MJ, et al. Processing diets containing  
351 corn distillers' dried grains with solubles in growing broiler chickens: Effects on performance,  
352 pellet quality, ileal amino acids digestibility, and intestinal microbiota. *Poult Sci.* 2018;97:2411–  
353 8. <https://doi.org/10.3382/ps/pey075>
- 354 17. Georgescu B, Boaru AM, Muntean L, Sima N, Struți DI, Păpuc TA, et al. Modulating the Fatty  
355 Acid Profiles of *Hermetia illucens* Larvae Fats by Dietary Enrichment with Different Oilseeds: A  
356 Sustainable Way for Future Use in Feed and Food. *Insects.* 2022.  
357 <https://doi.org/10.3390/insects13090801>
- 358 18. Grossule V, Lavagnolo MC. The treatment of leachate using Black Soldier Fly (BSF) larvae:  
359 Adaptability and resource recovery testing. *J Environ Manage.* 2020;253.  
360 <https://doi.org/10.1016/j.jenvman.2019.109707>
- 361 19. Bekker NS, Heidelberg S, Vestergaard SZ, Nielsen ME, Riisgaard-Jensen M, Zeuner EJ, et al.  
362 Impact of substrate moisture content on growth and metabolic performance of black soldier fly  
363 larvae. *Waste Manag.* 2021;127. <https://doi.org/10.1016/j.wasman.2021.04.028>
- 364 20. Mei H, Li C, Li X, Hu B, Lu L, Tomberlin JK, et al. Characteristics of tylosin and enrofloxacin  
365 degradation in swine manure digested by black soldier fly (*Hermetia illucens* L.) larvae. *Environ*  
366 *Pollut.* 2022;293. <https://doi.org/10.1016/j.envpol.2021.118495>
- 367 21. Hosseindoust A, Mun J, Ha SH, Kim J. Effects of meal processing of black soldier fly on  
368 standardized amino acids digestibility in pigs. *J Anim Sci Technol.* Korean Society of Animal  
369 Science and Technology; 2023; <https://doi.org/10.5187/jast.2023.e28>

370 22. Kim B, Bang HT, Kim KH, Kim MJ, Jeong JY, Chun JL, et al. Evaluation of black soldier fly larvae  
371 oil as a dietary fat source in broiler chicken diets. *J Anim Sci Technol.* 2020;62.  
372 <https://doi:10.5187/jast.2020.62.2.187>

373

ACCEPTED

## 374 TABLES

375 Table 1. Nutrient composition of botanical waste used in trial.

Substrate (%)	Food waste	Tofu By-product
Dry matter	16.95	23.96
Crude protein	19.68	30.52
Ether extract	21.65	12.95
Ash	6.85	7.96

376

Table 2. Effects of substrate (SUB) and development stage (STG) on growth performance of black soldier fly.

Treatments	Food waste		Tofu by-product		SEM	P-value		
	Larvae	Prepupae	Larvae	Prepupae		SUB	STG	SUB×STG
Larvae growth								
Weight, mg	177.92	191.05	194.97	209.39	2.178	<0.001	<0.001	0.802
Length, mm	13.83	15.20	14.20	15.21	0.132	0.305	<0.001	0.342
Width, mm	2.79	2.94	2.84	2.95	0.021	0.477	0.002	0.614
Survival rate, %	80.14	77.53	77.97	75.50	0.575	0.056	0.022	0.948
Substrate consumption, %	69.01	77.65	72.16	79.55	0.971	0.090	<0.001	0.671
.....								
Conversion efficiency rate, %								
Bioconversion	12.10	11.61	12.67	12.36	0.084	<0.001	0.001	0.440
Protein conversion	6.12	6.93	9.91	11.02	0.040	<0.001	<0.001	0.069
Lipid conversion	5.88	6.64	4.45	4.96	0.025	<0.001	<0.001	0.082
Protein yield, kg/c	1.13	1.32	1.32	1.47	0.029	0.008	0.005	0.859
Lipid yield, kg/c	1.03	1.16	1.13	1.26	0.022	0.039	0.006	0.902

377 SEM, standard error of mean. Each value is the average of 10 replicates.

Table 3. Effects of substrate (SUB) and development stage (STG) on nutrient composition of black soldier fly.

Treatments	Food waste		Tofu by-product		SEM	P-value		
	Larvae	Prepupae	Larvae	Prepupae		SUB	STG	SUB×STG
DM, %	40.60	42.73	41.57	42.97	0.689	0.662	0.207	0.792
GE, kcal	4,315	4,343	4,388	4,410	30.0	0.253	0.684	0.964
CP, % of DM	39.12	41.66	41.63	43.41	0.326	0.002	0.002	0.569
EE, % of DM	35.53	36.67	35.76	37.02	0.283	0.617	0.041	0.920
Ash, % of DM	7.04	7.24	7.10	7.29	0.167	0.872	0.564	0.916
Chitin, % of DM	7.85	8.45	8.15	8.60	0.125	0.257	0.049	0.882

378 SEM, standard error of mean; DM, dry matter; GE, gross energy; EE, ether extract. Each value is the  
 379 average of 10 replicates.

380

Table 4. Effects of substrate (SUB) and development stage (STG) on amino acid profile of black soldier fly.

Treatments	Food waste		Tofu by-product		SEM	P-value		
	Larvae	Prepupae	Larvae	Prepupae		SUB	STG	SUB×STG
Essential amino acid, %								
Arginine	1.86	1.95	1.74	1.93	0.036	0.361	0.061	0.520
Histidine	1.06	1.07	1.19	1.11	0.028	0.296	0.062	0.710
Isoleucine	1.45	1.16	1.66	1.83	0.031	0.002	0.016	0.958
Leucine	2.39	2.44	2.53	2.68	0.030	0.005	0.114	0.426
Lysine	2.81	2.92	2.84	2.93	0.062	0.836	0.423	0.931
Methionine	0.59	0.64	0.54	0.58	0.020	0.219	0.321	0.951
Phenylalanine	1.07	1.21	1.14	1.23	0.033	0.109	0.164	0.459
Threonine	1.36	1.48	1.38	1.58	0.022	0.205	0.002	0.400
Tryptophan	1.97	2.21	2.23	2.32	0.030	0.007	0.015	0.243
Valine	2.53	2.58	2.59	2.74	0.076	0.479	0.513	0.716
Non-essential amino acid, %								
Alanine	2.46	2.56	2.63	2.77	0.029	0.004	0.049	0.680
Aspartate	3.16	3.49	3.34	3.46	0.056	0.522	0.054	0.355
Cysteine	0.26	0.28	0.31	0.32	0.018	0.177	0.747	0.927
Glutamate	3.92	4.02	3.82	4.01	0.175	0.863	0.684	0.893
Glycine	1.93	2.06	2.11	2.13	0.031	0.062	0.243	0.335
Proline	2.50	2.64	2.72	2.73	0.026	0.006	0.183	0.203
Serine	1.37	1.58	1.42	1.55	0.025	0.894	0.002	0.449
Tyrosine	1.97	2.23	2.14	2.25	0.024	0.070	0.001	0.117

381 SEM, standard error of mean. Each value is the average of 10 replicates.

382

Table 5. Effects of substrate (SUB) and development stage (STG) on fatty acid composition of black soldier fly.

Treatments	Food waste		Tofu by-product		SEM	P-value		
	Larvae	Prepupae	Larvae	Prepupae		SUB	STG	SUB×STG
Fatty acid, %								
C10:0	0.88	0.91	0.94	0.97	0.009	0.006	0.160	0.827
C12:0	22.22	23.28	23.72	24.90	0.174	<0.001	0.004	0.871
C14:0	4.38	4.63	4.71	4.68	0.042	0.035	0.238	0.111
C14:1	0.12	0.10	0.16	0.15	0.010	0.032	0.437	0.814
C15:0	0.14	0.16	0.16	0.20	0.008	0.051	0.033	0.454
C16:0	15.43	16.36	14.64	14.35	0.335	0.050	0.641	0.373
C16:1	3.25	3.31	3.29	3.25	0.017	0.882	0.730	0.125
C:17:0	0.22	0.26	0.19	0.23	0.007	0.065	0.015	0.861
C18:0	1.90	2.27	2.03	2.14	0.045	0.985	0.016	0.160
C18:1	24.10	23.41	23.89	23.06	0.268	0.607	0.174	0.893
C18:2n6	19.25	17.04	17.86	17.51	0.328	0.496	0.065	0.170
C18:3n3	2.16	2.11	2.23	2.25	0.022	0.036	0.811	0.392
C18:4	0.90	0.91	0.70	0.74	0.007	<0.001	0.105	0.527
C20:1	0.24	0.27	0.45	0.47	0.010	<0.001	0.345	0.811
C20:4n6	0.26	0.21	0.22	0.21	0.009	0.290	0.067	0.252
C20:5n3	1.36	1.28	1.23	1.31	0.017	0.121	0.980	0.028
Other	2.01	2.14	2.35	2.30	0.049	0.016	0.685	0.374
SFA	45.17	47.86	46.39	47.46	0.400	0.612	0.030	0.325
USFA	53.64	50.77	52.38	51.25	0.417	0.647	0.026	0.311