

RESEARCH

Open Access

Effects of *Prunella vulgaris* labiatae extract on specific and non-specific immune responses in tilapia (*Oreochromis niloticus*)

Kwan-Ha Park and Sanghoon Choi*

Abstract

We examined the effects of *Prunella vulgaris* Labiatae (*P. vulgaris* L.) on specific and non-specific immune responses of Nile tilapia, *Oreochromis niloticus*. The optimal concentration without toxicity of *P. vulgaris* was determined to 30-40 µg/ml *in vitro* and 120 µg/100 g of fish *in vivo*. *P. vulgaris* significantly elicited an antibody titer compared to FCA or β-glucan. β-glucan plus *P. vulgaris* group synergistically enhanced antibody production. No significant difference in antibody production was observed between *P. vulgaris* and *P. vulgaris* plus β-glucan group. A respiratory burst activity of head kidney (HK) leucocytes of tilapia administered with 300 or 500 µg *P. vulgaris* was significantly ($p < 0.05$) enhanced compared with the PBS-injected control group and FCA-treated group. Maximum increase in the NBT reduction value was observed in 500 µg *P. vulgaris* group but no significant difference was found between 300 and 500 µg *P. vulgaris* group. The level of serum lysozyme activity was significantly ($p < 0.05$) higher in the 300 and 500 µg *P. vulgaris* than 100 µg *P. vulgaris* and FCA group. The phagocytic activities of HK leucocytes from tilapia administered with 300 and 500 µg *P. vulgaris* were significantly ($p < 0.05$) higher than 100 µg *P. vulgaris* and the control group. *P. vulgaris* was revealed with a good immunoadjuvant evoking the specific and non-specific immune responses of tilapia.

Keywords: *Prunella vulgaris*, Respiratory burst activity, Lysozyme activity, Tilapia

Background

One of the most promising methods for controlling diseases in aquaculture is strengthening the defense mechanisms of fish through prophylactic administration of immunostimulants [1]. Immunostimulants are naturally occurring compounds that modulate the immune system by increasing host resistance to infectious pathogens, and they have been widely used in aquaculture [2-4]. Traditional disease control strategies employing antibiotics and chemical disinfectants are no longer recommended due to the emergence of bacterial resistance as well as concerns over the environment and wildlife protection. Although vaccination has been shown to be an effective prophylactic method for disease control in fish [5], there are some methodological problems related to high costs and stress [6]. Already, remarkable success has been achieved using

immunostimulants as a more environmentally friendly approach to disease management [7-9].

Several compounds, including β-glucans, chitin, algal and mistletoe extracts, and bacterial polysaccharides, have been used to enhance immunity and disease resistance in a variety of fish species [7,10-12]. β-glucan administration has been reported to augment antibody production, complement activity, lysozyme activity, phagocytic activity, and respiratory bursts in channel catfish *Ictalurus punctatus* [13], Atlantic salmon *Salmo salar* [14], rainbow trout *Oncorhynchus mykiss* [15], gilthead rainbow trout *Sparus auratus* [16], and sea bass *Dicentrarchus labrax* [17].

Prunella vulgaris is a perennial herb that is used in traditional medicine for the clinical treatment of sore throat, fever, and accelerated wound healing [18,19]. The organic fraction of *P. vulgaris* exhibits antioxidative and antimicrobial activities [18], whereas aqueous extracts of *P. vulgaris* inhibit HIV-1 infection [20]. In aquaculture, *P. vulgaris* was reported to have efficacy as a dietary supplement, although only up-regulation of natural immunity was observed [21].

* Correspondence: shchoi@kunsan.ac.kr
Department of Aquatic Life Medicine, Kunsan National University, Gunsan,
Geonbuk 573-400, South Korea

To further study the availability of *P. vulgaris* as an immunoadjuvant to elicit a vaccination effect, we measured the specific antibody titer following intraperitoneal injection of *P. vulgaris* with an antigen in tilapia as a fish model.

To overcome the disease problem affecting fish culture systems, the present study applied indigenous *P. vulgaris* as an appropriate immunoadjuvant in order to augment specific and non-specific immunities as well as disease resistance in fish.

Materials and methods

Fish

Nile tilapia, *Oreochromis niloticus*, fish weighing about 100-150 g each were obtained from a fish farm in Kunsan National University, Korea and acclimated for 2 weeks to laboratory conditions in 70 L glass aquaria containing re-circulated and aerated water at 23-23°C. They were acclimated to this environment for at least 2 weeks prior to use and fed daily using a commercial diet during the adaptation and experimental periods. The health status of the animals was checked daily by observing fish behavior, and there were no clinical symptoms such as abnormal swimming patterns or body color changes.

Reagents

Nitroblue tetrazolium (NBT), Percoll, hemocyanin (HC), 3-(4,5-dimethylthiazol-2yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT), phorbol myristate acetate (PMA), and Minimum essential medium (MEM) were purchased from Sigma Chemicals CO. Hanks balanced salt solution (HBSS), fetal bovine serum (FBS), and antibiotic-antimycotic were obtained from Gibco BRL, Grand Island, NY. Sodium nitrite, sulfanilamide, and phosphoric acid were purchased from ICN Biomedicals. Bakers' yeast and *Saccharomyces cerevisiae* purchased from Oriental Yeast Co. Ltd., Tokyo, and thioglycollate broth was obtained from Difco Laboratories, Detroit, USA.

Extraction of *P. vulgaris*

P. vulgaris originating from South Korea was kindly donated by an herbal medicine company (Sanyacho-Nongwon, Namyangju, Korea). Extraction of *P. vulgaris* was performed according to Lee's method [22]. Briefly, the chopped flowers, stems, and leaves of *P. vulgaris* (100 g) were placed in distilled water (1,000 ml) and stirred at 4°C overnight. After centrifugation at 15,000 × *g* for 20 min, the supernatant was filtered through 0.2 μm pore-sized filters. Protein content of *P. vulgaris* extract was determined using a commercial protein assay kit (Bio-Rad Lab, USA) and stored at 4°C.

In vitro and *in vivo* toxicities of *P. vulgaris*

The *in vitro* toxicity of *P. vulgaris* was tested against EPC and CHSE-214 fish cell lines. Respective cell lines

(1 × 10⁶ cells) were dispensed into each well of a 24-well plate (Costar, USA), followed by administration of various concentrations of *P. vulgaris* extract ranging from 10 ng to 100 μg/ml. After incubation at 24°C for 3 days, MTT assay was performed according to the method of Daly et al. [23]. Briefly, the tissue culture plates were centrifuged at 500 × *g* for 10 min, after which the supernatant fluids were carefully removed without disturbing the cell pellet or formazan precipitate. The formazan crystals were then dissolved by addition of 200 μl of dimethyl sulphoxide (DMSO) (Sigma) to each well, followed by 25 μl of glycine buffer (0.1 M glycine, 0.1 M NaCl, pH 10.5). Contents of the wells were then thoroughly mixed with a multichannel pipette. After 10 min, formazan development was read at 595 nm using an ELISA reader (ASYS HITECH, Austria).

To determine whether or not *P. vulgaris* has serious toxicity *in vivo*, 100 and 1000 μg of *P. vulgaris*/100 g of fish were intraperitoneally (I.P.) injected into seven fish per group. Fish blood was collected 4 days after injection, and the concentrations of glutamic oxaloacetic transaminase (GOT), glutamic pyruvate transaminase (GPT), and c-creatin were determined on a Fuji Dry Chem System (Fuji Photo Film Co. Ltd, Japan).

Administration of *P. vulgaris* to elicit non-specific immune response in tilapia

Tilapia were divided into five groups of seven fishes per group. Fish in each group were I.P injected with 100, 300, and 500 μg of *P. vulgaris*/100 g of fish in 0.5 ml of phosphate-buffered saline (PBS). The remaining group of fish was injected with an equivalent volume of sterile PBS or 1:1 emulsified Freund's complete adjuvant (FCA) (Sigma) as a control. On day 4 post-injection, blood and head kidney leucocytes were obtained from each fish.

Antibody production upon administration of *P. vulgaris* plus other immunostimulants

The immunostimulating effect of *P. vulgaris* was compared with those of FCA and β-glucan based on elevated antibody production. HC was used as an antigen to evoke a specific antibody response. For I.P. injection, 300 μg of *P. vulgaris* and 100 μg of HC were mixed and administered in a volume of 200 μl. Tilapia were divided into six groups (five fish per group), after which HC was injected alone (control) or mixed and injected with 100 μl of other adjuvants. The total injection volume was adjusted to 200 μl in all experiments. *P. vulgaris* was I.P. injected at a concentration of 300 μg suspended in 200 μl of phosphate-buffered saline (PBS). In the FCA group, 100 μl of FCA was 1:1 emulsified with HC suspended in 100 μl of PBS. In the FCA plus *P. vulgaris* group, 100 μl of FCA was added to the *P. vulgaris* and HC mixture and then I.P. injected at a volume of 200 μl. The optimized concentration (50 μg) of β-glucan was injected with HC at a volume

of 200 μ l. In the β -glucan plus *P. vulgaris* group, 50 μ g of β -glucan and 300 μ g *P. vulgaris* were mixed together and I.P. injected at a volume of 200 μ l. On day 30 post-injection, blood was harvested from the fishes in each group, followed by antibody titer assay using an ELISA reader.

Isolation of head kidney (HK) leucocytes

The method described by Santarem et al. [24] was followed with some modifications. The tilapia HK was dissected out by ventral incision, cut into small fragments, and then transferred into 5 ml of HBSS. Cell suspensions of the HK were obtained by teasing HK tissues with two slide glasses in HBSS in a Petri dish (Corning, USA). After sedimentation of tissue debris at 4°C for 1 min, the supernatants were removed. HK cell suspensions were then layered over a 34-51% Percoll gradient and centrifuged at 1000 \times g for 40 min at 14°C. After centrifugation, leucocyte bands located above the 34-51% interfaces were collected using a Pasteur pipette and washed twice at 120 \times g for 8 min in HBSS. The concentration of viable cells was determined by trypan blue exclusion.

Serum

Blood was collected from the dorsal aorta of tilapia. Blood was allowed to clot at 20°C for 30 min and then cooled at 0°C for 1 h. Serum was obtained by centrifugation at 1000 \times g for 8 min. Sera were frozen at -20°C until used.

Reactive oxygen intermediates (ROI) production assay

ROI production by tilapia kidney cells after administration of *P. vulgaris* was assessed by monitoring reduction of NBT [25]. Leucocytes (1×10^5 cells) were washed once with HBSS at 60 \times g for 3 min at 4°C and then incubated in 100 μ l of complete media in the presence of PMA and 1 μ g/ml of NBT. After 1 h of incubation at 25°C, excess NBT was washed out with PBS, and the leucocytes were fixed with 70% methanol. After discarding the methanol, the leucocytes were washed twice with PBS. The reduced formazan was then solubilized with 120 μ l of KOH and 140 μ g of DMSO, after which optical density values were read at 620 nm on an ELISA reader.

Lysozyme activity

Serum lysozyme activity was measured using a modified turbidimetric microtiter plate technique according to Ellis [6]. Briefly, a standard suspension of 0.15 mg/ml of *Micrococcus lysodeikticus* (Sigma) was prepared in 66 mM phosphate buffer (pH 6.0). Tilapia serum (50 μ l) was then added to 1 ml of the bacterial suspension, after which the absorbance reduction was recorded at 0.5 and 4.5 min intervals at 450 nm on a spectrophotometer (SHIMADZU

UV-1600PC). One unit of lysozyme activity was defined as a reduction in absorbance of 0.001/min.

Phagocytic activity

Tilapia HK leucocytes were adjusted to 1×10^6 cells/200 μ l/well in 5% FBS-MEM and dispensed in an 8-well slide chamber (Nunc, Denmark), followed by overnight incubation at 25°C. Following incubation, 1×10^7 cells/ml of zymosan (Sigma) was added. The mixture was incubated at 25°C for 1 h with occasional shaking, after which 50 μ l of the mixture was smeared onto a glass slide, air-dried, and stained with Wright's solution. Phagocytic activity (PA) [26] was calculated by enumerating 500 leucocytes per fish under a microscope. PA = number of cells ingesting zymosans/number of cells observed \times 100.

Statistical analysis

Statistical significance of the differences between the groups was calculated by applying Student's 2-tailed *t*-test.

Results and discussion

Prunella vulgaris is a perennial plant known for its self-healing properties in Western herbal medicine [27,28], and it traditionally has been used for treating various diseases such as an allergies and inflammation in East Asian countries [29]. In addition, *P. vulgaris* has been reported to have immunomodulatory effects such as activation of macrophages [27,28,30]. The effect of *P. vulgaris* on fish immunity in aquaculture has only been reported by Harikrishnan et al. [21]. Specifically, they investigated the dietary effects of *P. vulgaris* on the non-specific immune response as well as disease resistance against *Uronema marinum*. In the




Figure 1 *In vitro* cytotoxicity of *P. vulgaris* on EPC and CHSE-214 fish cell lines. The cells (1×10^6) from each cell line were incubated with *P. vulgaris* ranging from 10 ng to 100 μ g/ml and incubated at 24°C for 3 days. Formazan development was read at 595 nm using ELISA reader (ASYS HITECH, Austria). Error bars represent SD from the mean of triplicate wells. The result is a representative of three experiments.

Table 1 The levels of GOT, GPT and c-creatin in tilapia sera following injection of *P. vulgaris*

Standards for toxicity	Doses of <i>P. vulgaris</i> injected in fish		
	PBS	P100/100 ¹	P1000/100
GOT	19 ± 3 mg/ML	23 ± 5 mg/ML	25 ± 3 mg/ML
GPT	40 ± 6 mg/ML	39 ± 5 mg/ML	42 ± 6 mg/ML
c-creatin	31 ± 4 mg/ML	28 ± 5 mg/ML	33 ± 7 mg/ML

¹µg of *P. vulgaris*/g of fish.

present study, we tested the availability of *P. vulgaris* as a potent immunoadjuvant to achieve an enhanced vaccination effect. Furthermore, the *in vitro* and *in vivo* toxicities of *P. vulgaris* were investigated in tilapia as a fish model. Lastly, to determine the optimal amount of *P. vulgaris* extract that evokes an immune response in tilapia, we administered *P. vulgaris* extract by intraperitoneal (I.P.) injection.

First, we tested the *in vitro* and *in vivo* toxicities of *P. vulgaris* against transformed fish cell lines and tilapia. The half-killing concentrations of *P. vulgaris* against EPC and RTG-2 cells were 30 and 40 µg/ml, respectively (Figure 1). These *in vitro* toxicities were similar to a previous study in which Korean mistletoe showed negligible toxicity on mammalian cell lines [31]. Table 1 shows the levels of GOT, GPT, and c-creatin in blood from tilapia sensitized with *P. vulgaris*. There were no significant differences in toxicity between the groups administered 100 and 1000 µg of *P. vulgaris*/100 g of fish. Further, the GOT, GPT, and c-creatin levels of the PBS control group were within normal ranges, indicating non-toxicity. In contrast, a previous study showed that a 10-fold greater

concentration of mistletoe injected into eel significantly augmented toxicity [11]. The immunomodulatory effect of *P. vulgaris* was compared with those of FCA and β-glucan based on HC-specific antibody production. As shown in Figure 2, *P. vulgaris* induced significantly stronger antibody production than either FCA or β-glucan. Although β-glucan failed to elicit efficient antibody production, β-glucan plus *P. vulgaris* synergistically enhanced antibody production. However, no significant difference in toxicity between the *P. vulgaris* and *P. vulgaris* plus β-glucan groups was observed, indicating that *P. vulgaris* alone has strong immunoadjuvant activity.

Phagocytes produce respiratory bursts as a form of attack against invasive pathogens. As such, they are common measure of the defense ability against pathogens, although excessive accumulation of reactive oxygen intermediates (ROIs) is extremely toxic to host cells [32]. ROIs such as superoxide (O₂⁻), hydrogen peroxide (H₂O₂), hydroxyl radical (OH), and singlet oxygen play important roles in the antimicrobial activity of phagocytic cells [33]. As shown in Figure 3, ROI production was significantly (*p* < 0.05) up-regulated in HK leucocytes from tilapia injected with 300 and 500 µg of *P. vulgaris*/100 g of fish compared to both the control and FCA groups, suggesting that ROIs are an indicator of *P. vulgaris*-induced non-specific immunity in tilapia. Although ROI production in the 100 µg of *P. vulgaris* group was higher than that in the control group, the difference was not significant. Maximum NBT reduction value was observed in the 500 µg of *P. vulgaris* group, but no significant difference was observed between the 300 and 500 µg of *P. vulgaris* groups. On the other hand, injection of more than 500 µg of






Figure 2 Antibody titer induced by *P. vulgaris* administered in tilapia. Six groups (5 fish/group) of fish were I.P. immunized with hemocyanin supplemented with *P. vulgaris* (PV), FCA, *P. vulgaris* + FCA (PV + FCA), β-glucan, *P. vulgaris* + β-glucan (PV + glucan) and PBS as a control, respectively. Data represent the mean + S.D. (n = 5). Statistical differences (*p* < 0.05) between groups are indicated by different letters over the bar.



P. vulgaris/100 g of fish failed to up-regulate ROI production (data not shown). Usually, the effect of immunostimulants is strongest at intermediate dosages with minimal activity and even toxicity at high doses [34,35]. This phenomenon has been established in fish through *in vivo* [36,37] and *in vitro* studies [38].


Lysozyme has both bactericidal as well as opsonin effects that activate the complement system and phagocytes to prevent infection and disease [39]. Figure 4 shows lysozyme activities in the sera of tilapia with or without *P. vulgaris* administration. Serum lysozyme activities were significantly ($p < 0.05$) higher in the 300 and 500 µg of *P. vulgaris* groups compared to the 100 µg of *P. vulgaris* and FCA groups. Further, there was no significant difference in



lysozyme activity between the 300 and 500 µg of *P. vulgaris* groups. Considering that serum from *P. vulgaris*-injected fish showed elevated lysozyme activity, *P. vulgaris* is likely to play a critical role in evoking lysozyme activity from tilapia kidney phagocytes. However, injection of 1000 µg of *P. vulgaris* reduced lysozyme activity compared to the 300 and 500 µg of *P. vulgaris* groups (data not shown), suggesting that an excess concentration of *P. vulgaris* interferes with lysozyme activity in tilapia.

Phagocytes are the first cells to recognize invading foreign bodies and are thus central to cell-mediated and humoral immunities [40]. To test whether or not *P. vulgaris* can influence phagocytic activity, tilapia kidney leucocytes sensitized with *P. vulgaris* (100, 300, and 500 µg of *P. vulgaris*) were incubated overnight with zymosans. In our study, instead of foreign pathogens, zymosans were treated to phagocytes from either *P. vulgaris*-injected tilapia or non-treated tilapia. As shown in Figure 5, the phagocytic activities of HK leucocytes isolated from tilapia injected with 300 and 500 µg of *P. vulgaris* were significantly higher compared to the 100 µg of *P. vulgaris* and PBS control groups. Further, was a significant difference ($p < 0.05$) between the 100 µg of *P. vulgaris* as well as 300 or 500 µg of *P. vulgaris* groups, but no significant difference between the 300 and 500 µg of *P. vulgaris* groups themselves. Lastly, excess injection of *P. vulgaris* (1000 µg) inhibited phagocytosis and respiratory bursts in HK leucocytes isolated from tilapia.

Although the materials used in the study were different, Gopalakannan and Aurl [41] and Luo et al. [42] also reported that fish treated with a high dosage of chitosan display significantly inhibited phagocytosis compared to low dosage. This result suggests that the high level of *P. vulgaris* directly induced phagocytosis, thereby exhausting the cells.



Conclusions

P. vulgaris elevated almost all non-specific immune parameters as well as specific humoral immunity. Therefore, *P. vulgaris* could be a promising immunomodulatory material for inducing specific and non-specific immune responses in fish. Further studies on using *P. vulgaris* as a dietary supplement in aquaculture are currently underway.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KH collected *Prunella vulgaris* samples, wrote the manuscript and performed *in vitro* toxicity test as well as a statistical analysis. SH carried out the specific and non-specific immune-related experiments. Both authors read and approved the final manuscript.

Acknowledgments

This work was supported by a grant from the Fisheries Science Institute in Kunsan National University in 2013.

Received: 7 April 2014 Accepted: 7 April 2014

Published: 15 May 2014

References

1. Robertsen B: Modulation of the non-specific defense of fish by structurally conserved microbial polymers. *Fish Shellfish Immunol* 1999, **9**:269–90.
2. Ai QH, Mai KS, Zhang I, Tan BP, Zhang WB, Xu W: Effects of dietary β -1,3 glucan on innate immune response of large yellow croaker, *Pseudosciaena crocea*. *Fish Shellfish Immunol* 2007, **22**:394–402.
3. Selvaraj V, Sampath K, Sekar V: Administration of lipopolysaccharide increased specific and non-specific immune parameters and survival in carp infected with *Aeromonas hydrophila*. *Aquaculture* 2009, **286**:176–183.
4. Ringø E, Olsen RE, Gifstad TØ, Dalmo RA, Amlund H, Hemre GI, Bakke AM: Prebiotics in aquaculture: a reviews. *Aquacul Nutr* 2010, **16**:117–136.
5. Midtlyng PJ, Reitan LJ, Lillehaug A, Ramstad A: Protection, immune responses and side effects in Atlantic salmon (*Salmo salar* L.) vaccinated against furunculosis by different procedures. *Fish Shellfish Immunol* 1996, **6**:599–613.
6. Ellis AE: Immunity to bacteria in fish. *Fish Shellfish Immunol* 1999, **9**:291–308.
7. Raa J: The use of immunostimulatory substances in fish and shellfish farming. *Rev Fish Sci* 1996, **4**:229–288.
8. Sakai M: Current research status of fish immunostimulants. *Aquaculture* 1999, **172**:63–92.
9. Peddie S, Zou J, Secombes J: Immunostimulation in the rainbow trout (*Oncorhynchus mykiss*) following intraperitoneal administration of Ergosan. *Vet Immunol Immunopathol* 2002, **86**:101–113.
10. Anderson DP: Environmental factors on fish health: immunological aspects. In *The fish immune system: organism, pathogen, and environment*. Edited by Iwama G, Nakanishi T. San Diego, CA: Academic; 1996:289–310.
11. Yoon TJ, Park KH, Choi SH: Korean mistletoe (*Viscum album Coloratum*) extract induces eel (*Anguilla japonica*) non-specific immunity. *Immune Network* 2008, **8**(4):124–129.
12. Park KH, Choi SH: The effect of mistletoe, *Viscum album coloratum*, extract on innate immune response of Nile tilapia (*Oreochromis niloticus*). *Fish Shellfish Immunol* 2012, **32**:1016–1021.
13. Chen D, Ainsworth AJ: Glucan administration potentiates immune defense mechanisms of channel catfish, *Ictalurus punctatus Rafinesque*. *J Fish Dis* 1992, **15**:295–304.
14. Engstad R, Robertsen B, Frivold E: Yeast glucan induces increase in lysozyme and complement-mediated haemolytic activity in Atlantic salmon blood. *Fish Shellfish Immunol* 1992, **2**:287–297.
15. Verlhac V, Gabaudan J, Obach A, Schuep W, Hole R: Influence of dietary glucan and vitamin D on non-specific responses of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 1996, **144**:123–133.
16. Castro R, Couso N, Obach A, Lamas J: Effect of different beta-glucans on the respiratory burst of turbot (*Psetta maxima*) and gilthead rainbow trout (*Sparus aurata*) phagocytes. *Fish Shellfish Immunol* 1999, **9**:529–541.
17. Bagni M, Romano N, Finoia MG, Abelli L, Scapigliate G, Tiscar PG: Short- and long-term effects of a dietary yeast β -glucan (Macrogard) and alginic acid (Ergosan) preparation on immune response in sea bass (*Dicentrarchus labrax*). *Fish Shellfish Immunol* 2005, **18**:311–325.
18. Psotova J, Kolar M, Sousek J, Svagera Z, Vicar J, Ulrichova J: Biological activities of *Prunella vulgaris* extract. *Phytother Res* 2003, **17**:1082–1087.
19. Psotova J, Svobodova A, Kotarova H, Walterova D: Photoprotective properties of *Prunella vulgaris* and rosmarinic acid on human keratinocytes. *J Photochem Photobiol B* 2006, **84**:167–174.
20. Oh C, Price J, Brindley MA, Widrechner MP, Qu L, McCoy JA, Murphy P, Hauck D, Maury W: Inhibition of HIV-1 infection by aqueous extracts of *Prunella vulgaris* L. *Virology* 2011, **23**:8–12.
21. Harikrishnan R, Kim JS, Kim MC, Balasundaram C, Heo MS: *Prunella vulgaris* enhances the non-specific immune response and disease resistance of *Paralichthys olivaceus* against *Uronema marinum*. *Aquaculture* 2011, **318**:61–66.
22. Lee JB, Kang TB, Choi SH, Lee U, Kim AJ, Jeong CJ, Lee HC, Cho YS, Won JG, Lim JC, Yoon TJ: Effect of *Prunella vulgaris* Labiatae extract on innate immune cells and anti-metastatic effect in mice. *Food Sci Biotechnol* 2009, **18**(1):218–222.
23. Daly JG, Moore AR, Olivier G: A colorimetric assay for the quantification of brook trout (*Salvelinus fontinalis*) lymphocyte mitogenesis. *Fish Shellfish Immunol* 1995, **5**:265–273.
24. Santarem M, Novoa B, Figueras A: Effect of beta-glucans on the non-specific immune responses of turbot (*Scophthalmus* L.). *Fish Shellfish Immunol* 1997, **7**:429–437.
25. Secombes CJ, Chung S, Jeffries AH: Superoxide anion production by rainbow trout macrophages detected by the reduction of ferricytochrome C. *Dev Comp Immunol* 1988, **12**:201–206.
26. Sahoo PK, Mukherjee SC: Effect of dietary β -1,3 glucan on immune responses and disease resistance of healthy and aflatoxin B1-induced immunocompromised rohu (*Labeo rohita* Hamilton). *Fish Shellfish Immunol* 2001, **11**:683–695.
27. Fang X, Chang RC, Yuen WH, Zee SY: Immune modulatory effects of *Prunella vulgaris* L. *Int J Mol Med* 2005, **15**:491–496.
28. Fang X, Yu MM, Yuen WH, Xee SY, Chang RC: Immune modulatory effects of *Prunella vulgaris* L. on monocytes/macrophages. *Int J Mol Med* 2005, **16**:1109–1116.
29. Kim SY, Kim SH, Shin HY, Lim JP, Chae BS, Park JS, Hong SG, Kim MS, Jo DG, Park WH, Shin TY: Effects of *Prunella vulgaris* on mast cell-mediated allergic reaction and inflammatory cytokine production. *Exp Biol Med* 2007, **232**:921–926.
30. Han EH, Choi JH, Hwang YP, Park HJ, Choi CY, Chung YC, Seo JK, Jeong HG: Immunostimulatory activity of aqueous extract isolated from *Prunella vulgaris*. *Food Chem Toxicol* 2009, **47**(1):62–69.
31. Yoon TJ, Yoo YC, Kang TB, Her E, Kim K, Azuma I, Kim JB: Cellular and humoral adjuvant activity of lectins isolated from Korean mistletoe (*Viscum album coloratum*). *Int Immunopharmacol* 2001, **1**:881–889.
32. Salinas I, Cuesta A, Esteban MA, Meseguer J: Dietary administration of *Lactobacillus delbrueckii* and *Bacillus subtilis*, single or combined, on gilthead seabream cellular innate immune responses. *Fish Shellfish Immunol* 2005, **19**:67–77.
33. Babior BM: Oxidants from phagocytes: agents of defense and destruction. *Bolld* 1984, **64**:959–966.
34. Bliznakov EG, Adler AD: Nonlinear response of the reticuloendothelial system upon stimulation. *Pathol Microbiol* 1972, **38**:393–410.
35. Gialdroni-Grassi G, Grassi C: Bacterial products as immunomodulating agents. *Int Arch Allergy Appl Immunol* 1985, **76**(Suppl.1):119–127.
36. Kenyon CJ, McKeever A, Oliver JA, Henderson IW: Control of renal and adrenocortical function by the renin-angiotensin system in two euryhaline teleost fishes. *Gen Comp Endocrinol* 1985, **58**:93–100.
37. Anderson DP, Jeney G: Immunostimulants added to injected *Aeromonas salmonicida* bacterin enhance the defense mechanisms and protection on rainbow trout (*Oncorhynchus mykiss*). *Vet Immunol Immunopathol* 1992, **34**:379–389.
38. Siwicki AK, Cossarini-Dunier M, Studnicka M, Demael A: In vivo effect of the organophosphorus insecticide trichlorphon on immune response of carp (*Cyprinus carpio*). II. Effect of high doses of trichlorphon on nonspecific immune response. *Ecotoxicol Environ Saf* 1990, **19**:99–105.
39. Alexander JB, Ingram GA: Noncellular non-specific defense mechanisms of fish. *Ann Rev Fish Dis* 1992, **2**:249–79.

40. Cavaillon JM: Cytokines and macrophages. *Biomed Pharmacother* 1994, **48**:445–453.
41. Gopalakannan A, Aurl V: Immunomodulatory effects of dietary intake of chitin, chitosan and levamisole on the immune system of *Cyprinus carpio* and control of *Aeromonas hydrophila* infection in ponds. *Aquaculture* 2006, **255**:179–187.
42. Luo L, Cai X, He C, Xue M, Wu X, Cao H: Immune response, stress resistance and bacterial challenge in juvenile rainbow trouts *Oncorhynchus mykiss* fed diets containing chitosan-oligosaccharides. *Curr Zool* 2009, **55**:416–22.

doi:10.1186/2055-0391-56-3

Cite this article as: Park and Choi: Effects of *Prunella vulgaris* labiatae extract on specific and non-specific immune responses in tilapia (*Oreochromis niloticus*). *Journal of Animal Science and Technology* 2014 **56**:3.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

