

Dietary turmeric (*Curcuma longa* L.) supplementation improves growth performance, short-chain fatty acid production, and modulates bacterial composition of weaned piglets

Neeraja Recharla¹, Balamuralikrishnan Balasubramanian¹, Minho Song², Pradeep Puligundla³, Soo-ki Kim⁴, Jin Young Jeong⁵ and Sungkwon Park^{1*}

¹Department of Food Science and Biotechnology, Sejong University, Seoul 05006, Korea

²Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Korea

³Department of Food Science and Biotechnology, Gachon University, Seongnam 13120, Korea

⁴Department of Animal Science and Technology, Konkuk University, Seoul 05029, Korea

⁵National Institute of Animal Science, RDA, Wanju 55365, Korea



Received: Feb 10, 2021
Revised: Mar 3, 2021
Accepted: Mar 10, 2021

*Corresponding author

Sungkwon Park
Department of Food Science and
Biotechnology, Sejong University, Seoul
05006, Korea.
Tel: +82-2-3408-2906
E-mail: sungkwonpark@sejong.ac.kr

Copyright © 2021 Korean Society of
Animal Sciences and Technology.
This is an Open Access article
distributed under the terms of the
Creative Commons Attribution
Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted
non-commercial use, distribution, and
reproduction in any medium, provided
the original work is properly cited.

ORCID

Neeraja Recharla
<https://orcid.org/0000-0001-9343-7436>
Balamuralikrishnan Balasubramanian
<https://orcid.org/0000-0001-6938-1495>
Minho Song
<https://orcid.org/0000-0002-4515-5212>
Pradeep Puligundla
<https://orcid.org/0000-0002-2457-0273>
Soo-ki Kim
<https://orcid.org/0000-0003-3499-3330>
Jin Young Jeong
<https://orcid.org/0000-0002-8670-7036>

Abstract

In livestock nutrition, natural feed additives are gaining increased attention as alternatives to antibiotic growth promoters to improve animal performance. This study investigated the effects of dietary turmeric supplementation on the growth performance and gut health of weaned piglets. A total of 48 weaned piglets (Duroc × [Landrace × Yorkshire]) were used in a 6-week feeding trial. All piglets were allotted to two dietary treatments: corn-soybean meal basal diet without turmeric (control) and with 1% weight per weight (w/w) turmeric powder (turmeric). The results showed that dietary inclusion of turmeric with the basal diet improved final body weight and total average daily gain ($p < 0.05$). The concentrations of short-chain fatty acids in the fecal samples, including acetic, butyric, and propionic acids, were higher in the turmeric group ($p < 0.05$). The villus height-to-crypt depth ratio was higher in the ileum of turmeric-fed piglets ($p = 0.04$). The 16S rRNA gene sequencing of fecal microbiota indicated that, at the phylum level, *Firmicutes* and *Bacteroidetes* were the most predominant taxa in all fecal samples. *Bacteroidetes* were significantly decreased in the turmeric group compared to the control group ($p = 0.021$). At the genus level, turmeric showed a decreased abundance of *Prevotella* ($p = 0.021$) and an increasing trend of *Lactobacillus* ($p = 0.083$). Among the total detected species, nine bacterial species showed significant differences between the two groups. The results of this study indicated that turmeric altered the gut microbiota and short-chain fatty acid production. This suggests that turmeric could be used as a potential alternative growth promoter for piglets.

Keywords: Weaned piglets, Turmeric, Gut health, Gut microbiota, Growth promoters, 16S rRNA sequencing

Sungkwon Park
<https://orcid.org/0000-0002-7684-9719>

Competing interests

Not applicable.

Funding sources

This work was carried out with the support of "Cooperative Research Program for Agriculture Science & Technology Development (Project No. PJ015002032020)," Rural Development Administration and Basic Science Research Foundation through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future planning (2019R1F1A1059043), Korea.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

conceptualization: Recharla N, Park S.
Data curation: Recharla N, Balasubramanian B.
Formal analysis: Recharla N, Jeong JY.
methodology: Kim S, Jeong JY.
validation: Song M, Puligundla P.
writing-original draft: Recharla N
writing-review & editing: Song M, Kim S, Jeong JY, Park S.

Ethics approval and consent to participate

Animal care procedures and experimental protocols were approved by the Animal Care and Use Committee of Chungnam National University (Approval# CNU-00611).

INTRODUCTION

Antibiotics have been widely used as growth promoters in livestock to improve animal performance and profitability by improving feed efficiency and animal growth, as well as reducing the incidence of diseases [1]. In particular, nursery pigs are supplemented with antibiotics for disease prevention or to reduce morbidity and mortality [2]. On the other hand, extensive use of antibiotics promotes antibiotic resistance, which could have a negative impact on both animal and human health. Antibiotics used for animal growth are closely related to the class of antibiotics used in human medicine to treat foodborne infections, including penicillin, aminoglycosides, and tetracyclines [3]. The use of antimicrobials in food-producing animals leads to multidrug resistance in both animals and humans [3,4]. Therefore, several countries have banned or placed restrictions on antibiotic use in animal feed [4–7]. Thus, it is necessary to adopt alternative approaches for antibiotic use. During post-weaning, various changes take place in the swine due to stress, new diet, and other factors, which cause increased invasion and colonization of pathogenic bacteria, resulting in infection and diarrhea [8,9]. The immune system is not mature enough in piglets to fight invading pathogens; hence, the post-weaning period is a critical time for maintaining animal health and performance. The gut microbiome plays a major role in immune system development, maintaining nutrient metabolism, performance, disease defense, and health status of the host [10]. Intestinal microbiota might be a potential novel strategy to modulate the general immune system and gut health [11,12]. To modulate the gut microbiota to exert beneficial effects on the host, most researchers have employed prebiotics, probiotics, essential oils [13–15], dietary enzymes, natural herbs, and medicinal plants [16] or phytobiotics [17–19].

Turmeric, also known as the golden spice, is a popular medicinal herb derived from *Curcuma longa* Linnaeus rhizomes. Turmeric plays a vital role in traditional medicinal purposes as an antimicrobial and anti-carcinogenic agent [20,21]. It contains approximately 69.4% carbohydrates, 5.1% fat, 6.3% protein, 3.5% minerals, and 13.1% moisture [22,23]. The turmeric rhizome contains a major fraction of starch (47%–56% w/w) on a dry basis [24,25]. Isolated turmeric starch contains 48%–50% (w/w) amylose [25]. Indigestible carbohydrates, such as resistant starch and other carbohydrates, are fermented by microbes in the large intestine and produce short-chain fatty acids (SCFAs) and other products. The main bioactive compounds in turmeric are curcuminoids, which constitute 1%–6% of the dry weight of turmeric [26]. The three major curcuminoids are curcumin (80%), desmethoxycurcumin (18%), and bisdemethoxycurcumin (2%) [27,28]. The bioactive compounds of turmeric consist of volatile and non-volatile phytochemicals that are less toxic and have beneficial effects, including antioxidant, antibacterial, anti-inflammatory, antiviral, antifungal, anticarcinogenic, and hypo-cholesteric activities [29–31].

Turmeric has gained attention in recent years as a potential alternative to antibiotic growth promoters in livestock feed. The beneficial effects of dietary inclusion of turmeric on growth performance and digestibility have been reported [32,33]. However, limited studies are available on the influence of turmeric on the gut microbiota of pigs. Furthermore, to the best of our knowledge, there is no metagenomic study on the effects of turmeric supplemented diets on gut microbiota in pigs. Therefore, the objective of this study was to investigate the effect of dietary turmeric supplementation on growth performance, blood parameters, fecal score, fecal SCFAs, branched-chain fatty acids (BCFAs), gut microbiota, and histomorphology of the ileum in weaned piglets.

MATERIALS AND METHODS

Experimental design, animal, diet, and housing

A total of 48 newly weaned piglets (Duroc × [Landrace × Yorkshire]) with an initial average bodyweight (BW) of 7.35 ± 0.3 kg were used in 6-week feeding trial. All piglets were obtained from one farm and weaned at 28 days of age. Animal experiments were performed at the Animal Research Center at Chungnam National University, Daejeon, Korea. Animal care procedures and experimental protocols were approved by the Animal Care and Use Committee of Chungnam National University (Approval# CNU-00611). All piglets were randomly assigned to two dietary treatments: the group fed with basal diet only (control) and the basal diet supplemented with 1% (w/w) turmeric powder (turmeric). Turmeric powder was purchased from a local supermarket in Seoul, Korea. Each dietary treatment had four replicates per treatment, with six piglets per pen. In total, 48 male piglets, 24 piglets in the control group, and 24 piglets in the turmeric group were allotted. Diets in mash form were formulated to meet the requirements suggested by the NRC 2012 [34]. Nutrient composition of the diet and chemical composition of turmeric are shown in Tables 1 and 2, respectively. A general maintenance program was used for sows and piglets during lactation. The diets did not include any antibiotics to avoid antibacterial activity during the lactation

Table 1. Nutrient composition of basal diet fed to experimental piglets

Ingredient (%)	Phase 1 ¹⁾	Phase 2 ²⁾
Corn	31.57	51.56
Soybean meal (44% CP)	18.00	26.56
Soy protein concentrate	16.96	8.00
Dried whey	24.00	10.00
Lactose	4.00	-
Soybean oil	3.00	1.35
Limestone	1.00	1.00
Monocalcium phosphate	0.90	0.90
Vitamin pre-mix ³⁾	0.20	0.20
Mineral pre-mix ⁴⁾	0.20	0.20
L-Lysine-HCl	0.08	0.17
DL-Methionine	0.09	0.07
Total	100	100
Calculated energy and nutrient content		
ME (Mcal/kg)	3.53	3.42
CP (%)	24.49	22.51
Calcium (%)	0.81	0.73
Phosphorus (%)	0.69	0.63
Lysine (%)	1.54	1.41

¹⁾Week 1 to 3 (21 days).

²⁾Week 4 to 6 (21 days).

³⁾Provided per kilogram of diet: vitamin A, 12,000 IU; vitamin D₃, 2,500 IU; vitamin E, 30 IU; vitamin K₃, 3 mg; D-pantothenic acid, 15 mg; nicotinic acid, 40 mg; choline, 400 mg; and vitamin B₁₂, 12 µg.

⁴⁾Fe, 90 mg from iron sulfate; Cu, 8.8 mg from copper sulfate; Zn, 100 mg from zinc oxide; Mn, 54 mg from manganese oxide; I, 0.35 mg from potassium iodide; Se, 0.30 mg from sodium selenite.

The calculation for the energy and nutrient contents was performed using the below formula:

Calculated energy or each nutrient content = sum of (energy or each nutrient value of each ingredient used in a diet × % concentration of each ingredient used in a diet / 100).

ME, metabolizable energy; CP, crude protein.

Table 2. Chemical composition of turmeric powder

Constituents	Quantity (%)
Moisture	10.86
Crude protein	37.39
Crude fat	2.78
Crude fiber	3.11
Crude ash	6.26
Carbohydrates	42.71
Starch	35.91
Neutral detergent fiber (NDF)	12.11
Acid detergent fiber (ADF)	9.68
Soluble dietary fiber (SDF)	2.24
Insoluble dietary fiber (ISDF)	17.37
Cellulose	8.77
β -Glucans	13.04
Lignin	0.91
Hemicellulose	2.43

and experimental periods. All experimental piglets were housed in an environmentally controlled, slatted-floor facility with a mechanical ventilation system. Each pen was equipped with a self-feeder and nipple water to allow ad libitum access to feed and water throughout the experimental period. The piglets were individually weighed at the start and at weeks 3 and 6 of the experimental period, and feed intake was recorded throughout the experiment to calculate average daily gain (ADG), average daily feed intake (ADFI), and the gain-to-feed ratio (G:F) was calculated using ADG and ADFI.

Sample collection

On the final day of the experiment, freshly voided fecal samples from one randomly selected piglet in each pen were collected by rectal stimulation for 16S sequencing and SCFA analysis. The number of samples was determined based on our previous pilot study (data not shown). All samples were stored at -80°C until analysis. Blood samples were collected aseptically through an external jugular vein puncture.

Serum hematological and biochemical indices

All blood samples were centrifuged at $1,200\times g$ for 10 min at 4°C , and plasma and serum samples were sent to Neodin Vet Lab (Seoul, Korea) on the same day of sample collection for analysis of concentrations of total proteins, creatine, urea, glucose, total cholesterol, triglycerides, and gamma-glutamyl transferase. Plasma samples were analyzed for red blood cell (RBS), white blood cell (WBC), and platelet counts.

Volatile fatty acids analysis

Volatile fatty acids (VFA) analysis was performed according to Cho et al. [35] with modifications using gas chromatography (GC) (6890 N, Agilent, Santa Clara, CA, USA), equipped with an HP-INNOWAX column and a flame ionization detector. Fresh fecal samples (1 g) were acidified with 1 mL of 25% phosphoric acid solution, 3 mL of distilled water, and 50 μL saturated mercury solution (Sigma-Aldrich, St. Louis, MO, USA). After 30 min, the samples were centrifuged at $3,000\times g$ for 20 min, and 3 mL of the supernatant was collected. Then, the 3 mL of the supernatant

was centrifuged at 13,800×g for 10 min and filtered through a 0.2 µm filter (Whatman, Uppsala, Sweden). The filtrates were mixed with an equal amount of methanol and then placed in 2.0 mL GC vials (Agilent) to measure the concentration of volatile fatty acids. The sample injection volume was 2 µL, with a split ratio of 10:1.

Histometric analysis of piglet ileum

On the final day of the experimental period, a total of 8 piglets (4 piglets per group) were slaughtered, and the intestinal tract was removed. The distal ileum segments were collected and fixed with 4% paraformaldehyde in 0.01 M phosphate buffered saline. The ileum sections were stained with hematoxylin and eosin for microscopic examination to determine villus height (VH), crypt depth (CD), VH:CD, villus width, villus area, and number of goblet cells.

Occurrence of diarrhea

The diarrhea score of each piglet was recorded at weeks 1 to 6 of the trial. Diarrhea was assessed visually based on consistency of the feces, and fecal scores were determined using the following fecal scoring system: 1 hard, dry pellet; 2 firm, formed stool; 3 soft, moist stool that retains shape; 4 soft, unformed stool that assumes shape of container; 5 watery liquid that can be poured. The fecal score was assessed in a treatment-blinded manner by two trained individuals. Scores were recorded on a pen-basis observation of individual piglets and signs of stool consistency in the pen [36].

DNA extraction and sequencing

Total DNA was extracted from fecal samples using the PowerSoil® DNA Isolation Kit according to the manufacturer's protocol. The quantification of DNA and DNA quality was measured using PicoGreen and Nanodrop (Thermo Scientific, Waltham, MA, USA). The primers used for 16S V3-V4 rRNA gene amplification are listed in Table 3.

Input gDNA (12.5 ng) was amplified with 16S V3-V4 primers, and a subsequent limited-cycle amplification step was performed to add multiplexing indices and Illumina sequencing adapters. Amplicons from PCR were pooled using PicoGreen and used as input for Illumina library preparation. The size of the libraries was verified using the LabChip GX HT DNA High Sensitivity Kit (PerkinElmer, Waltham, MA, USA). Samples were sequenced using an Illumina MiSeq (Macrogen, Seoul, Korea).

Sequence read processing and data analysis

Sequencing reads obtained from Illumina MiSeq were filtered and trimmed using CD-HIT-OUT software and rDNA Tools [37]. To perform taxonomic assignment, operative taxonomic units (OTUs) were selected based on a 97% threshold of sequence similarity using the QIIME-UCLUST program. The filtered reads were clustered and OTUs were generated using CD-HIT-DUP. The sequences that passed from the quality filters were analyzed using the QIIME pipeline, which included features to calculate diversity indices and phylogenetic diversity (PD) rarefaction curves. Alpha-diversity indices including OTUs, Shannon, Chao1, and Simpson index were measured for each sample, and beta-diversity of the two groups were illustrated using principal

Table 3. Primers used for 16S V3-V4 rRNA gene amplification

Direction	Primer
Forward	5'TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG
Reverse	5'GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC

component analysis (PCA) and principal coordinate analysis (PCoA) based on weighted UniFrac distances. The Ribosomal Database Project (RDP) classifier was used for taxonomic assignment of the fecal microbiome of the two groups.

Statistical analysis

Data of hematological and biochemical analyses, VFA concentrations, histometric analysis, and microbial diversity indices were analyzed by Student's *t*-test using the SPSS Statistics Version 23 software package (IBM, Armonk, NY, USA). Growth performance data were statistically analyzed using the GLM procedure of the Statistical Analysis System (SAS Institute, Cary, NC, USA). The Mann-Whitney U test was used to determine the statistical significance of the relative abundance of microbial communities in two groups at the phylum, class, genus, and species levels (SPSS version 23, IBM). Statistical significance was reported at $p < 0.05$, and trends were noted when $0.05 < p < 0.10$.

RESULTS

Effects of turmeric on growth performance and fecal score analysis

In the current study, piglets supplemented with turmeric diet had increased final BW compared to piglets fed the control diet (Table 4). Furthermore, dietary supplementation with turmeric had significant effects on ADFI and tendency effects on G:F at week 3 without affecting ADG. At week 6, increased ADG ($p = 0.026$) and tendency effects on G:F ($p = 0.09$) did not affect ADFI. Overall, turmeric supplementation had increased ($p < 0.04$) ADG and tendency toward G:F ($p = 0.078$) without affecting total ADFI ($p = 0.349$). The diarrhea incidence scores are presented in Table 5. During the experimental period, none of the piglets suffered from diarrhea. Supplementation of turmeric with the basal diet improved the fecal score during week 6 ($p = 0.009$).

Table 4. Growth performance of weaned piglets fed control and turmeric supplemented diets

Items	Control	Turmeric	SEM	<i>p</i> -value
BW (kg)				
Initial	7.33	7.36	0.038	0.392
Wk 3	16.64	17.40	0.275	0.115
Wk 6	24.96	26.08	0.29	0.029
Phase 1 (wk 1–3)				
ADG (g)	443	478	12.87	0.14
ADFI (g)	357	336	5.247	0.042
G:F	1.241	1.424	0.046	0.06
Phase 2 (wk 4–6)				
ADG (g)	394	414	6.945	0.026
ADFI (g)	822	780	17.087	0.158
G:F	0.480	0.532	0.071	0.09
Total				
ADG (g)	419	446	6.943	0.042
ADFI (g)	693	674	15.827	0.349
G:F	0.605	0.664	0.016	0.078

Control, basal diet; Turmeric, basal diet with 1% (w/w%) of turmeric powder; BW, body weight; ADG, average daily gain; ADFI, average daily feed intake; G:F, gain-to-feed ratio.

Table 5. Effects of turmeric supplementation on fecal score of piglets

Items	Control	Turmeric	SEM	p-value
Fecal score ¹⁾				
Week 1	3.29	3.50	0.10	0.22
Week 2	3.43	3.24	0.06	0.08
Week 3	3.29	3.21	0.06	0.22
Week 4	3.40	3.50	0.05	0.03
Week 5	3.41	3.24	0.04	0.08
Week 6	3.51	3.36	0.05	0.009

¹⁾Fecal scores were determined using the following fecal scoring system: 1 hard, dry pellet; 2 firm, formed stool; 3 soft, moist stool that retains shape; 4 soft, unformed stool that assumes shape of container; 5 watery liquid that can be poured.

Effect of dietary turmeric on hematological and biochemical indices

The hematological parameters for the turmeric and control groups are shown in Table 6. Dietary turmeric supplementation did not influence leukocyte, erythrocyte, and thrombocyte counts, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC). There were no significant differences between the two groups ($p > 0.05$). The effects of dietary turmeric on the biochemical variables are summarized in Table 7. Turmeric supplementation did not significantly affect the levels of total, LDL, and HDL cholesterol. Numerically, HDL-cholesterol was higher in the turmeric group, but not significantly ($p = 0.776$).

VFA analysis

As shown in Table 8, dietary turmeric supplementation increased SCFA production ($p < 0.05$). Acetic, propionic, and butyric acids were predominant. The highest acetic acid levels were observed in the turmeric group, followed by propionic and butyric acids. The levels of isobutyric and isovaleric acids were not significantly different between the two groups; however, the concentrations showed

Table 6. Effect of turmeric supplementation on hematological parameters in piglets

Items	Control	Turmeric	SEM	p-value
Leukocytes				
White blood cell (K/ μ L)	19.27	14.69	1.62	0.18
Neutrophil (%)	43.8	31.03	4.51	0.23
Lymphocyte (%)	49.03	61.27	3.98	0.2
Monocyte (%)	4.7	4.77	0.77	0.97
Eosinophil (%)	4.1	2.97	0.66	0.45
Erythrocytes				
Red blood cell (M/ mm^3)	6	5.8	0.14	0.54
Hemoglobin (g/dL)	10.8	10.33	0.29	0.48
Thrombocytes				
Platelet (K/ μ L)	392.67	450.67	101.92	0.81
MCV (fl)	63.6	62.17	0.77	0.41
MCH (pg)	18.03	17.83	0.41	0.84
MCHC (%)	28.43	28.63	0.82	0.92

MCV, mean corpuscular volume; MCH, mean corpuscular hemoglobin concentration; MCHC, mean corpuscular hemoglobin concentration.

Table 7. Effect of turmeric supplementation on biochemical parameters in piglets

Items	Control	Turmeric	SEM	p-value
TP (g/dL)	7.15	5.875	0.67	0.38
ALB (g/dL)	3.7	3.575	0.07	0.418
T.Bil (mg/dL)	0.3	0.2	0.04	0.272
Glucose (mg/dL)	100.25	81	9.0	0.321
BUN (mg/dL)	5.475	4.85	0.39	0.463
Creatinine (mg/dL)	0.735	0.8925	0.05	0.164
γ-GTP (U/L)	45.75	39.5	4.82	0.554
LDH (U/L)	1033.75	793.75	203.81	0.596
Chol (mg/dL)	93.75	97.75	4.03	0.668
TG (mg/dL)	52.25	53.5	2.06	0.787
HDL (mg/dL)	29.675	31.525	2.9	0.776
LDL (mg/dL)	41.9	52.7	4.37	0.244
AST (U/L)	113	108	24.73	0.928
ALT (U/L)	79.75	76.75	3.19	0.678

TP, total protein; ALB, albumin; T.Bil, total bilirubin; BUN, blood urea nitrogen; GTP, glutamyl transpeptidase; LDH, lactic acid dehydrogenase; Chol, cholesterol; TG, triglyceride; HDL, high density lipoprotein; LDL, low density lipoprotein; AST, aspartate amino transferase; ALT, alanine aminotransferase.

Table 8. Effects of turmeric diet on fecal volatile fatty acids (SCFAs and BCFAs) concentration.

Concentration (µg/g)	Control	Turmeric	SEM	p-value
Acetic acid	5.56	9.8	0.97	0.011
Propionic acid	2.5	4.51	0.4	0.001
Butyric acid	1.88	3.87	0.41	0.001
Valeric acid	0.61	0.96	0.08	0.022
Iso butyric acid	0.55	0.74	0.05	0.057
Iso valeric acid	0.83	1.22	0.11	0.057
Total SCFA	10.57	19.15	1.78	0.002
Total BCFA	1.38	1.96	0.16	0.056

SCFA, short-chain fatty acids; BCFA, branched-chain fatty acids.

an increasing tendency in turmeric group ($p = 0.057$).

Effects of dietary treatments on ileum morphology of weaned piglets

Turmeric supplementation had no effect ($p > 0.05$) on VH and CD (Table 9). However, the VH:CD ratios were higher in the turmeric group than in the control group ($p = 0.04$). There was no difference in the surface area and width of villi in the turmeric group compared to the control group.

DNA sequence data and bacterial diversity

A total of 437,000 read bases were obtained from the sequencing of fecal samples from the control and turmeric groups. After filtering and removing low-quality sequences, an average of 39,289 and 41,318 reads were obtained for control and turmeric group samples, respectively. A total of 870 OTUs belonging to bacteria and archaea were identified at the 97% threshold level.

Alpha-diversity analyses, including Shannon, Simpson, and Chao1 indices were analyzed to explore the effect of dietary turmeric on the richness and evenness of gut microbiota. The

Table 9. Effects of turmeric supplementation on ileum morphology

Item	Control	Turmeric	SEM	p-value
Villus height (μm)	396.77	409.70	16.65	0.596
Crypt depth (μm)	281.47	245.33	13.74	0.096
VH:CD	1.42	1.68	0.08	0.040
Villus width (μm)	143.62	169.88	11.65	0.145
Villus area (μm^2)	30,919	36,862	3,571	0.270
Number of goblet cells	15.67	13.83	1.17	0.246

VH:CD, villus height-to-crypt depth ratio.

α -diversity metrics are shown in Figs. 1A, B, and C. According to the data, all diversity indices were not significantly different ($p > 0.05$) between groups, while turmeric administration showed a decreasing tendency in the Shannon index ($p = 0.055$).

α -Diversity rarefaction curves based on observed OTUs, PD whole tree, and sequence for samples between the two groups indicated sufficient sequencing depth (Figs. 2A and B). PCoA based on weighted Unifrac distance showed two clusters containing each sample of both control




Fig. 1. Microbial diversity indices for control and turmeric fed piglets. (A) The Chao value of control and turmeric groups. (B) The Shannon index of control and turmeric groups. (C) The Simpson index of control and turmeric groups.




Fig. 2. α -Diversity rarefaction curves. (A) Rarefaction curves represent the number of sequences per sample against the number of observed OTUs in control and turmeric groups. (B) Rarefaction curves of PD whole tree in control and turmeric groups. OTUs, operative taxonomic units; PD, phylogenetic diversity.

and turmeric groups, except for one sample in each group (Fig. 3).

Taxonomic analysis

The effects of turmeric supplementation on fecal microbial composition were observed at different taxonomic levels. At the phylum level, 10 phyla were observed in each sample. *Bacteroidetes* and *Firmicutes* were the predominant phyla, accounting for 90% of the total relative abundance. The *Firmicutes* abundance ratio was similar in the control (51.8%) and turmeric (52.46%) groups (Fig. 4). *Bacteroides* abundance decreased in turmeric (27.62%) fed piglets than in control (39.61%) fed piglets ($p = 0.021$).

As shown in Fig. 5, Bacteroidia and Clostridia were the predominant classes in both the control and treatment groups. Twelve classes were identified. The relative abundance of Bacteroidia was




Fig. 3. PcoA analysis of control and turmeric groups. Three dimensional plot based on weighted UniFrac distances. PCoA, principal coordinate analysis.




Fig. 4. Bacterial composition and abundance ratio of the fecal microbiota of piglets at phylum level.




Fig. 5. Bacterial composition and abundance ratio of the fecal microbiota of piglets at class level. C, control group; T, turmeric group.

significantly decreased in turmeric-fed piglets ($p = 0.021$). However, the remaining bacterial abundance was not affected by the turmeric diet.

At the genus level, approximately 105 genera were found in fecal samples from both dietary groups. The 15 most abundant genera in the two groups are shown in Fig. 6. Among the abundant genera, eight belonged to *Firmicutes*, 6 from *Bacteroidetes*, and 1 from *Spirochaetes*. The unclassified bacteria at the genus level were higher in the turmeric group than in the control group ($p = 0.043$). The relative abundance of *Prevotella* genera was significantly lower in the turmeric group than in the control group ($p = 0.021$). The *Lactobacillus* genus showed an increasing trend in the turmeric group ($p = 0.083$). The remaining genera did not differ significantly between the two groups. At the




Fig. 6. Bacterial composition and abundance ratio of the fecal microbiota of piglets at genus level. C, control group; T, turmeric group.

Table 10. Effects of turmeric supplementation on fecal microbiota at species level

Phylum	Genus	Species	Abundance ratio (%)		p-value
			Control	Turmeric	
Bacteroidetes	<i>Bacteroides</i>	<i>Bacteroides stercoris</i>	1.28	0.32	0.021
Bacteroidetes	<i>Prevotella</i>	<i>Prevotella shahii</i>	4.33	0.51	0.043
Bacteroidetes	<i>Prevotella</i>	<i>Prevotella timonensis</i>	1.89	0.1	0.021
Bacteroidetes	<i>Prevotella</i>	<i>Prevotella oris</i>	2.01	1.21	0.021
Bacteroidetes	<i>Muribaculum</i>	<i>Muribaculum intestinale</i>	13	6.71	0.083
Firmicutes	<i>Lactobacillus</i>	<i>Lactobacillus reuteri</i>	0.43	3.38	0.021
Firmicutes	<i>Falcatimonas</i>	<i>Falcatimonas natans</i>	0.24	0.59	0.021
Firmicutes	<i>Clostridium</i>	<i>Clostridium bommimense</i>	0.06	0.26	0.021
Firmicutes	<i>Peptococcus</i>	<i>Peptococcus simiae</i>	0	0.2	0.018
Firmicutes	<i>Negativebacillus</i>	<i>Negativibacillus massiliensis</i>	0.05	0.15	0.021
Spirochaetes	<i>Treponema</i>	<i>Treponema berlinense</i>	0.65	2.38	0.059

species level, a total of nine species abundance ratios were statistically significant between the two groups, including *Lactobacillus* spp. (Table 10).

DISCUSSION

Turmeric root powder is commonly used for medicinal purposes and as a spice in traditional cooking. A couple of studies have investigated the effects of turmeric supplementation on the growth performance of piglets [32,33]. However, more studies are needed to explore the effects of turmeric on intestinal morphology and gut microbial communities. Gut microbiota provides not only fuel to colonocytes, but also helps in immune system development and maintenance of intestinal homeostasis [10]. In this study, we describe the impact of dietary turmeric on growth performance, gut morphology, and microbiota in a porcine model. Turmeric inclusion in piglet diets improved ADG and showed higher average final BW and lower feed intake than the control group. Similarly, a previous study indicated that diets containing turmeric powder at 2%, 4%, and 6% improved final live weight and feed conversion ratio compared with basal diet fed pigs [38]. Maneewan et al. [32] reported that the effects of low levels of dietary turmeric supplementation on nursery pigs at doses of 0.05%, 0.10%, and 0.20% did not influence ADFI, ADG, and feed efficiency. The beneficial effects of turmeric on growth performance might be due to the enhanced secretion of amylase, trypsin, chymotrypsin, and lipase enzymes [38]. Similarly, Singh et al. [39] reported that 1% dietary turmeric resulted in increased BW gain in broiler chickens, however, turmeric did not influence the feed efficiency in this study. Tubcharoen et al. [40] reported similar results in growing-finishing pigs. Furthermore, in this study, fecal scores were much higher in the control group. The incidence of diarrhea among piglets was reduced when the diet was supplemented with turmeric.

Hematological and biochemical variables were determined to determine the health and stress status of the animals. There were no significant differences in the values of RBC, WBC, MCV, MCHC, MCH, and platelets. However, all hematological parameters were within the normal range for swine [41]. This indicates that turmeric causes no variations in the hematology of piglets and that none of the animals experienced stress during the experimental period. The serum biochemical variables were not affected by turmeric. The main active compound, curcumin, in turmeric exhibits hypocholesterolemic activity. Curcumin showed beneficial effects in high-fat fed animals [42].

In the present study, turmeric did not influence cholesterol levels in piglets. These results contrast with those of a previous study that showed that curcumin supplementation decreased serum LDL cholesterol levels in weaned piglets [2]. The lipid-lowering effects of turmeric and curcuminoids are associated with the dose and solubility of curcumin. Considering the poor solubility and bioavailability of curcumin, Porn-anek et al. [43] developed a carrier-based turmeric oleoresin using a solid dispersion technique to enhance curcumin solubility. Pigs fed with the newly developed turmeric oleoresin had increased HDL cholesterol and lowered LDL cholesterol, total cholesterol, and triglycerides.

Intestinal metabolites such as SCFAs play a major role in the regulation of gut homeostasis [42]. SCFAs are the end products produced by bacterial fermentation of non-digestible carbohydrates in the colon [13,44–46]. Primary SCFAs are acetic, propionic, and butyric acids, with butyric acid being the main energy substrate for colonic epithelial cells [44,47]. In this study, turmeric diet-fed piglets produced higher levels of SCFAs. Acetic, propionic, and butyric acid concentrations were higher among all the SCFAs. Catabolism of turmeric polysaccharides provides energy for fermentative bacteria such as *Lactobacillus* spp. and *Clostridium* spp., which results in an increase in SCFA production [28]. Similarly, Han et al. [48] reported that *in vitro* fermentation of spent turmeric powder with pig fecal bacteria resulted in higher concentrations of acetate and propionate. Microbial abundance, particularly *Lactobacillus* abundance in the gut, is associated with gut SCFA production. The status of gut health can be determined by intestinal morphology, such as VH, CD, and the VH:CD ratio. Villi plays a major role in increasing nutrient absorption, especially in the small intestine [12,49]. In this study, turmeric-fed piglets showed a higher VH:CD ratio than the control group. Increased VH:CD ratios indicate improved nutrient absorption function [50]. This result is consistent with a previous study that piglets consuming dietary curcumin showed an improved VH:CD ratio [49].

Turmeric has been reported to be an antimicrobial and anti-inflammatory agent [21,22]. Avanço et al. [51] reported that α -turmerone, β -turmerone, and ar-turmerone components of turmeric showed antifungal, antimycotoxigenic, and antioxidant activities. Thus, turmeric could alter microbial communities in the intestine by inhibiting pathogenic bacteria. The taxonomic analysis in this study showed that *Firmicutes* and *Bacteroidetes* were the most abundant phyla in both groups of piglet gut microbiota, as reported in previous studies [52,53]. The abundance ratio of *Bacteroidetes* decreased in turmeric-fed piglets. However, *Firmicutes* phyla were not altered by turmeric. *Bacteroidetes* are gram-negative anaerobic bacteria normally present in the intestinal flora. *Bacteroides* are generally beneficial to the host through their metabolism of dietary polysaccharides; however, *Bacteroidetes* are involved in inflammatory pathology when the gut microbiota is in an imbalanced state [54,55]. Moreover, Zhao et al. [56] reported that fecal *Bacteroides* were negatively correlated with SCFAs and amino acids in mice. We also observed a decreased abundance of *Bacteroidetes* and increased concentrations of SCFAs in turmeric-fed piglets. Hence, low *Bacteroides* and higher SCFAs are predicted to promote the gut health of piglets in the post-weaning period. Turmeric had no impact on the overall microbial diversity and richness, except for the lower trend of the Shannon index. Similarly, Shen et al. [57] reported that oral administration of curcumin tended to decrease microbial diversity and richness with no significant differences. Despite no significant differences in microbial diversity, the abundance of specific bacteria, including *Lactobacillus* and *Prevotella*, were altered in turmeric fed piglets. At the genus level, *Prevotella* genera decreased in the turmeric group, similar to previous studies [57]. Little is known about the role of *Prevotella* in health promotion. Like other bacteria in normal microflora, *Prevotella* spp. act as opportunistic pathogens and have been associated with infections [58,59]. Turmeric increased the abundance of *Lactobacillus* spp. and *Clostridium* spp. and decreased the abundance of *Prevotella* spp. Similarly, Han et al. [48] reported

that spent turmeric fermented with swine microbiota showed increased *Lactobacillus* populations compared to other groups. Kosti et al. [60] also observed higher *Lactobacillus* counts and lower *E. coli* counts in turmeric-fed hens than in the basal diet group. Some studies have suggested that the phenolic compound curcumin in turmeric root powder possesses alterations in gut microbial composition [28]. Moreover, curcumin has been found to improve the barrier function of the intestine by modulating intracellular signaling pathways [61]. The results obtained from this study revealed that dietary turmeric influences gut microbial fermentation and improves gut health by enhancing beneficial bacteria, SCFAs, and gut morphology. However, in the current study, we used whole turmeric root powder as a dietary supplement. Further studies that use turmeric extract or curcumin alone are needed to clarify the microbial alteration effects in the intestine of pigs.

REFERENCES

1. Cheng G, Hao H, Xie S, Wang X, Dai M, Huang L, et al. Antibiotic alternatives: the substitution of antibiotics in animal husbandry? *Front Microbiol.* 2014;5:217. <https://doi.org/10.3389/fmicb.2014.00217>
2. Li J. Current status and prospects for in-feed antibiotics in the different stages of pork production—a review. *Asian-Australas J Anim Sci.* 2017;30:1667-73. <https://doi.org/10.5713/ajas.17.0418>
3. Tollefson L, Karp BE. Human health impact from antimicrobial use in food animals. *Med Mal Infect.* 2004;34:514-21. <https://doi.org/10.1016/j.medmal.2004.07.028>
4. Omonijo FA, Ni L, Gong J, Wang Q, Lahaye L, Yang C. Essential oils as alternatives to antibiotics in swine production. *Anim Nutr.* 2018;4:126-36. <https://doi.org/10.1016/j.aninu.2017.09.001>
5. Thacker PA. Alternatives to antibiotics as growth promoters for use in swine production: a review. *J Anim Sci Biotechnol.* 2013;4:35. <https://doi.org/10.1186/2049-1891-4-35>
6. Brown K, Uwiera RRE, Kalmokoff ML, Brooks SPJ, Inglis GD. Antimicrobial growth promoter use in livestock: a requirement to understand their modes of action to develop effective alternatives. *Int J Antimicrob Agents.* 2017;49:12-24. <https://doi.org/10.1016/j.ijantimicag.2016.08.006>
7. Bengtsson B, Wierup M. Antimicrobial resistance in Scandinavia after a ban of antimicrobial growth promoters. *Anim Biotechnol.* 2006;17:147-56. <https://doi.org/10.1080/10495390600956920>
8. Laine TM, Lyytikäinen T, Yliäho M, Anttila M. Risk factors for post-weaning diarrhoea on piglet producing farms in Finland. *Acta Vet Scand.* 2008;50:21. <https://doi.org/10.1186/1751-0147-50-21>
9. Heo JM, Opapeju FO, Pluske JR, Kim JC, Hampson DJ, Nyachoti CM. Gastrointestinal health and function in weaned pigs: a review of feeding strategies to control post-weaning diarrhoea without using in-feed antimicrobial compounds. *J Anim Physiol Anim Nutr.* 2013;97:207-37. <https://doi.org/10.1111/j.1439-0396.2012.01284.x>
10. Sekirov I, Russell SL, Antunes LCM, Finlay BB. Gut microbiota in health and disease. *Physiol Rev.* 2010;90:859-904. <https://doi.org/10.1152/physrev.00045.2009>
11. Bengmark S. Gut microbiota, immune development and function. *Pharmacol Res.* 2013;69:87-113. <https://doi.org/10.1016/j.phrs.2012.09.002>
12. Liu Y, Chen F, Odle J, Lin X, Jacobi SK, Zhu H, et al. Fish oil enhances intestinal integrity and inhibits TLR4 and NOD2 signaling pathways in weaned pigs after LPS challenge. *J Nutr.* 2012;142:2017-24. <https://doi.org/10.3945/jn.112.164947>

13. Issara U, Park S, Park S. Determination of fat accumulation reduction by edible fatty acids and natural waxes in vitro. *Food Sci Anim Resour.* 2019;39:430-45. <https://doi.org/10.5851/kosfa.2019.e38>
14. Issara U, Park S, Lee S, Lee J, Park S. Health functionality of dietary oleogel in rats fed high-fat diet: a possibility for fat replacement in foods. *J Funct Foods.* 2020;70:103979. <https://doi.org/10.1016/j.jff.2020.103979>
15. Park S, Cho S, Hwang O. Effects of Italian Ryegrass (IRG) supplementation on animal performance, gut microbial compositions and odor emission from manure in growing pigs. *Agronomy.* 2020;10:647. <https://doi.org/10.3390/agronomy10050647>
16. Lee J, Seo HG, Lee CH. Effects of lotus (*Nelumbo nucifera*) leaf hot water extracts on the quality and stability of eggs using ultrasonication treatment during storage. *Food Sci Anim Resour.* 2020;40:1044-54. <https://doi.org/10.5851/kosfa.2020.e81>
17. Fang J, Yan FY, Kong XF, Ruan Z, Liu ZQ, Huang RL, et al. Dietary supplementation with *Acanthopanax senticosus* extract enhances gut health in weanling piglets. *Livest Sci.* 2009;123:268-75. <https://doi.org/10.1016/j.livsci.2008.11.020>
18. Bird AR, Vuaran M, Crittenden R, Hayakawa T, Playne MJ, Brown IL, et al. Comparative effects of a high-amylose starch and a fructooligosaccharide on fecal bifidobacteria numbers and short-chain fatty acids in pigs fed *Bifidobacterium animalis*. *Dig Dis Sci.* 2009;54:947-54. <https://doi.org/10.1007/s10620-008-0451-3>
19. Kiarie E, Romero LF, Nyachoti CM. The role of added feed enzymes in promoting gut health in swine and poultry. *Nutr Res Rev.* 2013;26:71-88. <https://doi.org/10.1017/S0954422413000048>
20. Azuine MA, Bhide SV. Chemopreventive effect of turmeric against stomach and skin tumors induced by chemical carcinogens in Swiss mice. *Nutr Cancer.* 1992;17:73-83. <https://doi.org/10.1080/01635589209514174>
21. Panpatil VV, Tattari S, Kota N, Nimgulkar C, Polasa K. In vitro evaluation on antioxidant and antimicrobial activity of spice extracts of ginger, turmeric and garlic. *J Pharmacogn Phytochem.* 2013;2:143-8.
22. Chattopadhyay I, Biswas K, Bandyopadhyay U, Banerjee RK. Turmeric and curcumin: biological actions and medicinal applications. *Curr Sci.* 2004;87:44-53.
23. Dono ND. Turmeric (*Curcuma longa* Linn.) supplementation as an alternative to antibiotics in poultry diets. *Indones Cent Anim Res Dev* 2018;23:41-93.
24. Leonel M, Sarmiento SBS, Cereda MP. New starches for the food industry: *Curcuma longa* and *Curcuma zedoaria*. *Carbohydr Polym.* 2003;54:385-8.
25. Kuttigounder D, Lingamallu JR, Bhattacharya S. Turmeric powder and starch: selected physical, physicochemical, and microstructural properties. *J Food Sci.* 2011;76:C1284-91. <https://doi.org/10.1111/j.1750-3841.2011.02403.x>
26. Zam W. Gut microbiota as a prospective therapeutic target for curcumin: a review of mutual influence. *J Nutr Metab.* 2018;2018:1367984. <https://doi.org/10.1155/2018/1367984>
27. Gupta SC, Patchva S, Aggarwal BB. Therapeutic roles of curcumin: lessons learned from clinical trials. *AAPS J.* 2013;15:195-218. <https://doi.org/10.1208/s12248-012-9432-8>
28. Peterson CT, Vaughn AR, Sharma V, Chopra D, Mills PJ, Peterson SN, et al. Effects of turmeric and curcumin dietary supplementation on human gut microbiota: a double-blind, randomized, placebo-controlled pilot study. *J Evid Based Integr Med.* 2018;23:2515690X18790725. <https://doi.org/10.1177/2515690X18790725>
29. Ammon HP, Anazodo MI, Safayhi H, Dhawan BN, Srimal RC. Curcumin: a potent inhibitor of leukotriene B4 formation in rat peritoneal polymorphonuclear neutrophils (PMNL).

- 1992;58:226. <https://doi.org/10.1055/s-2006-961438>
30. Negi PS, Jayaprakasha GK, Jagan Mohan Rao L, Sakariah KK. Antibacterial activity of turmeric oil: a byproduct from curcumin manufacture. *J Agric Food Chem*. 1999;47:4297-300. <https://doi.org/10.1021/jf990308d>
 31. Miquel J, Bernd A, Sempere JM, Díaz-Alperi J, Ramírez A. The curcuma antioxidants: pharmacological effects and prospects for future clinical use. a review. *Arch Gerontol Geriatr*. 2002;34:37-46. [https://doi.org/10.1016/S0167-4943\(01\)00194-7](https://doi.org/10.1016/S0167-4943(01)00194-7)
 32. Maneewan C, Yamauchi KE, Mekbungwan A, Maneewan B, Siri S. Effect of turmeric (*Curcuma longa* Linnaeus) on growth performance, nutrient digestibility, hematological values, and intestinal histology in nursery pigs. *J Swine Health Prod*. 2012;20:231-40.
 33. Alagbe JO. Growth performance and blood parameters of weaner pigs fed diets supplemented with turmeric powder. *Sch J Agric Sci*. 2017;7:57-61.
 34. National Research Council. Nutrient requirements of swine. Washington, DC: National Academies Press; 2012.
 35. Cho S, Hwang O, Park S. Effect of dietary protein levels on composition of odorous compounds and bacterial ecology in pig manure. *Asian-Australas J Anim Sci*. 2015;28:1362-70. <https://doi.org/10.5713/ajas.15.0078>
 36. Balasubramanian B, Lee SI, Kim IH. Inclusion of dietary multi-species probiotic on growth performance, nutrient digestibility, meat quality traits, faecal microbiota and diarrhoea score in growing-finishing pigs. *Ital J Anim Sci*. 2018;17:100-6. <https://doi.org/10.1080/1828051X.2017.1340097>
 37. Kang SJ, Jang CS, Son JM, Hong KW. Comparison of seven commercial TaqMan master mixes and two real-time PCR platforms regarding the rapid detection of porcine DNA. *Food Sci Anim Res*. 2021;41:85-94. <https://doi.org/10.5851/kosfa.2020.e80>
 38. Platel K, Srinivasan K. Influence of dietary spices and their active principles on pancreatic digestive enzymes in albino rats. *Food/Nahrung*. 2000;44:42-6. [https://doi.org/10.1002/\(SICI\)1521-3803\(20000101\)44:1<42::AID-FOOD42>3.0.CO;2-D](https://doi.org/10.1002/(SICI)1521-3803(20000101)44:1<42::AID-FOOD42>3.0.CO;2-D)
 39. Singh PK, Kumar A, Tiwari DP, Kumar A, Palod J. Effect of graded levels of dietary turmeric (*Curcuma longa*) powder on performance of broiler chicken. *Ind J Anim Nutr*. 2018;35:428-35. <http://doi.org/10.5958/2231-6744.2018.00065.8>
 40. Tubcharoen S, Chomchai S, Saardrak K, Pisarnpong C, Dirokwirat W, Ponglumjeak S. Effects of dietary turmeric meal supplementation on production performance and carcass quality of growing-finishing pigs. In: Proceedings of 45 Kasetsart University Annual Conference; 2007; Bangkok.
 41. Etim NN, Offiong EEA, Williams ME, Asuquo LE. Influence of nutrition on blood parameters of pigs. *Am J Biol Life Sci*. 2014;2:46-52.
 42. Shao W, Yu Z, Chiang Y, Yang Y, Chai T, Foltz W, et al. Curcumin prevents high fat diet induced insulin resistance and obesity via attenuating lipogenesis in liver and inflammatory pathway in adipocytes. *PLOS ONE*. 2012;7:e28784. <https://doi.org/10.1371/journal.pone.0028784>
 43. Porn-anek P, Uriyapongson S, Vichasilp C. Enhanced curcumin solubility by solid dispersion technique reduces fat accumulation in pigs. *Food Appl Biosci J*. 2017;5:104-14. <https://doi.org/10.14456/fabj.2017.9>
 44. Henningsson Å, Björck I, Nyman M. Short-chain fatty acid formation at fermentation of indigestible carbohydrates. *Näringsforskning*. 2001;45:165-8. <https://doi.org/10.3402/fnr.v45i0.1801>
 45. Williams BA, Grant LJ, Gidley MJ, Mikkelsen D. Gut fermentation of dietary fibres: physico-chemistry of plant cell walls and implications for health. *Int J Mol Sci*. 2017;18:2203.

- <https://doi.org/10.3390/ijms18102203>
46. Park S, Cho E, Chung H, Cho K, Sa S, Balasubramanian B, et al. Digestibility of phosphorous in cereals and co-products for animal feed. *Saudi J Biol Sci.* 2019;26:373-7. <https://doi.org/10.1016/j.sjbs.2018.12.003>
 47. Zhao J, Bai Y, Tao S, Zhang G, Wang J, Liu L, et al. Fiber-rich foods affected gut bacterial community and short-chain fatty acids production in pig model. *J Funct Foods.* 2019;57:266-74. <https://doi.org/10.1016/j.jff.2019.04.009>
 48. Han KH, Azuma S, Fukushima M. In vitro fermentation of spent turmeric powder with a mixed culture of pig faecal bacteria. *Food Funct.* 2014;5:2446-52. <https://doi.org/10.1039/C4FO00142G>
 49. Xun W, Shi L, Zhou H, Hou G, Cao T, Zhao C. Effects of curcumin on growth performance, jejunal mucosal membrane integrity, morphology and immune status in weaned piglets challenged with enterotoxigenic *Escherichia coli*. *Int Immunopharmacol.* 2015;27:46-52. <https://doi.org/10.1016/j.intimp.2015.04.038>
 50. Pluske JR, Thompson MJ, Atwood CS, Bird PH, Williams IH, Hartmann PE. Maintenance of villus height and crypt depth, and enhancement of disaccharide digestion and monosaccharide absorption, in piglets fed on cows' whole milk after weaning. *Br J Nutr.* 1996;76:409-22. <https://doi.org/10.1079/BJN19960046>
 51. Avançaço GB, Ferreira FD, Bomfim NS, de Souza Rodrigues dos Santos PA, Peralta RM, Brugnari T, et al. *Curcuma longa* L. essential oil composition, antioxidant effect, and effect on *Fusarium verticillioides* and fumonisin production. *Food Control.* 2017;73:806-13. <https://doi.org/10.1016/j.foodcont.2016.09.032>
 52. Chen L, Xu Y, Chen X, Fang C, Zhao L, Chen F. The maturing development of gut microbiota in commercial piglets during the weaning transition. *Front Microbiol.* 2017;8:1688. <https://doi.org/10.3389/fmicb.2017.01688>
 53. Hu J, Nie Y, Chen J, Zhang Y, Wang Z, Fan Q, et al. Gradual changes of gut microbiota in weaned miniature piglets. *Front Microbiol.* 2016;7:1727. <https://doi.org/10.3389/fmicb.2016.01727>
 54. Zhai SS, Ruan D, Zhu YW, Li MC, Ye H, Wang WC, et al. Protective effect of curcumin on ochratoxin a-induced liver oxidative injury in duck is mediated by modulating lipid metabolism and the intestinal microbiota. *Poult Sci.* 2020;99:1124-34. <https://doi.org/10.1016/j.psj.2019.10.041>
 55. Lukiw WJ. *Bacteroides fragilis* lipopolysaccharide and inflammatory signaling in Alzheimer's disease. *Front Microbiol.* 2016;7:1544. <https://doi.org/10.3389/fmicb.2016.01544>
 56. Zhao Y, Wu J, Li JV, Zhou NY, Tang H, Wang Y. Gut microbiota composition modifies fecal metabolic profiles in mice. *J Proteome Res.* 2013;12:2987-99. <https://doi.org/10.1021/pr400263n>
 57. Shen L, Liu L, Ji HF. Regulative effects of curcumin spice administration on gut microbiota and its pharmacological implications. *Food Nutr Res.* 2017;61:1361780. <https://doi.org/10.1080/16546628.2017.1361780>
 58. Alauzet C, Marchandin H, Lozniewski A. New insights into *Prevotella* diversity and medical microbiology. *Future Microbiol.* 2010;5:1695-718. <https://doi.org/10.2217/fmb.10.126>
 59. Amat S, Lantz H, Munyaka PM, Willing BP. *Prevotella* in pigs: the positive and negative associations with production and health. *Microorganisms.* 2020;8:1584. <https://doi.org/10.3390/microorganisms8101584>
 60. Kosti D, Dahiya DS, Dalal R, Vinus. Effect of supplementing different levels of turmeric powder on ileal morphology in laying hens. *Int J Curr Microbiol Appl Sci.* 2018;7:1524-32. <https://doi.org/10.3390/ijms18102203>

doi.org/10.20546/ijcmas.2018.707.179

61. Wang J, Ghosh SS, Ghosh S. Curcumin improves intestinal barrier function: modulation of intracellular signaling, and organization of tight junctions. *Am J Physiol Cell Physiol.* 2017;312:C438-45. <https://doi.org/10.1152/ajpcell.00235.2016>