

The current state of phage therapy in livestock and companion animals

Youbin Choi[#], Woongji Lee[#], Joon-Gi Kwon[#], Anna Kang[#], Min-Jin Kwak, Ju-Young Eor and Younghoon Kim*

Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea



Received: Dec 29, 2023
 Revised: Jan 6, 2024
 Accepted: Jan 6, 2024

[#]These authors contributed equally to this work.

*Corresponding author

Younghoon Kim
 Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea.
 Tel: +82-2-880-4808
 E-mail: ykeys2584@snu.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Youbin Choi
<https://orcid.org/0000-0002-9444-3237>
 Woongji Lee
<https://orcid.org/0000-0001-7243-0702>
 Joon-Gi Kwon
<https://orcid.org/0000-0001-9953-6945>
 Anna Kang
<https://orcid.org/0000-0003-0208-6234>
 Min-Jin Kwak
<https://orcid.org/0000-0001-9832-3251>
 Ju-Young Eor
<https://orcid.org/0000-0002-3764-3339>
 Younghoon Kim
<https://orcid.org/0000-0001-6769-0657>

Competing interests

No potential conflict of interest relevant to this article was reported.

Abstract

In a global context, bacterial diseases caused by pathogenic bacteria have inflicted sustained damage on both humans and animals. Although antibiotics initially appeared to offer an easy treatment for most bacterial infections, the recent rise of multidrug-resistant bacteria, stemming from antibiotic misuse, has prompted regulatory measures to control antibiotic usage. Consequently, various alternatives to antibiotics are being explored, with a particular focus on bacteriophage (phage) therapy for treating bacterial diseases in animals. Animals are broadly categorized into livestock, closely associated with human dietary habits, and companion animals, which have attracted increasing attention. This study highlights phage therapy cases targeting prominent bacterial strains in various animals. In recent years, research on bacteriophages has gained considerable attention, suggesting a promising avenue for developing alternative substances to antibiotics, particularly crucial for addressing challenging bacterial diseases in the future.

Keywords: Bacteriophage, Antibiotics alternatives, Livestock, Companion animals

INTRODUCTION

The ongoing threat of pathogenic bacteria has persisted throughout human history, and recent advancements in public health awareness and hygiene practices were expected to alleviate infections caused by these microbes [1]. However, the invasion of bacteria from the external environment, beyond human control, has remained challenging in terms of prevention [2–4]. In particular, damages caused by livestock (swine, chicken, and bovine) closely associated with human dietary habits and escalating issues involving companion animals (canines and felines) worldwide present an ongoing challenge that requires resolution [5–8].

The most effective and straightforward method for treating bacterial infections is the use of antibiotics specifically tailored to the bacteria in question [9,10]. However, since the initial discovery of antibiotics, their potent effects have led to global misuse, resulting in various side effects and contributing to substantial environmental pollution and public health concerns. Of these issues, the emergence of antibiotic-resistant bacteria poses the most serious threat [11,12]. In particular, multidrug-resistant bacteria, resistant to two or more antibiotics, require high antibiotic concentrations or additional administration of different antibiotics for treatment, exacerbating the situation [13]. Therefore, while proper antibiotic use to minimize the emergence of resistant strains and active

Funding sources

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the National Research Foundation of Korea Grant, funded by the Korean government (MEST) (NRF-2021R1A2C3011051) and by the Korea government (MSIT) (No. RS-2023-00218476).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Choi Y, Lee W, Kwon JG, Kang A, Kwak MJ, Eor JY, Kim Y.
 Data curation: Choi Y, Lee W, Kwon JG, Kang A.
 Formal analysis: Choi Y, Lee W, Kwon JG, Kang A.
 Writing - original draft: Choi Y, Lee W, Kwon JG, Kang A, Kwak MJ, Eor JY, Kim Y.
 Writing - review & editing: Choi Y, Lee W, Kwon JG, Kang A, Kwak MJ, Eor JY, Kim Y.

Ethics approval and consent to participate

This article does not require IRB/ACUC approval because there are no human and animal participants.

infection management is crucial, the urgent development of antibiotic alternatives is paramount.

Among various alternatives to antibiotic substances, such as probiotics, organic acids, bacteriocins, and antimicrobial peptides, we have focused on the potential of bacteriophages as antimicrobial agents (Fig. 1) [9,14–17]. In contrast to indiscriminate bombardment by antibiotics, bacteriophages (phages) possess characteristics akin to guided missiles, selectively infecting and killing specific bacteria (Fig. 2). Some phages exhibit distinct specificity, infecting only a single species of bacteria, whereas others can infect multiple species. In either case, the bacteria serving as hosts are closely related taxonomically [18]. Virulent phages carry only a lytic life cycle, invading the host’s interior, replicating, assembling, and ultimately destroying host cells through release. It is estimated that there are approximately 10³¹ phages on earth, surpassing the number of bacteria. The specificity, prolificacy, and abundance of phages distinguish them from other antibiotic alternatives, garnering renewed attention as substitutes for antibiotics (Table 1) [19].

In an environment where pathogenic bacteria are prevalent, such as the livestock environment, the need for antibiotics is deemed critical [20]. However, there is a global trend toward the prohibition of antibiotic use in the livestock industry. Despite this trend, the prevalence of antibiotic-resistant strains continues to rise, attributed to the persistent use of antibiotics for purposes such as the treatment of bacterial diseases [21]. To address these challenges, experimental studies using phages for the inhibition of various pathogenic bacteria are underway (Table 2) [22]. On a national scale, legal permissions have been granted for the use of antibiotic alternatives within the livestock industry as veterinary drugs to mitigate these issues.

The pet industry is continuously growing globally, and recently, the term “Petconomy” has emerged. According to the American Pet Products Association (APPA), the expenditure in the U.S. pet industry surpassed \$95.7 billion in 2019 [23]. According to Euromonitor, the global pet food market size will reach \$102.1 billion in 2020 [24]. With the increasing scale of the Petconomy,




Fig. 1. Interaction among bacteriophages, pathogenic bacteria, and the host animals. Pathogens infect the host, causing various diseases, whereas bacteriophages, either as single phages or phage cocktails, are used to control the pathogens. Consequently, phage therapy can be used as a treatment to address bacterial diseases caused by pathogenic bacteria. *S. aureus*, *Staphylococcus aureus*; *E. coli*, *Escherichia coli*; *S. Enteritidis*, *Salmonella Enteritidis*; *S. Typhimurium*, *Salmonella Typhimurium*; *C. jejuni*, *Campylobacter jejuni*; *S. Gallinarum*, *Salmonella Gallinarum*; *P. aeruginosa*, *Pseudomonas aeruginosa*; *S. pseudintermedius*, *Staphylococcus pseudintermedius*; *L. infantum*, *Leishmania infantum*.




Fig. 2. Bacteriophages have advantages as alternatives to antibiotics, particularly with high specificity for host cells and self-replication capabilities. This allows effective targeting, elimination, and proliferation of pathogenic bacteria even at low concentrations. Additionally, phages can prevent the development of antibiotic-resistant strains and dysbiosis in microbial communities associated with antibiotic use.

Table 1. Difference between phage and antibiotics

	Bacteriophage	Antibiotics
Specificity	High specificity	Little specificity
Dosage	Auto-dosage	Fixed dosage
Side effects	Little side effects	Multiple side effects
Replication	Self-replication	No replication
Resistance	Phage-resistance bacteria remains susceptible to other phages	Resistance is not limited to target bacteria
Application	Can penetrate biofilms	Cannot penetrate biofilms
Host range	Narrow host range	Wide host range
Cost	Cost-efficient	Time and cost consuming

there has been a growing interest in the health and well-being of animals, leading to a rise in the use of antibiotics for treating animal diseases [25,26]. However, the use of antibiotics has raised health issues, particularly issues related to the reduction of probiotics in the gut microbiota, which is becoming a significant consideration for pet owners when addressing animal health [27]. As a response to preventing intestinal pathogen infections, various products using phages have been introduced, and research on phage-based treatments for pets has increased (Table 3) [28,29]. This study aims to introduce successful cases of treating bacterial diseases in animals using phages, emphasizing the potential utility of phages in treating pathogenic bacterial infections, considering the close connection between human life and animals, especially livestock and pets, where

Table 2. Phage therapy in the livestock industry over the last two decades

Animals	Target bacteria	Symptoms	Phage application	Therapeutic activity	References
Bovine	<i>S. aureus</i> (MDR)	Mastitis	SAH-1	Significantly reduced bacterial growth at MOIs of 1–100	[111]
	<i>S. aureus</i> (MRSA)		Six phage cocktails	Reduced <i>S. aureus</i> CFU counts by 64%–95%	[112]
	<i>S. aureus</i> (MSSA, MRSA VISA)		Five phage cocktails	Reduced colonization in the mouse mammary gland 8 h after treatment and prophylactically 4 h before challenge was most effective	[113]
	<i>S. aureus</i> (MRSA)		PhiSA012, PhiSA039	Intravenous and intraperitoneal administration of SA012 reduced bacterial colonization and inflammation of the mammary gland.	[114–116]
	<i>S. aureus</i> (MRSA)		vB_SauM_SDQ	Lysed 20 of the 24 strains reduced established biofilms on polystyrene, milk, and mammary gland tissue after treatment.	[117]
	<i>S. aureus</i> (MRSA)		Phage 24 A2	Lysed 19 of the 30 strains examined. Phage-cleared bacterial cultures on agar at a MOI of 10, supporting topical application for therapeutic use	[118]
	<i>S. aureus</i> (MRSA)		SLPW	Lysed 36 of the 40 isolates examined. Phage administration remedially reduced colonization and inflammation of cytokines in mice.	[119]
	<i>E. coli</i> (MDR)	Diarrhea	Three-phage cocktails	A cocktail of the phages reduced colonization, somatic cells, and inflammatory factors and alleviated symptoms of mastitis in cattle.	[42]
	<i>E. coli</i> (MPEC)		vB_EcoM_UFV13	A 10-fold reduction in bacterial load was observed at a MOI of 10 in mice.	[120]
	<i>E. coli</i> (ETEC, EPEC)		Three-phage cocktails	The probiotic-phage suppositories reduced the duration of diarrhea in calves, completely eliminating it within 24–48 h after use.	[42]
<i>E. coli</i> (ETEC)	VTCCBPA9		VTCCBPA9 showed bactericidal activity against 47.3% (62/131) <i>E. coli</i> isolates, including three ETEC strains.	[121]	
Swine	<i>S. Enteritidis</i>	Salmonellosis	Phage cocktail	3.5 log CFU reduction of <i>S. Enteritidis</i> PT4 per gram of cecal content	[122]
	<i>S. Enteritidis</i>		Felix 01	100% efficacy in eliminating <i>S. Enteritidis</i> strains from tonsils 6 h after application of bacteriophage suspension	[123]
	<i>S. Typhimurium</i>		Phage with probiotics	Significant influence on the growth of weaned pigs in comparison with pigs not treated with phages	[124]
	<i>S. Typhimurium</i>		Single phage	Significant reduction (99%) or complete elimination (100%) of <i>S. Typhimurium</i> strains in ileum, tonsils, and cecum samples within 48 h after the first administration	[48]
	<i>S. Typhimurium</i>		Phage cocktail	Significant reduction in the concentration of 2 of 3 serovars (<i>S. Enteritidis</i> and <i>Typhimurium</i>) by 2–4 log CFU after administration of bacteriophage suspension at 1011 PFU	[125]
	<i>S. Typhimurium</i>		Phage cocktail	100% reduction of the <i>Salmonella</i> ATCC 14028 reference strain and 92.5% of field isolates	[126]
	<i>E. coli</i> (ETEC)		Diarrhea	Phage cocktail	In comparison with the control group, the <i>E. coli</i> K88 CFUs in the duodenum, jejunum, ileum, cecum, colon, and mesenteric lymph nodes were lower in each phage-treated group, with differences at log levels of 0.83, 1.61, 1.67, 2.4, 1.47, and 1.65, respectively.
	<i>E. coli</i>	A221		With the treatment of phage A221, the body weight of piglets increased, and the percentage of Enterobacteriaceae in duodenum decreased to 0.64%.	[127]
	<i>E. coli</i> (MDR)	PT-10		When comparing the <i>E. coli</i> CFU in the feces between the phage-treated group and the control group, a difference of 4 log levels was observed on the 10th day.	[103]
	<i>E. coli</i> (ETEC)	Single phage		When comparing the number of <i>E. coli</i> isolated from challenged pig feces to the control group, there was an average difference of 1 log level throughout the experimental period.	[128]

Table 2. Continued

Animals	Target bacteria	Symptoms	Phage application	Therapeutic activity	References
Chicken	<i>C. jejuni</i>	Campylobacteriosis	Cj6	The maximum (2 log ₁₀ levels) reduction was achieved in samples that were treated with high densities of <i>C. jejuni</i> and high MOI of the phage at both storage temperatures.	[66]
	<i>C. jejuni</i>		Three-phage cocktails	2 log ₁₀ levels of reduction were achieved using the cocktail consisting of the three phages.	[129]
	<i>C. jejuni</i>		CP81	No reduction was observed at 4°C <i>in situ</i> on meat or <i>in vitro</i> . 1 log ₁₀ reduction was observed <i>in vitro</i> at 37°C.	[67]
	<i>C. jejuni</i>		Phage cocktail	3.2 log ₁₀ CFU/g lower <i>C. jejuni</i> counts than in the control until slaughter	[71]
	<i>C. jejuni</i>		Φ3, Φ15	Modest reduction of 0.2 log ₁₀ level. 0.8 log ₁₀ at 4°C	[130]
	<i>S. Gallinarum</i>	Salmonellosis	CJø01	Treatment using bacteriophages as a feed additive for chickens having contact with infected individuals led to a mortality rate of only 5%, as compared with 30% in the group that did not receive phage therapy.	[75]
	<i>S. Enteritidis</i>		PSE	100% efficacy in eliminating <i>S. Enteritidis</i> strains from the tonsils; 6 h after application of bacteriophage suspension	[131]
	<i>S. Enteritidis</i>		CJ07	The highest doses of bacteriophage significantly inhibited the replication of pathogens in the digestive tract of the chickens.	[132]
	<i>S. Typhimurium</i>		Three-phage cocktails	10-fold reduction in bacteria in the chicken ileum, cecum, liver, and spleen	[76]

In the livestock industry, various studies have been conducted to address the prevalent issue of pathogenic bacteria and bacterial diseases. This summary compiles research efforts aimed at preventing bacterial diseases in livestock through the application of bacteriophages. Numerous studies have been conducted, and further advancements in research are anticipated. The "Phage application" section provides information on the names or numbers of phages used in the treatment.

S. aureus, *Staphylococcus aureus*; MDR, multi-drug resistance; MSSA, methicillin-sensitive *Staphylococcus aureus*; MRSA, methicillin-resistant *Staphylococcus aureus*; VISA, vancomycin Intermediate *Staphylococcus aureus* (VISA); *E. coli*, *Escherichia coli*; MPEC, mammary pathogenic *E. coli*; ETEC, enterotoxigenic *Escherichia coli*; EPEC, enteropathogenic *E. coli*; *C. jejuni*, *Campylobacter jejuni*.

antibiotics are the most commonly used.

PHAGE THERAPY IN THE LIVESTOCK INDUSTRY

Bovine

Staphylococcus aureus

Bovine mastitis, characterized by inflammation of the mammary gland due to bacterial infection via the teat canal, stands out as one of the most crucial diseases [30]. The condition is classified into clinical and subclinical mastitis based on symptoms. Clinical mastitis presents visible abnormalities such as redness, udder swelling, and milk clot formation. In contrast, subclinical mastitis lacks visible abnormalities or milk clotting but can be confirmed by an increase in somatic cells [31]. Mastitis is further categorized as contagious or environmental depending on the pathogen involved. Contagious mastitis, caused by pathogens like *Staphylococcus aureus*, *Streptococcus agalactiae*, and *Corynebacterium bovis*, can transmit from one cattle to another. Environmental mastitis is instigated by pathogens such as *Escherichia coli*, *Klebsiella* spp., *Streptococcus dysgalactiae*, and *Streptococcus uberis* [8,32].

Among these pathogens, *S. aureus*, along with nonaureus staphylococci, is the most frequently detected contagious pathogen [33]. *E. coli*, a common Gram-negative bacterium, is also recognized as a frequent mastitis-causing pathogen [34]. Notably, while *E. coli* rapidly induces an immune response, *S. aureus* does not trigger a similar response. Consequently, *S. aureus* infections tend to be milder, leading to chronic mastitis lasting for months. In the previous study, the efficacy of a phage cocktail composed of two phages in inhibiting *S. aureus* N305, isolated from bovine mastitis,

Table 3. Phage therapy in companion animal industry

Animals	Target bacteria	Symptoms	Phage application	Therapeutic activity	References
Canine	<i>P. aeruginosa</i>	Otitis media	Pa12, Pa18	This study explored two sewage-derived phages capable of lysing a variety of <i>P. aeruginosa</i> strains, even those resistant to fluoroquinolones. Notably, the phages showed effectiveness against isolates with high enrofloxacin and orbifloxacin resistance.	[78]
	<i>P. aeruginosa</i>		Phage cocktail	A bacteriophage treatment for chronic <i>P. aeruginosa</i> otitis in 10 dogs resulted in a significant improvement within 48 h, with a mean clinical score fall of 30.1% and a mean <i>P. aeruginosa</i> count fall of 67%.	[77]
	<i>E. coli</i> (UPEC)	Urinary tract infections	Single phage	Bacteriophages effectively lysed 94% of UPEC strains, with 10 of them individually lysing $\geq 51\%$ of the strains.	[79]
	<i>S. pseudintermedius</i> (MRSP)	Skin infections	Single phage	All phages (n = 4) successfully lysed all MRSP isolates (n = 17); however, their lytic activity was restricted to <i>S. pseudintermedius</i> and <i>S. schleiferi</i> .	[80]
	<i>Staphylococcus</i> spp. (MDR)	Conjunctivitis	Single phage	The phage eye drops tested in the experiment demonstrated 100% effectiveness <i>in vitro</i> against the <i>Staphylococcus</i> isolates under investigation.	[133]
	<i>L. infantum</i>	Visceral leishmaniasis	CVL serodiagnosis	This study found eight mimotopes for accurate CVL serodiagnosis using a cost-effective and straightforward phage-ELISA assay.	[83]
Feline	<i>E. coli</i> (UPEC)	Urinary tract infections	Single phage	Bacteriophages effectively lysed 94% of UPEC strains, with 10 of them individually lysing $\geq 51\%$ of the strains.	[79]

In the companion animal industry, various studies have been conducted to address the prevalent issue of pathogenic bacteria and bacterial diseases, using bacteriophages for prevention. The use of phages in the companion animal industry is relatively limited compared with livestock, but it is predicted that more research will be conducted in the future. The "Phage application" section provides information on the names or numbers of phages used in the treatment.

P. aeruginosa, *Pseudomonas aeruginosa*; *E. coli*, *Escherichia coli*; UPEC, UroPathogenic *Escherichia coli*; *L. infantum*, *Leishmania infantum*.

was verified through *in vitro* testing. Subsequently, an *in vivo* test was conducted, confirming the efficacy of the phage cocktail in a mouse mastitis model created by inoculating the *S. aureus* N305 strain into a lactating mouse [35]. Another research team isolated five *S. aureus* lytic phages from mastitis cow milk samples, belonging to the *Podoviridae* family. After *in vitro* characterization of the five phages, the efficacy of a phage cocktail was confirmed in a mouse mastitis model inoculated with *S. aureus* CVCC 546. Phage cocktail therapy alleviated the immune response, as evidenced by the reduction in proinflammatory cytokines tumor necrosis factor alpha (TNF- α) and interleukin 6 (IL-6), along with histopathological analysis [36]. Given that previous experiments utilized a mouse model, the actual application of bacteriophages to bovine mastitis remains uncertain. Prior study conducted an experiment with 24 lactating Holstein cows with preexisting asymptomatic *S. aureus* mastitis. Using a single bacteriophage (K - ATCC 19685-B1) for all experiments, intramammary infusions of either 1.25×10^{11} plaque forming unit (PFU) or saline were administered once per day for 5 days. Following bacteriophage infusion, mean milk production increased [37]. In prior study, the recombinant endolysin (Trx-SA1) from a novel bacteriophage (IME-SA1) was obtained for the experiment. The udder-infected group with *S. aureus* showed a decrease in both somatic cell count and pathogen level when treated with Trx-SA1, whereas the udder-infected group with *E. coli* did not exhibit the same response [38].

Escherichia coli

Neonatal calf diarrhea, defined as diarrhea occurring within the first month of life, is a common occurrence [39]. This condition can be caused by various factors, including viruses, bacteria, and stress. Antibiotics have historically been extensively used to treat bacterial-induced diarrhea in calves. However, the emergence of antibiotic-resistant bacteria and the presence of antibiotic residues have raised concerns [40]. In response to these issues, a study utilizing bacteriophages is

currently underway.

Screening a cocktail of four unique bacteriophages based on 36 *E. coli* strains isolated from cattle with clinical mastitis, researchers observed that the bacteriophage cocktail significantly inhibited the growth of 58% of strains from Washington State and 54% of *E. coli* mastitis isolates from New York State. This suggests a relatively broad spectrum of action against related strains in two distinct regions [41]. In another study, three lytic phages—SYGD1, SYGE1, and SYGMH1—were isolated from the sewage of a dairy farm. These phages were administered in cocktail form to cattle with mastitis. Similar to antibiotic treatment, reductions in *E. coli* CFU and milk somatic cell count were noted, accompanied by a decrease in proinflammatory cytokines, IL- β and TNF- α [42]. In a related study, a combination of probiotics and bacteriophages was explored as an alternative to antibiotics for controlling *E. coli*, a common cause of neonatal calf diarrhea. The study comprised four groups: a healthy control group, a positive control group with diarrhea, a group treated with probiotics and bacteriophages in healthy calves, and a group treated with probiotics and bacteriophages in calves with diarrhea. Over an observation period of 11 days, the use of probiotic-phage suppositories led to a reduction in the duration of diarrhea in calves, with symptoms completely disappearing within 24–48 hours post-treatment. This therapeutic approach stimulated the activation of the calf's immune system, enhancing both specific and nonspecific responses and increasing resistance to infection [43].

Swine

***Salmonella* spp.**

The domestic pig, *Sus scrofa domesticus*, holds significant economic and nutritional importance, being raised in various industrial systems and habitats. Susceptibility to emerging and re-emerging infections is common among swine [44]. The introduction of pigs into diverse environments by humans can lead to viral adaptability and genetic recombination within infectious agents. Specific circumstances, such as changes in land use, trade, and other human-induced occurrences, can promote the transmission and dissemination of infections, potentially resulting in the development of new viruses with zoonotic potential [45]. Swine health is a critical consideration for producers from farrowing through slaughter, impacting profitability through reduced weight gain or animal mortality in the event of herd infections [46]. Enteric viruses, especially highly contagious ones like the porcine epidemic diarrhea virus, pose significant economic challenges for farmers and the entire industry [47].

Examining phage dispersion tactics, researchers explored oral gavage and the addition of phages to drinking water and feed. Investigating the administration of phages through oral gavage, challenged pigs received phage cocktails every 2 hours for 6 hours. While this approach reduced *Salmonella* levels in cecal samples compared to a control, it is considered technically challenging for practical use on multiple infected pigs [48]. Instead, a delivery method enabling simultaneous treatment of several animals, such as in feed or water, becomes essential. One research was exemplified by the assessment of a three-phage cocktail in drinking water, revealing a noteworthy reduction in intestinal inflammation and *Salmonella* invasion. [49]. Although water-based phage dissemination is simple, it reduces product shelf life, and transporting large volumes of liquid phages is challenging [50]. Moreover, although high-titer phage stocks may be introduced to drinking water as a concentrate or powder, there is still the chance of human error [51]. Feed-based distribution offers various benefits, and recent advances in phage formulation studies demonstrated that phages can be transformed into a fine powder using excipients through lyophilizing, freeze-drying, or spray drying [52]. Excipients keep phages from drying out and act as agents that replace water to keep virion particles stable while they dry [53]. FDA-approved phage drying excipients,

including trehalose, sucrose, mannitol, leucine, and the pH-sensitive polymer Eudragit S100, maintain phage stability during drying [54,55]. To date, no published study has documented improving phage drying at a commercial scale, but companies have filed patents based on drying phages, indicating a growing interest in developing dried phage products.

Administering a microencapsulated phage cocktail to young pigs challenged with *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) 2, 4, and 6 h later, researchers observed significantly lower ileal concentrations of *S. Typhimurium* in phage-treated pigs compared to untreated pigs. Moreover, the ileal *S. Typhimurium* concentrations were more frequently below the detectable limit in phage-treated pig samples than in untreated pig samples [56]. In two trials, researchers administered a microencapsulated phage cocktail to young pigs infected with *S. Typhimurium*. Concentrations of *S. Typhimurium* in ileal, cecal, and cecal tonsil samples were numerically lower (not statistically compared) in phage-treated pigs at necropsy (6 h post-challenge) than in untreated pigs. While some lymph node samples from both untreated and phage-treated pigs were positive for the challenge organism 6 h after exposure, all fecal samples were positive for the challenge organism. *S. Typhimurium* levels in phage-treated pigs were below the detection limit in several samples [48]. Using a phage cocktail suspended in sodium bicarbonate, researchers treated *S. Typhimurium*-challenged market-weight pigs. At necropsy, the quantities of *S. Typhimurium* in cecal and ileal samples were not significantly different ($p > 0.05$) between phage-treated and untreated pigs (18 ± 2 h post-treatment). Numerical variations in *S. Typhimurium* concentrations were most apparent between groups of pigs treated with 10^{10} PFU/pig and untreated pigs. Similarly, despite numerical changes in incidence, the incidence of *S. Typhimurium* in necropsy fecal samples was not significantly different ($p > 0.05$) in phage-treated versus untreated pigs [57].

Escherichia coli

In a study, 28-day-old pigs were infected with *E. coli* K88 and K99 using a phage cocktail. The phage treatment began on the day of the challenge and continued until the experiment concluded (7 days post-challenge). Fecal *E. coli* K99 concentrations did not differ significantly ($p > 0.05$) between phage-treated and untreated pigs during the study (1, 3, and 7 days post-challenge). However, overall fecal *E. coli* K88 concentrations in phage-treated pigs were significantly lower ($p < 0.01$) than those in untreated pigs. Adhesion of *E. coli* K88 to the ileum and cecum of phage-treated pigs was significantly lower ($p < 0.05$) than that of untreated pigs. The adherence of *E. coli* K88 to the duodenum, jejunum, colon, and mesenteric lymph nodes did not differ significantly ($p > 0.05$) between treatment groups. *E. coli* K99 adhesion did not differ significantly ($p > 0.05$) between phage-treated and untreated pigs in any sample type [58].

Regarding another investigation, initial antibiotic resistance was found in 87.3% of *E. coli* strains isolated from healthy or diarrheal fecal samples from three pig farms. Using these 87 *E. coli* strains as indicator hosts, 45 coliphages were extracted, showing a higher abundance in the postweaning stage than in the preweaning stage (24 versus 17 in the Nanjing farm and 13 versus 4 in the Chuzhou farm). Furthermore, each farm included the most common coliphage strain. Pathogenic *E. coli*-specific bacteriophages were detected in the intestines of examined pigs (9/10 in Nanjing farm and 7/10 in Chuzhou farm), and the majority had significant bacteriostatic effects ($p < 0.05$) on pathogenic *E. coli* strains. Polyvalent bacteriophages N24, N30, and C5 were found. With relatively few differences in infection characteristics, the N30 and C5 strains shared 89.67% genetic similarity. Pathogenic *E. coli*-specific bacteriophages and polyvalent bacteriophages are common in piglet stomachs, and weaning is an important event that determines coliphage abundance [59].

Chicken

Campylobacter jejuni

Campylobacter jejuni is a prevalent pathogenic bacterium globally and a leading cause of foodborne illness. Typically associated with poultry, *C. jejuni* naturally inhabits the digestive tracts of various bird species and is most commonly found in animal feces [60]. Fortunately, infections caused by *C. jejuni* can lead to severe gastroenteritis but are rarely life-threatening [61]. Nonetheless, in the United States alone, approximately 2 million cases of *Campylobacter* enteritis occur annually, constituting 5%–7% of gastroenteritis cases, with the majority attributed to poultry, including chickens, turkeys, and waterfowl [62]. In fact, the cornerstone of *Campylobacter* enteritis treatment lies in maintaining electrolyte balance rather than antibiotic therapy, as most patients with this infection experience self-limiting illnesses that do not necessitate antibiotics. However, antibiotics are warranted in specific clinical scenarios such as high fever, bloody stools, pregnancy, and human immunodeficiency virus infection [63]. To prevent such situations, minimizing bacterial infection during the early stages of poultry growth is crucial. Consequently, several studies exploring the inhibition of *C. jejuni* using bacteriophage, an antibiotic alternative, are underway. Still, there is a scarcity of patents related to phage products targeting this bacterium. In conclusion, as global chicken consumption continues to rise, ongoing research on *Campylobacter* phages is essential.

Until now, various *in situ* and *in vivo* studies related to *C. jejuni*-infecting phages have been conducted. Atterbury *et al.* preserved chicken skin sections inoculated with varying concentrations of *C. jejuni* and bacteriophages at 4°C and 20°C. At the maximum phage concentration (10^7), there was a reduction of bacteria by 1.1–1.3 log levels in the 4°C treatment setting and a 2.3–2.5-log reduction in the 20°C treatment setting compared with the control group [64]. Previous study observed a 1-log reduction in chicken skin treated with phages at a concentration of 10^6 PFU/cm² compared with untreated control groups [65]. Another research team treated beef samples inoculated with *C. jejuni* with bacteriophages and stored them at 5°C and 24°C. High-density *C. jejuni* and phage-treated samples achieved a maximum reduction (2 log levels) at both storage temperatures [66]. The other study observed no reduction *in situ* or *in vitro* in meat at 4°C, whereas a 1-log reduction was observed *in vitro* within the test tube at 37°C [67].

In addition to the aforementioned *in situ* experiments, various *in vivo* experiments have been conducted. Previous study observed an initial 3-log reduction, followed by a resurgence in *C. jejuni* CFU counts within 5 days, which were 1 log lower than the control group [68]. Similarly, another research demonstrated reductions ranging from 0.5 to 5 log levels depending on the gut region and phage-host combinations, with the largest bacterial reduction achieved within 24–48 h [69]. Additionally, prior study achieved a 2-log reduction using a cocktail of three phages. Interestingly, phage delivery through food was more effective than oral gavage in nutrient broth [70]. The other research group administered a phage cocktail to birds through drinking water, resulting in *C. jejuni* counts 3.2 log CFU/g lower than the control group until slaughter [71].

***Salmonella* spp.**

Salmonella is one of the most problematic pathogenic bacteria in poultry, along with *C. jejuni* [72]. There are two well-known species of *Salmonella*, *Salmonella enterica* and *Salmonella bongori*, with *S. enterica* further divided into six subspecies containing over 2,600 serotypes. In particular, the nontyphoidal serotypes, unlike typhoidal serovars that can only be transmitted among humans, pose a zoonotic threat, being capable of common infections between animals and humans, thereby giving rise to potential public health concerns [44]. In the poultry industry, the most frequently occurring serotypes implicated in the majority of *Salmonella* infection incidents are *S. Enteritidis*, *S. Typhimurium*, and *S. Gallinarum* [73]. Research on *Salmonella* phages, similar to the studies

conducted on *C. jejuni*, has been extensively conducted to suppress this bacterium without relying on antibiotics.

In a previous study conducted to reduce contamination by foodborne pathogens in poultry products and decrease the incidence, severity, and mortality of diseases, a concentration of 10^8 CFU of *S. Enteritidis* PT4 culture in 100 μ L was orally administered to chickens. Subsequently, a phage cocktail consisting of CNPSA1, CNPSA3, and CNPSA4 at a concentration of 10^{11} PFU was administered orally as a single dose. Thus, a significant reduction of pathogenic bacterial populations in the digestive tract was observed with the single administration of a high-titer bacteriophage mixture. Furthermore, a reduction of 3.5 log levels in colony-forming units of *S. Enteritidis* PT4 per gram of cecal content was observed [74]. In prior research, an oral challenge with *S. Gallinarum* at a concentration of 5×10^8 CFU/mL was conducted. Following the challenge, bacteriophage CJø01 was used as a feed additive at a concentration of 10^6 PFU/kg. The administration of these bacteriophages to chickens that had contact with infected individuals resulted in a mortality rate of only 5%, compared with 30% in the group that did not receive phage therapy [75]. Finally, an oral challenge with a suspension of *S. Typhimurium* at a concentration of approximately 10^6 CFU/mL, administered in a volume of 0.5 mL, was conducted concurrently with oral administration of a bacteriophage cocktail (S2a, S9, S11) at a concentration of 10^6 PFU to 4–6 and 8–10-day-old chickens. The results revealed a 10-fold reduction in bacterial counts in the chicken's ileum, cecum, liver, and spleen. Furthermore, when a commercial probiotic supplement was administered orally along with the bacteriophage cocktail, a synergistic antibacterial effect was observed [76].

PHAGE THERAPY IN COMPANION ANIMALS

Canine

Pseudomonas aeruginosa

The first *in vivo* experimental investigation using phage therapy to treat external otitis caused by *Pseudomonas aeruginosa* in dogs was reported recently. In the experiment, 10 dogs with a preexisting case of *P. aeruginosa*-related external otitis were used. The researchers isolated phages and then selected which virus would be most effective in killing the strains often seen in canine otitis to create a phage cocktail for administration to the animals. For the *in vivo* trial, the researchers opted for the phage combination that killed 90% or more of the *P. aeruginosa* strains tested [77,78].

Escherichia coli

The effectiveness of 40 bacteriophages against 53 uropathogenic *E. coli* (UPEC) strains was tested *in vitro* by a New Zealand research team. Phage mixtures have been shown to efficiently lyse over 90% of bacterial strains, suggesting that such mixtures may constitute a legitimate alternative to traditional medicines in the case of urinary infections [79].

Staphylococcus pseudintermedius

Methicillin-resistant *S. pseudintermedius* (MRSP) strains are becoming increasingly common and are limiting the effectiveness of antibiotic molecules in treating infections in dogs. Thus, a team of researchers has recently set out to isolate and characterize specific bacteriophages for MRSP strains. Dog feces bacteriophages were tested on 66 different strains of *S. pseudintermedius* to determine their host range (17 strains resistant to methicillin, 43 sensitive to methicillin, and 6 isolated directly from dogs). When tested against a panel of resistant bacteria, all phages demonstrated lytic activity; however, they killed only 16%–28% of susceptible strains [80–82].

Leishmania infantum

This study reported that *L. infantum* did not use a serological panel that included samples from dogs infected with *Leishmania braziliensis* because the researchers collected dog serum samples only in the Belo Horizonte metropolitan area, where a comparatively low incidence of *Leishmania* infection in dogs has been recently described. Therefore, the findings of this study should be used as proof-of-concept of the ability of the suggested synthetic antigens to help in the serodiagnosis of canine visceral leishmaniasis (CVL) and may serve as a reference for future tests. However, owing to their simplicity, ease of use, repeatability, and cheap cost, these new extremely accurate phage-fused peptides and their application in phage-enzyme-linked immunosorbent assays (ELISAs) may be swiftly used in the serodiagnosis used in CVL-monitoring programs [83].

Feline

Escherichia coli

UPEC, the primary infectious cause of urinary tract infections in dogs, cats, and humans, is a serious issue. This study aimed to determine whether a large variety of phages capable of lysing canine and feline UPEC exist in nature. Using 53 UPEC strains and 7 *E. coli* strains obtained from dog and cat feces, the ability of phages to cause lysis *in vitro* was studied. Between January 2002 and April 2004, four New Zealand animal health laboratories were contacted to obtain a total of 31 canine UPEC strains and 22 feline UPEC strains. Once lysis profiles suggested that a specific phage may be a potential candidate for *in vivo* testing, electron microscopy and DNA sequencing were used to perform more morphological and genetic studies on the phage in issue.

The majority of the canine and feline UPEC strains were lysed *in vitro* by naturally occurring phages that may be obtained by visiting a sewage treatment plant only once, according to the findings of the previous experiments. Of the UPEC strains that were examined, 17%–72% were eradicated by certain phages. More than half of all UPEC strains were lysed by the top 10 phages with the greatest host versatility. The phages were able to lyse 92% of the UPEC strains when combined, and each had a distinct lysis profile. In contrast to a previous study on *E. coli* urinary tract infections in children, however, 14 out of 44 phages lysed >15% of the UPEC strains; the present study demonstrates a far more positive result. Drulis-Kawa *et al.* discovered that only three of the phages they studied lysed >50% of the UPEC strains they analyzed [84]. This experiment used *E. coli* phages collected from a hospital collection. The phage propagation on UPEC may have contributed to the increased lysis success. The idea that UPEC isolated from cats and dogs is more susceptible to phage lysis than UPEC obtained from children cannot be eliminated at this time [79,85].

Staphylococci

Most commonly found on the skin of animals, staphylococci are Gram-positive cocci that are asymptomatic, do not produce spores, and are facultative anaerobes. Two staphylococcal species, *S. aureus* and *Staphylococcus schleiferi*, are becoming more prevalent as they are harmful to rodents and birds. *Staphylococcus intermedius* has been linked to instances of superficial dermatitis, bacterial folliculitis, and superficial pyoderma in cats, although staphylococci do not appear to be a prominent cause of any particular illnesses in cats. An array of drugs used to treat staphylococcal infections in canines and felines has contributed to the development of drug-resistant forms of the bacteria. Penicillins, cephalosporins, macrolides, lincosamides, fusidic acid, tetracyclines, chloramphenicol, potentiated sulfonamides, aminoglycosides, and fluoroquinolones are all regularly used antibiotics in canine and feline treatment.

Nosocomial strains of staphylococci have exhibited rapid development of resistance to practically

all the main types of antibiotics since their widespread usage became commercialized. Not much is understood about how staphylococci in cats and dogs become resistant to antibiotics. Conversely, plasmids are anticipated to play a crucial role in the dissemination of antibiotic resistance in these species. Either transposable elements or plasmids may serve as resistance gene vectors. Some studies have shown evidence of horizontal gene transfer between *S. aureus* and *S. intermedius* and between *S. aureus* and coagulase-negative staphylococci. Staphylococci are genetically diverse because of their interactions with a broad variety of other bacteria on the skin, mucosal surfaces, and respiratory tracts of mammals, all of which provide them with a rich source of genetic diversity. In a complex microbial community, resistance genes may be acquired rapidly. Staphylococci thrive on mammalian hosts, making them easily transmitted between animals and even to humans through cat and dog bites. There are extremely few accounts in the literature of staphylococci spreading from one species to another since staphylococci tend to be very species-specific. For example, cattle and cats are susceptible to infection from human *S. aureus*, presumably *S. epidermidis* strains and antibiotic-resistant staphylococci [86–89].

SUMMARY AND CONCLUSION

Since the discovery of antibiotics, phage therapy has not received considerable attention. However, the emergence of superbugs has emphasized the importance of bacteriophages in various industries. This study focuses on the use of phages in animal farming, specifically in the livestock and companion animal industries. In livestock farming, phages can be used throughout the entire production chain, with a particular emphasis on bacterial control in farms vulnerable to pathogenic bacteria. As animals in farming industries cannot manage hygiene on their own, external interventions are crucial to remove pathogenic factors [90]. Historically, antibiotics were efficient substances not only in eliminating bacteria but also in promoting animal growth [91,92]. Nevertheless, the rise of multidrug-resistant bacteria has prompted research into the potential of bacteriophages as an alternative to antibiotics [93–95]. Recent experiments have demonstrated the synergistic effect of bacteriophages and probiotics in controlling pathogenic bacteria and promoting animal growth [96–98]. Although the societal perception of phage use in humans remains negative, its application in animal farming is relatively acceptable, with some phage uses being legally permitted. In conclusion, research on phage therapy in animal farming is expected to become more active in the future.

Phage use in the livestock industry is strategically focused on vulnerable animals during their early stages. For instance, research indicates that a low concentration of *Salmonella* infection can be lethal to piglets but has a minimal impact on mature pigs [99]. Accordingly, in livestock, phage treatments primarily aim at preventing pathogenic bacterial infections and growth performance [100,101]. In contrast, with companion animals that have assured individual hygiene, the focus rests on bacterial diseases arising from their unique characteristics and lifestyle habits. This study observes a similar trend, where livestock research primarily centers around intestinal diseases directly linked to growth, such as *Salmonella* and *E. coli* [40,102–104]. In companion animals, studies extend beyond intestinal diseases to encompass conditions such as otitis and skin disorders [77,78]. Chronic illnesses, while quickly treatable with antibiotic products, pose the risk of developing antibiotic resistance with continuous use. Additionally, antibiotic treatments can disrupt the balance of symbiotic microorganisms by reducing beneficial bacteria [27,105]. Conversely, phage therapy alleviates concerns about antibiotic resistance, and according to the literature, phage resistance often lowers the pathogenicity of bacteria, significantly reducing the associated risks [106–108]. Consequently, while research in phage therapy for companion animals may be in its early stages, the

growing societal recognition of pets as integral family members suggests that phage therapy holds potential as a research topic for treating various diseases in companion animals [109,110].

The extensive use of phage treatment, notwithstanding its benefits, raises new issues that extend beyond the scope of traditional medical practice and need new approaches. These include the need to increase phage strains in reference phage banks, the development of efficient phage screening techniques for the rapid identification of therapeutic phages, the design of effective treatment strategies that target infectious biofilms, the establishment of manufacturing procedures that ensure the quality and safety of preparations, and the assurance of the stability of preparations during storage and transportation. As infectious illnesses know no boundaries, a global action plan is required to make phage treatment accessible globally. This clearly necessitates active cooperation across nations to overcome logistical and regulatory obstacles, as well as between physicians and scientists to close the present knowledge gap and progress the area.

REFERENCES

1. Song D, Lee J, Kim K, Oh H, An J, Chang S, et al. Effects of dietary supplementation of *Pediococcus pentosaceus* strains from kimchi in weaned piglet challenged with *Escherichia coli* and *Salmonella enterica*. *J Anim Sci Technol*. 2023;65:611–26. <https://doi.org/10.5187/jast.2023.e31>
2. Lee J, Oh S, Kim M. Response to environmental enrichment of weanling pigs on growth, behaviour and welfare after weaning. *J Anim Sci Technol*. Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e128>
3. Lim CI, Choo HJ, Park JH. Effect of phytase supplementation on performance, fecal excretion, and compost characteristics in broilers fed diets deficient in phosphorus and calcium. *J Anim Sci Technol*. Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e59>
4. Hwang S, Lee W, Lee Y. Development of a nucleic acid detection method based on the CRISPR-Cas13 for point-of-care testing of bovine viral diarrhoea virus-1b. *J Anim Sci Technol*. Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e77>
5. Song D, Lee J, Yoo Y, Oh H, Chang S, An J, et al. Effects of probiotics on growth performance, intestinal morphology, intestinal microbiota weaning pig challenged with *Escherichia coli* and *Salmonella enterica*. *J Anim Sci Technol*. Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e119>
6. Oh H, Yoon Y, Yoon JW, Oh SW, Lee S, Lee H. Quantitative risk assessment of foodborne *Salmonella* illness by estimating cooking effect on eggs from retail markets. *J Anim Sci Technol*. 2023;65:1024–39. <https://doi.org/10.5187/jast.2023.e18>
7. Sykes JE. *Canine and feline infectious diseases*. London: Elsevier Health Sciences; 2013.
8. Abebe R, Hatiya H, Abera M, Megersa B, Asmare K. Bovine mastitis: prevalence, risk factors and isolation of *Staphylococcus aureus* in dairy herds at Hawassa milk shed, South Ethiopia. *BMC Vet Res*. 2016;12:270. <https://doi.org/10.1186/s12917-016-0905-3>
9. Raza M, Eungyung K, Shakeel M, Fiaz M, Ma L, Kim H, et al. Evaluation of zinc oxide and copper oxide nanoparticles as potential alternatives to antibiotics for managing fowl typhoid in broilers. *J Anim Sci Technol*. Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e91>
10. Zhong Y, Zuo B, Li J, Zhai Y, Mudarra R. Effects of paraformic acid supplementation, as an antibiotic replacement, on growth performance, intestinal morphology and gut microbiota of nursery pigs. *J Anim Sci Technol*. Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e95>
11. Keum GB, Kim ES, Cho J, Song M, Oh KK, Cho JH, et al. Analysis of antibiotic resistance genes in pig feces during the weaning transition using whole metagenome shotgun

- sequencing. *J Anim Sci Technol.* 2023;65:175-82. <https://doi.org/10.5187/jast.2022.e103>
12. Pandey S, Doo H, Keum GB, Kim ES, Kwak J, Ryu S, et al. Antibiotic resistance in livestock, environment and humans: one health perspective. *J Anim Sci Technol.* Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e129>
 13. van Duin D, Paterson DL. Multidrug-resistant bacteria in the community: trends and lessons learned. *Infect Dis Clin North Am.* 2016;30:377-90. <https://doi.org/10.1016/j.idc.2016.02.004>
 14. Jo H, Han G, Kim EB, Kong C, Kim BG. Effects of supplemental bacteriophage on the gut microbiota and nutrient digestibility of ileal-cannulated pigs. *J Anim Sci Technol.* Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e96>
 15. Han S, Elnar A, Lim C, Kim GB. Complete genome sequence of bacteriocin-producing *Ligilactobacillus salivarius* B4311 isolated from fecal samples of broiler chicken with anti-listeria activity. *J Anim Sci Technol.* Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e40>
 16. Sampath V, Park JH, Pineda L, Han Y, Cho S, Kim IH. Sows fed with synergistic blend of short- and medium chain organic acid has a carryover effect on post-weaning growth rate. *J Anim Sci Technol.* 2022;64:302-11. <https://doi.org/10.5187/jast.2022.e11>
 17. Jeon K, Song M, Lee J, Oh H, Chang S, Song D, et al. Effects of single and complex probiotics in growing-finishing pigs and swine compost. *J Anim Sci Technol.* Forthcoming. 2023. <https://doi.org/10.5187/jast.2023.e88>
 18. de Jonge PA, Nobrega FL, Brouns SJJ, Dutilh BE. Molecular and evolutionary determinants of bacteriophage host range. *Trends Microbiol.* 2019;27:51-63. <https://doi.org/10.1016/j.tim.2018.08.006>
 19. Lenski RE, Levin BR. Constraints on the coevolution of bacteria and virulent phage: a model, some experiments, and predictions for natural communities. *Am Nat.* 1985;125:585-602. <https://doi.org/10.1086/284364>
 20. Gustafson RH. Use of antibiotics in livestock and human health concerns. *J Dairy Sci.* 1991;74:1428-32. [https://doi.org/10.3168/jds.S0022-0302\(91\)78299-4](https://doi.org/10.3168/jds.S0022-0302(91)78299-4)
 21. Vivas R, Barbosa AAT, Dolabela SS, Jain S. Multidrug-resistant bacteria and alternative methods to control them: an overview. *Microb Drug Resist.* 2019;25:890-908. <https://doi.org/10.1089/mdr.2018.0319>
 22. Gordillo Altamirano FL, Barr JJ. Phage therapy in the postantibiotic era. *Clin Microbiol Rev.* 2019;32:e00066-18. <https://doi.org/10.1128/CMR.00066-18>
 23. APPA [American Pet Products Association]. Pet industry market size, trends & ownership statistics [Internet]. 2023 [cited 2023 Nov 12]. <https://www.americanpetproducts.org/news/News-Public-Relations/pet-industry-market-size-trends-ownership-statistics>
 24. Alexander P, Berri A, Moran D, Reay D, Rounsevell MDA. The global environmental paw print of pet food. *Glob Environ Change.* 2020;65:102153. <https://doi.org/10.1016/j.gloenvcha.2020.102153>
 25. Joksimovic M, Ford BA, Lazic T, Soldatovic I, Luzetsky S, Grozdanic S. Antibiotic recommendations for treatment of canine stromal corneal ulcers. *Vet Sci.* 2023;10:66. <https://doi.org/10.3390/vetsci10020066>
 26. Cho HW, Choi S, Seo K, Kim KH, Jeon JH, Kim CH, et al. Gut microbiota profiling in aged dogs after feeding pet food contained *Hericium erinaceus*. *J Anim Sci Technol.* 2022;64:937-49. <https://doi.org/10.5187/jast.2022.e66>
 27. Duan H, Yu L, Tian F, Zhai Q, Fan L, Chen W. Antibiotic-induced gut dysbiosis and barrier disruption and the potential protective strategies. *Crit Rev Food Sci Nutr.* 2022;62:1427-52. <https://doi.org/10.1080/10408398.2020.1843396>

28. Loponte R, Pagnini U, Iovane G, Pisanelli G. Phage therapy in veterinary medicine. *Antibiotics*. 2021;10:421. <https://doi.org/10.3390/antibiotics10040421>
29. Ferriol-González C, Domingo-Calap P. Phage therapy in livestock and companion animals. *Antibiotics*. 2021;10:559. <https://doi.org/10.3390/antibiotics10050559>
30. Zduńczyk S, Janowski T. Bacteriophages and associated endolysins in therapy and prevention of mastitis and metritis in cows: current knowledge. *Anim Reprod Sci*. 2020;218:106504. <https://doi.org/10.1016/j.anireprosci.2020.106504>
31. Cheng WN, Han SG. Bovine mastitis: risk factors, therapeutic strategies, and alternative treatments — a review. *Asian-Australas J Anim Sci*. 2020;33:1699-713. <https://doi.org/10.5713/ajas.20.0156>
32. Kim SD, Kim GB, Lee GY, Yang SJ. Multilocus sequence type-dependent activity of human and animal cathelicidins against community-, hospital-, and livestock-associated methicillin-resistant *Staphylococcus aureus* isolates. *J Anim Sci Technol*. 2022;64:515-30. <https://doi.org/10.5187/jast.2022.e32>
33. Heikkilä AM, Liski E, Pyörälä S, Taponen S. Pathogen-specific production losses in bovine mastitis. *J Dairy Sci*. 2018;101:9493-504. <https://doi.org/10.3168/jds.2018-14824>
34. Gilbert FB, Cunha P, Jensen K, Glass EJ, Foucras G, Robert-Granié C, et al. Differential response of bovine mammary epithelial cells to *Staphylococcus aureus* or *Escherichia coli* agonists of the innate immune system. *Vet Res*. 2013;44:40. <https://doi.org/10.1186/1297-9716-44-40>
35. Breyne K, Honaker RW, Hobbs Z, Richter M, Żaczek M, Spangler T, et al. Efficacy and safety of a bovine-associated *Staphylococcus aureus* phage cocktail in a murine model of mastitis. *Front Microbiol*. 2017;8:2348. <https://doi.org/10.3389/fmicb.2017.02348>
36. Teng F, Xiong X, Zhang S, Li G, Wang R, Zhang L, et al. Efficacy assessment of phage therapy in treating *Staphylococcus aureus*-induced mastitis in mice. *Viruses*. 2022;14:620. <https://doi.org/10.3390/v14030620>
37. Gill JJ, Pacan JC, Carson ME, Leslie KE, Griffiths MW, Sabour PM. Efficacy and pharmacokinetics of bacteriophage therapy in treatment of subclinical *Staphylococcus aureus* mastitis in lactating dairy cattle. *Antimicrob Agents Chemother*. 2006;50:2912-8. <https://doi.org/10.1128/AAC.01630-05>
38. Fan J, Zeng Z, Mai K, Yang Y, Feng J, Bai Y, et al. Preliminary treatment of bovine mastitis caused by *Staphylococcus aureus*, with trx-SA1, recombinant endolysin of *S. aureus* bacteriophage IME-SA1. *Vet Microbiol*. 2016;191:65-71. <https://doi.org/10.1016/j.vetmic.2016.06.001>
39. Porter J, Anderson J, Carter L, Donjacour E, Paros M. In vitro evaluation of a novel bacteriophage cocktail as a preventative for bovine coliform mastitis. *J Dairy Sci*. 2016;99:2053-62. <https://doi.org/10.3168/jds.2015-9748>
40. Guo M, Gao Y, Xue Y, Liu Y, Zeng X, Cheng Y, et al. Bacteriophage cocktails protect dairy cows against mastitis caused by drug resistant *Escherichia coli* infection. *Front Cell Infect Microbiol*. 2021;11:690377. <https://doi.org/10.3389/fcimb.2021.690377>
41. Al Mawly J, Grinberg A, Prattley D, Moffat J, Marshall J, French N. Risk factors for neonatal calf diarrhoea and enteropathogen shedding in New Zealand dairy farms. *Vet J*. 2015;203:155-60. <https://doi.org/10.1016/j.tvjl.2015.01.010>
42. Kim HS, Whon TW, Sung H, Jeong YS, Jung ES, Shin NR, et al. Longitudinal evaluation of fecal microbiota transplantation for ameliorating calf diarrhea and improving growth performance. *Nat Commun*. 2021;12:161. <https://doi.org/10.1038/s41467-020-20389-5>
43. Alomari MMM, Dec M, Nowaczek A, Puchalski A, Wernicki A, Kowalski C, et al.

- Therapeutic and prophylactic effect of the experimental bacteriophage treatment to control diarrhea caused by *E. coli* in Newborn calves. *ACS Infect Dis.* 2021;7:2093-101. <https://doi.org/10.1021/acsinfecdis.1c00010>
44. Won K, Kim D, Shin D, Hur J, Lee HK, Heo J, et al. High-throughput sequencing-based metagenomic and transcriptomic analysis of intestine in piglets infected with *Salmonella*. *J Anim Sci Technol.* 2022;64:1144-72. <https://doi.org/10.5187/jast.2022.e73>
 45. Thomson DMP, Krupsey J, Freedman SO, Gold P. The radioimmunoassay of circulating carcinoembryonic antigen of the human digestive system. *Proc Natl Acad Sci.* 1969;64:161-7. <https://doi.org/10.1073/pnas.64.1.161>
 46. Conceição-Neto N, Theuns S, Cui T, Zeller M, Yinda CK, Christiaens I, et al. Identification of an enterovirus recombinant with a torovirus-like gene insertion during a diarrhea outbreak in fattening pigs. *Virus Evol.* 2017;3:vex024. <https://doi.org/10.1093/ve/vex024>
 47. Nantel-Fortier N, Gauthier M, LHomme Y, Lachapelle V, Fravallo P, Brassard J. The swine enteric virome in a commercial production system and its association with neonatal diarrhea. *Vet Microbiol.* 2022;266:109366. <https://doi.org/10.1016/j.vetmic.2022.109366>
 48. Wall SK, Zhang J, Rostagno MH, Ebner PD. Phage therapy to reduce preprocessing *Salmonella* infections in market-weight swine. *Appl Environ Microbiol.* 2010;76:48-53. <https://doi.org/10.1128/AEM.00785-09>
 49. Borie C, Albala I, Sánchez P, Sánchez ML, Ramírez S, Navarro C, et al. Bacteriophage treatment reduces *Salmonella* colonization of infected chickens. *Avian Dis.* 2008;52:64-7. <https://doi.org/10.1637/8091-082007-Reg>
 50. Jończyk E, Kłak M, Międzybrodzki R, Górski A. The influence of external factors on bacteriophages—review. *Folia Microbiol.* 2011;56:191-200. <https://doi.org/10.1007/s12223-011-0039-8>
 51. Thanki AM, Hooton S, Gigante AM, Atterbury RJ, Clokie MRJ. Potential roles for bacteriophages in reducing *Salmonella* from poultry and swine. In: Lamas A, Regal P, Franco M, editors. *Salmonella spp: a global challenge*. London: IntechOpen; 2021.
 52. Vinner GK, Richards K, Leppanen M, Sagona AP, Malik DJ. Microencapsulation of enteric bacteriophages in a pH-responsive solid oral dosage formulation using a scalable membrane emulsification process. *Pharmaceutics.* 2019;11:475. <https://doi.org/10.3390/pharmaceutics11090475>
 53. Zhang Y, Zhang H, Ghosh D. The stabilizing excipients in dry state therapeutic phage formulations. *AAPS PharmSciTech.* 2020;21:133. <https://doi.org/10.1208/s12249-020-01673-5>
 54. Chang RY, Wong J, Mathai A, Morales S, Kutter E, Britton W, et al. Production of highly stable spray dried phage formulations for treatment of *Pseudomonas aeruginosa* lung infection. *Eur J Pharm Biopharm.* 2017;121:1-13. <https://doi.org/10.1016/j.ejpb.2017.09.002>
 55. Vinner GK, Rezaie-Yazdi Z, Leppanen M, Stapley AGF, Leaper MC, Malik DJ. Microencapsulation of *Salmonella*-specific bacteriophage Felix O1 using spray-drying in a pH-responsive formulation and direct compression tableting of powders into a solid oral dosage form. *Pharmaceutics.* 2019;12:43. <https://doi.org/10.3390/ph12010043>
 56. Saez AC, Zhang J, Rostagno MH, Ebner PD. Direct feeding of microencapsulated bacteriophages to reduce *Salmonella* colonization in pigs. *Foodborne Pathog Dis.* 2011;8:1269-74. <https://doi.org/10.1089/fpd.2011.0905>
 57. Albino LAA, Rostagno MH, Húngaro HM, Mendonça RCS. Isolation, characterization, and application of bacteriophages for *Salmonella* spp. biocontrol in pigs. *Foodborne Pathog Dis.* 2014;11:602-9. <https://doi.org/10.1089/fpd.2013.1600>

58. Han SJ, Oh Y, Lee CY, Han JH. Efficacy of dietary supplementation of bacteriophages in treatment of concurrent infections with enterotoxigenic *Escherichia coli* K88 and K99 in postweaning pigs. *J Swine Health Prod.* 2016;24:259-63.
59. Lin Y, Zhou B, Zhu W. Pathogenic *Escherichia coli*-specific bacteriophages and polyvalent bacteriophages in piglet guts with increasing coliphage numbers after weaning. *Appl Environ Microbiol.* 2021;87:e00966-21. <https://doi.org/10.1128/AEM.00966-21>
60. Oosterom J, Notermans S, Karman H, Engels GB. Origin and prevalence of *Campylobacter jejuni* in poultry processing. *J Food Prot.* 1983;46:339-44. <https://doi.org/10.4315/0362-028X-46.4.339>
61. Hansson I, Sandberg M, Habib I, Lowman R, Engvall EO. Knowledge gaps in control of *Campylobacter* for prevention of campylobacteriosis. *Transbound Emerg Dis.* 2018;65:30-48. <https://doi.org/10.1111/tbed.12870>
62. Duncan DL. Gastroenteritis: an overview of the symptoms, transmission and management. *Br J Sch Nurs.* 2018;13:484-8. <https://doi.org/10.12968/bjns.2018.13.10.484>
63. Acheson D, Allos BM. *Campylobacter jejuni* infections: update on emerging issues and trends. *Clin Infect Dis.* 2001;32:1201-6. <https://doi.org/10.1086/319760>
64. Atterbury RJ, Connerton PL, Dodd CER, Rees CED, Connerton IF. Application of host-specific bacteriophages to the surface of chicken skin leads to a reduction in recovery of *Campylobacter jejuni*. *Appl Environ Microbiol.* 2003;69:6302-6. <https://doi.org/10.1128/AEM.69.10.6302-6306.2003>
65. Goode D, Allen VM, Barrow PA. Reduction of experimental *Salmonella* and *Campylobacter* contamination of chicken skin by application of lytic bacteriophages. *Appl Environ Microbiol.* 2003;69:5032-6. <https://doi.org/10.1128/AEM.69.8.5032-5036.2003>
66. Bigwood T, Hudson JA, Billington C, Carey-Smith GV, Heinemann JA. Phage inactivation of foodborne pathogens on cooked and raw meat. *Food Microbiol.* 2008;25:400-6. <https://doi.org/10.1016/j.fm.2007.11.003>
67. Orquera S, Götz G, Hertwig S, Hammerl J, Sparborth D, Joldic A, et al. Control of *Campylobacter* spp. and *Yersinia enterocolitica* by virulent bacteriophages. *J Mol Genet Med.* 2012;6:273-8. <https://doi.org/10.4172/1747-0862.1000049>
68. Wagenaar JA, Mevius DJ, Havelaar AH. *Campylobacter* in primary animal production and control strategies to reduce the burden of human campylobacteriosis. *Rev Sci Tech Off Int Epizoot.* 2006;25:581-94.
69. Loc Carrillo C, Atterbury RJ, El-Shibiny A, Connerton PL, Dillon E, Scott A, et al. Bacteriophage therapy to reduce *Campylobacter jejuni* colonization of broiler chickens. *Appl Environ Microbiol.* 2005;71:6554-63. <https://doi.org/10.1128/AEM.71.11.6554-6563.2005>
70. Basso B, Amato M, Bitella G, Rossi R, Kravchenko A, Sartori L, et al. Two-dimensional spatial and temporal variation of soil physical properties in tillage systems using electrical resistivity tomography. *Agron J.* 2010;102:440-9. <https://doi.org/10.2134/agronj2009.0298>
71. Kittler S, Fischer S, Abdulmawjood A, Glünder G, Klein G. Effect of bacteriophage application on *Campylobacter jejuni* loads in commercial broiler flocks. *Appl Environ Microbiol.* 2013;79:7525-33. <https://doi.org/10.1128/AEM.02703-13>
72. Felix L, Moreira Filho A, Santos MR, Saraiva M, Freitas Neto O, Givisiez P, et al. Intestinal morphometric changes associated with the use of non-antibiotic feed additives in broiler chicks challenged with *Salmonella* Enteritidis. *J Anim Sci Technol.* Forthcoming 2023. <https://doi.org/10.5187/jast.2023.e113>
73. Tariq S, Samad A, Hamza M, Ahmer A, Muazzam A, Ahmad S, et al. *Salmonella* in poultry; an overview. *Int J Multidiscip Sci Arts.* 2022;1:80-4. <https://doi.org/10.47709/ijmdsa>.

- v1i1.1706
74. Fiorentin L, Vieira ND, Barioni W Jr. Oral treatment with bacteriophages reduces the concentration of Salmonella Enteritidis PT4 in caecal contents of broilers. *Avian Pathol.* 2005;34:258-63. <https://doi.org/10.1080/01445340500112157>
 75. Lim TH, Lee DH, Lee YN, Park JK, Youn HN, Kim MS, et al. Efficacy of bacteriophage therapy on horizontal transmission of Salmonella Gallinarum on commercial layer chickens. *Avian Dis.* 2011;55:435-8. <https://doi.org/10.1637/9599-111210-Reg.1>
 76. Toro H, Price SB, McKee AS, Hoerr FJ, Krehling J, Perdue M, et al. Use of bacteriophages in combination with competitive exclusion to reduce Salmonella from infected chickens. *Avian Dis.* 2005;49:118-24. <https://doi.org/10.1637/7286-100404R>
 77. Hawkins C, Harper D, Burch D, Ånggård E, Soothill J. Topical treatment of Pseudomonas aeruginosa otitis of dogs with a bacteriophage mixture: a before/after clinical trial. *Vet Microbiol.* 2010;146:309-13. <https://doi.org/10.1016/j.vetmic.2010.05.014>
 78. Furusawa T, Iwano H, Higuchi H, Yokota H, Usui M, Iwasaki T, et al. Bacteriophage can lyse antibiotic-resistant Pseudomonas aeruginosa isolated from canine diseases. *J Vet Med Sci.* 2016;78:1035-8. <https://doi.org/10.1292/jvms.15-0310>
 79. Freitag T, Squires RA, Schmid J. Naturally occurring bacteriophages lyse a large proportion of canine and feline uropathogenic Escherichia coli isolates in vitro. *Res Vet Sci.* 2008;85:1-7. <https://doi.org/10.1016/j.rvsc.2007.09.004>
 80. Moodley A, Kot W, Nälgård S, Jakociune D, Neve H, Hansen LH, et al. Isolation and characterization of bacteriophages active against methicillin-resistant Staphylococcus pseudintermedius. *Res Vet Sci.* 2019;122:81-5. <https://doi.org/10.1016/j.rvsc.2018.11.008>
 81. Lynch SA, Helbig KJ. The complex diseases of Staphylococcus pseudintermedius in canines: where to next? *Vet Sci.* 2021;8:11. <https://doi.org/10.3390/vetsci8010011>
 82. Urban-Chmiel R, Balicki I, Świąder K, Nowaczek A, Pyzik E, Stepien-Pyśniak D, et al. The in vitro efficacy of eye drops containing a bacteriophage solution specific for Staphylococcus spp. isolated from dogs with bacterial conjunctivitis. *Ir Vet J.* 2020;73:21. <https://doi.org/10.1186/s13620-020-00175-x>
 83. Costa LE, Lima MI, Chávez-Fumagalli MA, Menezes-Souza D, Martins VT, Duarte MC, et al. Subtractive phage display selection from canine visceral leishmaniasis identifies novel epitopes that mimic Leishmania infantum antigens with potential serodiagnosis applications. *Clin Vaccine Immunol.* 2014;21:96-106. <https://doi.org/10.1128/CVI.00583-13>
 84. Drulis-Kawa Z, Weber-Dabrowska B, Lewczyk E, Jankowski S, Doroszkiewicz W. The sensitivity of the uropathogenic Escherichia coli strains to antibiotics, bacteriophages and bactericidal serum activity. *Pol Merkur Lekarski.* 2002;13:470-2.
 85. Russo TA, Johnson JR. Medical and economic impact of extraintestinal infections due to Escherichia coli: focus on an increasingly important endemic problem. *Microbes Infect.* 2003;5:449-56. [https://doi.org/10.1016/S1286-4579\(03\)00049-2](https://doi.org/10.1016/S1286-4579(03)00049-2)
 86. Conner JG, Smith J, Erol E, Locke S, Phillips E, Carter CN, et al. Temporal trends and predictors of antimicrobial resistance among Staphylococcus spp. isolated from canine specimens submitted to a diagnostic laboratory. *PLOS ONE.* 2018;13:e0200719. <https://doi.org/10.1371/journal.pone.0200719>
 87. Feßler AT, Scholtzek AD, Schug AR, Kohn B, Weingart C, Schink AK, et al. Antimicrobial and biocide resistance among feline and canine Staphylococcus aureus and Staphylococcus pseudintermedius isolates from diagnostic submissions. *Antibiotics.* 2022;11:127. <https://doi.org/10.3390/antibiotics11020127>
 88. Machado AB, Machado MFR, Picoli SU. An investigation of methicillin-resistant

- Staphylococcus pseudintermedius* (MRSP) in domestic and shelter dogs in Montenegro (RS-Brazil). *Rev Bras Saúde Prod Anim.* 2017;18:542-8. <https://doi.org/10.1590/S1519-99402017000400005>
89. Malik S, Peng H, Barton MD. Antibiotic resistance in staphylococci associated with cats and dogs. *J Appl Microbiol.* 2005;99:1283-93. <https://doi.org/10.1111/j.1365-2672.2005.02699.x>
 90. Sureshkumar S, Park JH, Kim IH. A preliminary evaluation on mixed probiotics as an antimicrobial spraying agent in growing pig barn. *J Anim Sci Technol.* 2022;64:1035-45. <https://doi.org/10.5187/jast.2022.e69>
 91. Butaye P, Devriese LA, Haesebrouck F. Antimicrobial growth promoters used in animal feed: effects of less well known antibiotics on gram-positive bacteria. *Clin Microbiol Rev.* 2003;16:175-88. <https://doi.org/10.1128/cmr.16.2.175-188.2003>
 92. Graham JP, Boland JJ, Silbergeld E. Growth promoting antibiotics in food animal production: an economic analysis. *Public Health Rep.* 2007;122:79-87. <https://doi.org/10.1177/003335490712200111>
 93. Manohar P, Loh B, Athira S, Nachimuthu R, Hua X, Welburn SC, et al. Secondary bacterial infections during pulmonary viral disease: phage therapeutics as alternatives to antibiotics? *Front Microbiol.* 2020;11:1434. <https://doi.org/10.3389/fmicb.2020.01434>
 94. Romero-Calle D, Guimarães Benevides R, Góes-Neto A, Billington C. Bacteriophages as alternatives to antibiotics in clinical care. *Antibiotics.* 2019;8:138. <https://doi.org/10.3390/antibiotics8030138>
 95. Kim H, Cho JH, Cho JH, Song M, Shin H, Kim S, et al. Complete genome sequence of *Salmonella enterica* strain K_SA184, multidrug resistance bacterium isolated from lamb (Ovis aries). *J Anim Sci Technol.* 2021;63:194-7. <https://doi.org/10.5187/jast.2021.e6>
 96. Porter SB, Johnston BD, Kisiela D, Clabots C, Sokurenko EV, Johnson JR. Bacteriophage cocktail and microcin-producing probiotic *Escherichia coli* protect mice against gut colonization with multidrug-resistant *Escherichia coli* sequence type 131. *Front Microbiol.* 2022;13:887799. <https://doi.org/10.3389/fmicb.2022.887799>
 97. Zadeh RG, Zadeh MH, Sholeh M, Asgharzadeh S, Darbandi A, Fashami PA, et al. Bacteriophage and probiotic therapy in the treatment of *Pseudomonas aeruginosa* burn infection and their synergistic effect. *Egypt J Exp Biol.* 2023;19:21-9. <https://doi.org/10.5455/egyjebb.20221207072545>
 98. Shaufi MAM, Sieo CC, Chong CW, Geok Hun T, Omar AR, Han Ming G, et al. Effects of phage cocktail, probiotics, and their combination on growth performance and gut microbiota of broiler chickens. *Animals.* 2023;13:1328. <https://doi.org/10.3390/ani13081328>
 99. Bernad-Roche M, Casanova-Higes A, Marín-Alcalá CM, Cebollada-Solanas A, Mainar-Jaime RC. *Salmonella* infection in nursery piglets and its role in the spread of salmonellosis to further production periods. *Pathogens.* 2021;10:123. <https://doi.org/10.3390/pathogens10020123>
 100. Biswas S, Dang DX, Kim IH. Comparison of the effects of zinc oxide and zinc aspartic acid chelate on the performance of weaning pigs. *J Anim Sci Technol.* Forthcoming 2023. <https://doi.org/10.5187/jast.2023.e39>
 101. Park JH, Sureshkumar S, Kim IH. Effects of dietary lysozyme supplementation on growth performance, nutrient digestibility, intestinal microbiota, and blood profiles of weanling pigs challenged with *Escherichia coli*. *J Anim Sci Technol.* 2021;63:501-9. <https://doi.org/10.5187/jast.2021.e54>
 102. Clavijo V, Morales T, Vives-Flores MJ, Reyes Muñoz A. The gut microbiota of chickens in a commercial farm treated with a *Salmonella* phage cocktail. *Sci Rep.* 2022;12:991. <https://doi.org/10.1038/s41598-022-12991-2>

- org/10.1038/s41598-021-04679-6
103. Imklin N, Sriprasong P, Phuttapatimok S, Kaminsonsakul T, Woonwong Y, Jirawattanapong P, et al. In vivo assessment of bacteriophages specific to multidrug resistant *Escherichia coli* on fecal bacterial counts and microbiome in nursery pigs. *Res Vet Sci.* 2022;151:138-48. <https://doi.org/10.1016/j.rvsc.2022.07.012>
 104. Thanki AM, Clavijo V, Healy K, Wilkinson RC, Sicheritz-Pontén T, Millard AD, et al. Development of a phage cocktail to target *Salmonella* strains associated with swine. *Pharmaceuticals.* 2022;15:58. <https://doi.org/10.3390/ph15010058>
 105. McDonnell L, Gilkes A, Ashworth M, Rowland V, Harries TH, Armstrong D, et al. Association between antibiotics and gut microbiome dysbiosis in children: systematic review and meta-analysis. *Gut Microbes.* 2021;13:1870402. <https://doi.org/10.1080/19490976.2020.1870402>
 106. Song L, Yang X, Huang J, Zhu X, Han G, Wan Y, et al. Phage selective pressure reduces virulence of hypervirulent *Klebsiella pneumoniae* through mutation of the *wzc* gene. *Front Microbiol.* 2021;12:739319. <https://doi.org/10.3389/fmicb.2021.739319>
 107. Li J, Yan B, He B, Li L, Zhou X, Wu N, et al. Development of phage resistance in multidrug-resistant *Klebsiella pneumoniae* is associated with reduced virulence: a case report of a personalised phage therapy. *Clin Microbiol Infect.* 2023;29:1601. <https://doi.org/10.1016/j.cmi.2023.08.022>
 108. Markwitz P, Olszak T, Gula G, Kowalska M, Arabski M, Drulis-Kawa Z. Emerging phage resistance in *Pseudomonas aeruginosa* PAO1 is accompanied by an enhanced heterogeneity and reduced virulence. *Viruses.* 2021;13:1332. <https://doi.org/10.3390/v13071332>
 109. Lee D, Goh TW, Kang MG, Choi HJ, Yeo SY, Yang J, et al. Perspectives and advances in probiotics and the gut microbiome in companion animals. *J Anim Sci Technol.* 2022;64:197-217. <https://doi.org/10.5187/jast.2022.e8>
 110. Cho HW, Seo K, Chun JL, Jeon J, Kim CH, Lim S, et al. Effects of resistant starch on anti-obesity status and nutrient digestibility in dogs. *J Anim Sci Technol.* 2023;65:550-61. <https://doi.org/10.5187/jast.2023.e11>
 111. Han JE, Kim JH, Hwang SY, Choresca CH Jr, Shin SP, Jun JW, et al. Isolation and characterization of a Myoviridae bacteriophage against *Staphylococcus aureus* isolated from dairy cows with mastitis. *Res Vet Sci.* 2013;95:758-63. <https://doi.org/10.1016/j.rvsc.2013.06.001>
 112. Basdew IH, Laing MD. Investigation of the lytic ability of South African bacteriophages specific for *Staphylococcus aureus*, associated with bovine mastitis. *Biocontrol Sci Technol.* 2015;25:429-43. <https://doi.org/10.1080/09583157.2014.983458>
 113. Brouillette E, Millette G, Chamberland S, Roy JP, Ster C, Kiros T, et al. Effective treatment of *Staphylococcus aureus* intramammary infection in a murine model using the bacteriophage cocktail StaphLyse™. *Viruses.* 2023;15:887. <https://doi.org/10.3390/v15040887>
 114. Tanji Y, Tanaka A, Tani K, Kurimoto M, Miyanaga K. IgG-dependent aggregation of *Staphylococcus aureus* inhibits bacteriophage attack. *Biochem Eng J.* 2015;97:17-24. <https://doi.org/10.1016/j.bej.2015.01.007>
 115. Synnott AJ, Kuang Y, Kurimoto M, Yamamichi K, Iwano H, Tanji Y. Isolation from sewage influent and characterization of novel *Staphylococcus aureus* bacteriophages with wide host ranges and potent lytic capabilities. *Appl Environ Microbiol.* 2009;75:4483-90. <https://doi.org/10.1128/AEM.02641-08>
 116. Iwano H, Inoue Y, Takasago T, Kobayashi H, Furusawa T, Taniguchi K, et al. Bacteriophage Φ SA012 has a broad host range against *Staphylococcus aureus* and effective lytic capacity in a

- mouse mastitis model. *Biology*. 2018;7:8. <https://doi.org/10.3390/biology7010008>
117. Song J, Ruan H, Chen L, Jin Y, Zheng J, Wu R, et al. Potential of bacteriophages as disinfectants to control of *Staphylococcus aureus* biofilms. *BMC Microbiol*. 2021;21:57. <https://doi.org/10.1186/s12866-021-02117-1>
 118. Srujana AS, Sonalika J, Akhila DS, Juliet MR, Sheela P. Isolation of phages and study of their in vitro efficacy on *Staphylococcus aureus* isolates originating from bovine subclinical mastitis. *Indian J Anim Res*. 2022;56:754-8. <https://doi.org/10.18805/IJAR.B-4331>
 119. Wang Z, Zheng P, Ji W, Fu Q, Wang H, Yan Y, et al. SLPW: a virulent bacteriophage targeting methicillin-resistant *Staphylococcus aureus* in vitro and in vivo. *Front Microbiol*. 2016;7:934. <https://doi.org/10.3389/fmicb.2016.00934>
 120. da Silva Duarte V, Dias RS, Kropinski AM, Campanaro S, Treu L, Siqueira C, et al. Genomic analysis and immune response in a murine mastitis model of vB_EcoM-UFV13, a potential biocontrol agent for use in dairy cows. *Sci Rep*. 2018;8:6845. <https://doi.org/10.1038/s41598-018-24896-w>
 121. Anand T, Vaid RK, Bera BC, Barua S, Riyesh T, Virmani N, et al. Isolation and characterization of a bacteriophage with broad host range, displaying potential in preventing bovine diarrhoea. *Virus Genes*. 2015;51:315-21. <https://doi.org/10.1007/s11262-015-1222-9>
 122. Kim KH, Ingale SL, Kim JS, Lee SH, Lee JH, Kwon IK, et al. Bacteriophage and probiotics both enhance the performance of growing pigs but bacteriophage are more effective. *Anim Feed Sci Technol*. 2014;196:88-95. <https://doi.org/10.1016/j.anifeedsci.2014.06.012>
 123. Lee N, Harris DH. The effect of bacteriophage treatment to reduce the rapid dissemination of *Salmonella typhimurium* in pigs. *Swine Rep 2000*. 2001;50:196-7.
 124. Gebru E, Lee JS, Son JC, Yang SY, Shin SA, Kim B, et al. Effect of probiotic-, bacteriophage-, or organic acid-supplemented feeds or fermented soybean meal on the growth performance, acute-phase response, and bacterial shedding of grower pigs challenged with *Salmonella enterica* serotype Typhimurium. *J Anim Sci*. 2010;88:3880-6. <https://doi.org/10.2527/jas.2010-2939>
 125. Skoblikow N, Zimin A. Experience of application of nontransducing bacteriophages for prophylaxy and therapy of intestinal colibacteriosis of pigs. In: Proceedings of International Conference 'Bacteriophages Theoretical and Practical Aspects of Medicine, Veterinary and Food Industry'; 2013; Ulianowsk, Russia. p. 54-7.
 126. Seo BJ, Song ET, Lee K, Kim JW, Jeong CG, Moon SH, et al. Evaluation of the broad-spectrum lytic capability of bacteriophage cocktails against various *Salmonella* serovars and their effects on weaned pigs infected with *Salmonella Typhimurium*. *J Vet Med Sci*. 2018;80:851-60. <https://doi.org/10.1292/jvms.17-0501>
 127. Mao X, Wu Y, Ma R, Li L, Wang L, Tan Y, et al. Oral phage therapy with microencapsulated phage A221 against *Escherichia coli* infections in weaned piglets. *BMC Vet Res*. 2023;19:165. <https://doi.org/10.1186/s12917-023-03724-y>
 128. Lee CY, Kim SJ, Park BC, Han JH. Effects of dietary supplementation of bacteriophages against enterotoxigenic *Escherichia coli* (ETEC) K88 on clinical symptoms of post-weaning pigs challenged with the ETEC pathogen. *J Anim Physiol Anim Nutr*. 2017;101:88-95. <https://doi.org/10.1111/jpn.12513>
 129. Carvalho CM, Gannon BW, Halfhide DE, Santos SB, Hayes CM, Roe JM, et al. The in vivo efficacy of two administration routes of a phage cocktail to reduce numbers of *Campylobacter coli* and *Campylobacter jejuni* in chickens. *BMC Microbiol*. 2010;10:232. <https://doi.org/10.1186/1471-2180-10-232>
 130. Firlieyanti AS, Connerton PL, Connerton IF. *Campylobacters* and their bacteriophages

- from chicken liver: the prospect for phage biocontrol. *Int J Food Microbiol.* 2016;237:121-7. <https://doi.org/10.1016/j.ijfoodmicro.2016.08.026>
131. Ahmadi M, Karimi Torshizi MA, Rahimi S, Dennehy JJ. Prophylactic bacteriophage administration more effective than post-infection administration in reducing *Salmonella enterica* serovar Enteritidis shedding in quail. *Front Microbiol.* 2016;7:1253. <https://doi.org/10.3389/fmicb.2016.01253>
132. Lim TH, Kim MS, Lee DH, Lee YN, Park JK, Youn HN, et al. Use of bacteriophage for biological control of *Salmonella* Enteritidis infection in chicken. *Res Vet Sci.* 2012;93:1173-8. <https://doi.org/10.1016/j.rvsc.2012.06.004>
133. Dec M, Wernicki A, Urban-Chmiel R. Efficacy of experimental phage therapies in livestock. *Anim Health Res Rev.* 2020;21:69-83. <https://doi.org/10.1017/S1466252319000161>