

Sperm hyperactivation and the CatSper channel: current understanding and future contribution of domestic animals

Jae Yeon Hwang^{1,2*}

¹Department of Molecular Biology, Pusan National University, Busan 46241, Korea

²Institute of Systems Biology, Pusan National University, Busan 46241, Korea



Received: Sep 8, 2023
Revised: Nov 10, 2023
Accepted: Nov 29, 2023

*Corresponding author

Jae Yeon Hwang
Department of Molecular Biology,
Pusan National University, Busan
46241, Korea.
Tel: +82-51-510-2289
E-mail: jyhwang@pusan.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Jae Yeon Hwang
<https://orcid.org/0000-0002-6493-4182>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. RS-2023-00210046).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the

Abstract

In female tract, mammalian sperm develop hyperactivated motility which is a key physiological event for sperm to fertilize eggs. This motility change is triggered by Ca^{2+} influx via the sperm-specific Ca^{2+} channel, CatSper. Although previous studies in human and mice largely contributed to understanding CatSper and Ca^{2+} signaling for sperm hyperactivation, the differences on their activation mechanisms are not well understood yet. There are several studies to examine expression and significance of the CatSper channel in non-human and non-mouse models, such as domestic animals. In this review, I summarize key knowledge for the CatSper channel from previous studies and propose future aspects for CatSper study using sperm from domestic animals.

Keywords: Domestic animals, Sperm, hyperactivation, CatSper, Ca^{2+}

INTRODUCTION

Sperm is the male germ cells which deliver the haploid paternal genome to the eggs to generate offspring. In mammals, sperm are ejaculated into female reproductive tract and migrate toward the fertilizing site using their cilia-like sub-cellular organelle, flagella [1,2]. Although this hair-like organelle could be simply considered to generate mechanical force for sperm navigation in female tract, complex cellular signaling pathways determine flagellar movement and sperm swimming pattern [3,4]. In female reproductive tract, sperm sense the extracellular environment and initiate various signaling pathways [5]. Eventually, the activated signaling pathways in responding to the environment changes physiological characteristics of sperm to fertilize eggs, which is the process called capacitation [6,7]. Interestingly, the capacitated sperm change their motility pattern, characterized by asymmetric flagellar beating with increased amplitude, called hyperactivated motility [8]. By developing this unique swimming strategy, mammalian sperm successfully reach the fertilizing site and penetrate the eggs' barrier, zona pellucida, followed by fertilization.

Hyperactivated motility is achieved by influx of extracellular Ca^{2+} into the sperm via sperm-specific cation channel, CatSper [9,10]. CatSper is one of the most complex ion channels composed of at least 14 subunits. Previous studies using mouse models identified four six-transmembrane (TM) subunits,

corresponding author.

Authors' contributions

The article is prepared by a single author.

Ethics approval and consent to participate

This article does not require IRB/IACUC approval because there are no human and animal participants.

CatSper1–4 [9,11,12], to form heterotetrameric pore; single TM subunits, CatSper β , γ , δ , and ϵ [13–16] to form the huge extracellular canopy structure; and cytoplasmic subunits, CatSper ζ and EFCAB9 [16,17]. In addition, recent studies to resolve atomic structure of the CatSper channel further report CatSper θ (TMEM249), CatSper η (TMEM262), and Slco6C1 as the CatSper components [18, 19]. The CatSper complexes arrange quad-linearly to form unique Ca²⁺ signaling domain on the sperm principal piece [16,17,19,20]. With the unique domain structure, the CatSper channel serves as the major Ca²⁺ entry site in sperm to develop hyperactivated motility in mammalian sperm. Thus, the CatSper deficiency or its altered function cause defective sperm hyperactivation and impaired male fertility in human and mouse [5]. However, the physiological significance of the CatSper channel in sperm Ca²⁺ signaling and hyperactivation is not well understood in other mammals including domestic animals.

CatSper subunits are even conserved in unicellular flagellates [21]. Although the subunits are lineage-specifically conserved in certain animals [16,17,22], their functional regulation and activation mechanisms seem to be variable within species. In mouse, CatSper channel is mainly activated by the intracellular alkalinization [23]. By contrast, although intracellular alkalinization can contribute to activating CatSper channel [24], an additional ligand, progesterone, is further required to fully activate the CatSper in human [25,26]. The difference between human and mouse sperm suggests the species-specific characteristics of the CatSper channel in mammalian sperm. Yet, our knowledge of the mammalian CatSper channel is only limited to human and mouse sperm, which requires more insight into the CatSper channel in other mammals. In that context, domestic animals could be good models to expand our knowledge for CatSper channel in mammals due to their commercially available semen samples.

In this review, I briefly introduce sperm hyperactivation by CatSper channel and walk through what was previously known for the CatSper channel in domestic animals with comparative aspect.

Sperm hyperactivation and CatSper Ca²⁺ for sperm hyperactivated motility

In female reproductive tract, ejaculated mammalian sperm gradually acquire the fertilizing ability via capacitation [6,7]. Interestingly, ejaculated sperm retrieved from the fertilizing site, ampulla, have vigorous flagellar beating with increased amplitude distinguished from epididymal sperm [8]. This motility pattern is also triggered by incubation with follicular fluid *in vitro* [27]. The changed motility pattern of mammalian sperm by female factors are now termed hyperactivated motility. After the first observation of the hyperactivated motility in hamster sperm, this motility pattern has been also observed from sperm from various mammals [28]. Hyperactivated motility provides increased mechanical force to sperm. Thus, by developing this unique motility pattern, sperm can successfully pass the lumen filled with mucus with escaping the oviductal reservoir [29,30]. In addition, sperm require hyperactivated motility to penetrate zona pellucida followed by fertilizing eggs after their arrival to the fertilizing site [31]. Therefore, hyperactivated motility is indispensable for successful migration and fertilization of the mammalian sperm in female tract.

Although follicular fluids can induce sperm hyperactivation *in vitro* [27], downstream signaling pathway to develop hyperactivated motility had not been well understood. Later, individual factors, such as bicarbonate, cAMP, and Ca²⁺, have been examined to test whether they are the direct physiological triggers to induce sperm hyperactivation. Finally, it was turned out that Ca²⁺ is the fundamental trigger for sperm hyperactivation in mammals. Sperm fail to develop hyperactivated motility in absence of the extracellular Ca²⁺ during *in vitro* capacitation [32,33]. In opposite, treating Ca²⁺ ionophore (A23187) transiently induces hyperactivated motility in boar and golden hamster sperm [33]. The Ca²⁺ requirement to develop hyperactivated motility was further

supported by demembrated sperm of which flagella form huge arch by the Ca^{2+} exposure [34]. Furthermore, sperm to develop hyperactivated sperm have relatively higher intracellular Ca^{2+} levels compared to the non-hyperactivated sperm [10]. All these results clearly demonstrate that sperm develop hyperactivated motility by the increased intracellular Ca^{2+} level, which are sourced from the extracellular environment. Although Ca^{2+} from intracellular storage sites, such as mitochondria, has been suggested to develop hyperactivated motility [35], its physiological triggers in female tract, which secret Ca^{2+} into the sperm cytoplasm, is unidentified yet.

CatSper, the major Ca^{2+} entry path in sperm, for hyperactivated motility

The requirement of the extracellular Ca^{2+} for sperm hyperactivation indicates directly that sperm would carry Ca^{2+} channel to introduce extracellular Ca^{2+} into their cytoplasm. Various voltage-gated Ca^{2+} channels and transient receptor potential channels had been suggested for the Ca^{2+} channel for sperm hyperactivation [36–38]. However, those channels are not considered for the major Ca^{2+} entry site because of their marginal significance in sperm hyperactivation [39]. In 2001, the sperm-specific Ca^{2+} channel, CatSper, has been identified which is now known for the major Ca^{2+} channel for sperm hyperactivation [9]. The first identified CatSper subunit, CatSper1, has six TM, like the pore subunits of the voltage-dependent K^+ channel. Previous patch-clamping analysis recorded Ca^{2+} current which is not present in *CatSper1*-null sperm [23]. These studies demonstrate that CatSper is a Ca^{2+} channel to introduce the cation into mammalian sperm.

After the first identification of CatSper1, other three six TM CatSper subunits (CatSper2, 3, and 4) were further identified, which form pore of the channel [11,40]. Accumulated studies for the last 20 years reported that CatSper channel is one of the most complex ion channel composed of over 14 subunits [41]. Contrary to the other Ca^{2+} -permeable channels, genetic alterations of the CatSper subunits result in severe defects in male reproduction (Table 1). In mouse, genetic depletion of the CatSper TM subunits results in 100% male infertility *in vivo* and *in vitro* [9,12,15,42–44]. Although their absences do not impair spermatogenesis without morphologically abnormal development of the sperm, sperm fail to develop hyperactivated motility. Thus, mouse sperm with impaired CatSper barely pass the utero-tubal junction and arrive at fertilizing site in female tract [16,20,45] and fail to penetrate zona pellucida of eggs [9]. Together with the mouse studies, various genetic mutations in genes encoding CatSper subunits have been identified from infertile male patients (Table 1). Currently, mutations in *CATSPER1* [46], *CATSPER2* [47–50] and *CATSPERE* [51] have been identified and the sperm carrying genetic mutation show motility defects. In addition, a missense mutation of *CATSPERT* was identified from an infertile male with globozoospermia whose sperm motility defect was not confirmed [52]. Thus, currently, CatSper is known for the major Ca^{2+} channel critical for sperm hyperactivation, which highlights physiological significances of the CatSper channel in male reproduction.

Molecular properties of the CatSper in human and mouse

The CatSper is one of the most complex ion channels and previous studies have demonstrated that unique molecular features of the Ca^{2+} channel is indispensable for sperm hyperactivation in mammals [41].

Notably, CatSper subunits are extremely interdependent in mouse sperm. Although it is still not well understood how the individual subunits are assembled to form functional CatSper channel in developing germ cells, genetic depletion of the CatSper TM subunits results in the absence of the entire CatSper subunits in mature sperm [15–17]. For example, *CatSper1*- and *CatSper4*-null mouse sperm do not carry any CatSper subunits although the other CatSper subunits are still expressed in testis [16,17,53]. Considering the partial interaction within the CatSper subunits in

Table 1. Identified CatSper subunits and their physiological significance in male reproduction

Subunit	Topology	First identification	Knockout mouse	Human patients
CatSper1	6 TM with VSD	EST library screening	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	Male fertility: infertile Spermatogenesis: impaired Motility: reduced
CatSper2	6 TM with VSD	Signaling peptide trapping	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	Male fertility: infertile Spermatogenesis: impaired Motility: reduced
CatSper3	6 TM with VSD	Sequence comparison	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	ND
CatSper4	6 TM with VSD	Sequence comparison	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	ND
CatSperβ	1 TM	CatSper1 interactome	ND	ND
CatSperγ	1 TM	CatSper1 interactome	ND	ND
CatSperδ	1 TM	CatSper1 interactome	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	ND
CatSperε	1 TM	Sequence comparison	ND	Male fertility: infertile Spermatogenesis: normal Motility: normal
CatSperζ	Non-TM	CatSper1 interactome	Male fertility: Subfertile Spermatogenesis: normal Hyperactivation: defective	ND
EFCAB9	Non-TM	Sperm proteome comparison	Male fertility: Subfertile Spermatogenesis: normal Hyperactivation: defective	ND
CatSperη	Non-TM	Sperm proteome comparison	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	Male fertility: infertile spermatogenesis: ND Motility: ND
CatSperη	3 TM	Atomic structure analysis	ND	ND
CatSperθ	2 TM	Atomic structure analysis	Male fertility: infertile Spermatogenesis: normal Hyperactivation: defective	ND
Slco6C1	12 TM	Atomic structure analysis	ND	ND

TM, transmembrane; VSD, voltage-sensing domain; EST, expression sequencing tag; ND, non-determined.

absence of the pore subunits [15,53], the CatSper subunits could be only present in the mature sperm tail when they are fully assembled as the functional channel. Interestingly, however, this all-in or all-out of the CatSper subunits in mouse sperm are not observed in human sperm [47]. A previous study detected CatSper subunits in sperm from infertile males who have *CATSPER2* mutation [47], which demonstrates the discrepancy for the interdependency within the CatSper subunits in human and mouse sperm.

Interestingly, individual CatSper complexes are linked to each other and arranged on the tapering sperm tail to form unique domain structure [19,20]. CatSper channel localizes specifically at the principal piece of the sperm tail with the quadrilinear arrangement resolved by super-resolution microscopy [20]. Although how the linearly arranged CatSper channels coordinate the spatiotemporal Ca^{2+} influx into the sperm, the integrity of the domain structure would be crucial for sperm hyperactivation [20,45]. All hyperactivated sperm maintain the domain structure after inducing capacitation *in vitro* [20] and sperm to arrive at fertilizing site carries intact CatSper arrangement in female tract [45]. These observations strongly support that not only the CatSper channel itself, but its continuous arrangement is required for the successful Ca^{2+} influx into sperm to develop hyperactivated motility. One possibility is that linearly arranged CatSper would enrich local Ca^{2+} concentration for the successful sperm hyperactivation like Ca^{2+} domains in neuronal and

immune pre-synapses [54], which requires further studies. This quadrilateral CatSper arrangement is also observed from human sperm [16], which indicates the overall CatSper nanodomain structure could be preserved in mammalian sperm. Recent cryo-electron microscopy (cryo-EM) studies resolved that mouse CatSper channel carries anionic transporter, Slco6C1 [18], which forms the wing-like structure at the zig-zag CatSper domain in mouse [19]. Interestingly, the wing-like structure is not clearly resolved in human sperm [19]. Although this could be simply due to the resolution issue but still there could be differences on the CatSper domain structures between human and mouse sperm, which remains to be further examined.

CatSper activation in human and mouse sperm

As mentioned, a few molecular features of the CatSper channel are different between human and mouse sperm despite their common significance to developing hyperactivated motility. Especially, human and mouse sperm requires different activation factors for CatSper channel [23,25,26].

Initially, intracellular alkalinization is known for the key trigger to activate CatSper channel in mouse sperm [23]. Kirichok et al., clearly demonstrated that inward CatSper current is significantly increased by increasing pH in intrapipette solution from patch clamping recording [23]. The CatSper activation by the intracellular alkalinization in mouse sperm is further supported by the other mouse models lacking pH regulators in sperm [5]. The absence of Na⁺/H⁺ exchangers, such as sNHE, NHA1, and NHA2, impairs sperm motility with severe fertility defects [55,56]. A previous study demonstrated that the non-TM CatSper subunits, EFCAB9 and CatSper ζ , are direct binding partner, and their binary complex serves as pH-dependent Ca²⁺ sensor to modulate CatSper channel activity [17]. Considering ancient mammals, such as monotremes and marsupials, also carry the EFCAB9 and CatSper ζ orthologues (Fig. 1) [57], the pH-dependent CatSper activation and underlying molecular mechanisms to modulate CatSper activity could be conserved in mammals. Protein kinase A signaling by the cyclic AMP (cAMP) was also expected to be another factor to activate CatSper channel by observing increased intracellular Ca²⁺ level after treating cell-permeable cAMP in mouse sperm [9]. A recent study, however, demonstrated




Fig. 1. Orthologues of the CatSper subunits are conserved in mammals. A heatmap shows preservation of the CatSper (CS) subunits, CatSper1, 2, 3, 4, β, γ, δ, ε, ζ, τ, η, θ, EFCAB9, and Slco6c1, in mammals. Except Slco6c1, orthologues for all reported subunits are conserved in mammals (dark gray). Sequence homologues for Slco6c1 is not preserved in non-eutherian mammals, Tasmanian devil and echidna (white). Sequence homologues of the mouse Slco6c1 are named to SLCO6A1 in non-rodent eutherians (light gray). Common (left) and scientific (right) names of each species are marked. Orthologue information was searched from NCBI (<https://www.ncbi.nlm.nih.gov/gene/>).

inconsistent effects of the cAMP or cAMP analogues to CatSper activation [58]. Thus, whether the cAMP can activate CatSper channel remains to be clarified in mouse sperm.

As similar to mouse sperm, previous patch clamping studies demonstrated that the CatSper channel can be also activated by intracellular alkalization in human sperm [24]. Instead of the sNHE in mouse sperm, human sperm carries voltage-dependent proton channel, Hv1, which pumps out the proton from the sperm cells [24,59]. Thus, the proton channel can maintain intracellular alkaline conditions in human sperm to potentiate the CatSper current. However, CatSper channel in human sperm could be further potentiated by steroid ligands, such as progesterone and prostaglandin E1, which is secreted from cumulus cells in female tract [25,26]. Interestingly, different from human sperm, progesterone does not elicit CatSper channel in mouse sperm [9]. In human, α/β hydrolase domain-containing protein 2 (ABHD2) has been suggested as a non-genomic progesterone receptor to activate CatSper channel [60]. ABHD2, a lipid hydrolase, governs unconventional endocannabinoid signaling in a progesterone-dependent manner by depleting 2-arachidonylglycerol, which prevents CatSper activation in human sperm [60]. This progesterone-mediated evoke of the CatSper current is also observed from sperm from macaque monkey [61], suggesting lineage-specific activation mechanisms for CatSper channel in mammal.

CatSper in domestic animals

Intensive studies in human and mouse sperm have expanded our knowledge of CatSper channel and hyperactivated motility. However, how human and mouse CatSper have different characteristics, especially activation mechanisms, are not well understood. In addition, it is also unclear whether the differences are simply due to the species-specificity of the CatSper or atypical characteristics in either human or mouse sperm. It should be further clarified by expanding our understanding of the CatSper in other species. Thus, domestic animals could be good candidates to explore non-human and non-mouse CatSper with benefits of the easy access of the semen samples. Here, I go over what have been known for the CatSper in sperm from domestic mammals.

Bull

After the first observation in Yanagimachi's study, hyperactivated motility was also reported from bull sperm [62]. In bull, sperm hyperactivation is physiologically significant for not only the successful migration toward fertilizing site, but the dissociation from the epithelial cells at oviduct [63,64]. As like the human and mouse sperm, bull sperm can develop hyperactivated motility by increased intracellular Ca^{2+} level [34,65,66].

CatSper subunits in bull sperm were identified and characterized first in 2017 [67]. In this study, CatSper1-4 were identified by genomic sequence comparison and CatSper localization at principal piece was confirmed by immunostaining in bull sperm like mouse and human sperm. In addition, the authors demonstrated that blocking CatSper function by treating antibody or Mibefradil, a CatSper inhibitor, significantly reduces sperm hyperactivation and rheotactic response, which is CatSper-dependent in mouse sperm [68]. This study demonstrates the physiological significance of the CatSper in bull sperm hyperactivation.

Interestingly, Johnson et al., also revealed that caffeine, which inhibits phosphodiesterase to increase cAMP level, can trigger sperm hyperactivation [67]. This result suggests cAMP signaling pathways could be involved in CatSper activation followed by sperm hyperactivation. One note is that *in vitro* capacitating medium for domestic animals usually carries ~mM levels of caffeine. Thus, to test pharmacological treatment with physiological concentration are still required to confirm significance of the caffeine in bull sperm hyperactivation like studies in mouse sperm [58]. Progesterone has been also suggested to activate CatSper channel in bull sperm [69]. Progesterone

significantly enhances sperm dissociation from the oviductal epithelium as well as sperm hyperactivation. This was further confirmed reciprocally by treating antagonist for progesterone receptor. Considering its effectiveness at \sim nm ranges, like in human CatSper activation, progesterone could be a factor to activate CatSper channel in bull sperm. Yet, its significance in CatSper activation is further examined directly using patch clamping and/or functional imaging assay.

Boar

Hyperactivated motility in boar sperm was first reported in 1984 [70]. Influx of extracellular Ca^{2+} triggers boar sperm hyperactivation, which releases sperm from the oviductal epithelium like bull sperm [71,72].

Porcine CatSper subunits, CatSper1–4, were molecularly cloned in 2011. *CatSper1–4* genes highly and/or exclusively express in testis and testicular expression levels increase by sexual maturity [73]. The tissue expression patterns of the CatSper subunits suggest that procedures for CatSper biogenesis in developing germ cells could be conserved in boar. Although the detailed intracellular localization is still unclear, CatSper channel also expresses in ejaculated boar sperm and pharmacological treatment revealed that CatSper inhibition using NNC 55-0396 severely damages boar sperm motility during capacitation [74].

Like bull sperm, progesterone can develop hyperactivated motility in boar sperm, which suggests that progesterone could be also an activator for CatSper channel [75]. Progesterone releases boar sperm bound to glycan-coated beads, which mimic the oviductal epithelium. In addition, to treat a CatSper inhibitor prevents its release from the glycan-coated beads regardless of the progesterone treatment. Furthermore, more directly, Ca^{2+} imaging assay demonstrated that the intracellular Ca^{2+} level increases by progesterone, which is blocked by the CatSper inhibitor. All these results elucidate that progesterone is also required for Ca^{2+} influx via CatSper channel to develop hyperactivated motility in boar sperm. Physiological significance of the cAMP in sperm hyperactivation was also highlighted in boar, but its contribution in CatSper activation requires further studies [76,77].

Horse

A previous study characterized CatSper channel in horse sperm [78]. Increasing intracellular pH by NH_4Cl enhances curvilinear swimming velocity and amplitude of lateral head displacement in horse sperm, which indicates increased sperm hyperactivation. Interestingly, increasing intracellular pH elevates intracellular Ca^{2+} level in horse sperm, which is blocked by a CatSper inhibitor, mibefradil. These results suggest that CatSper channel is activated by intracellular alkalinization in horse sperm like the channel in mouse sperm [23]. One note is that progesterone does not affect hyperactivated motility in horse sperm. In addition, another steroid hormone, prostaglandin E1, which potentiates human CatSper current, affects horse sperm hyperactivation neither [78]. All these results indicate that mechanisms for CatSper activation in horse sperm could be distinguished from those in bull and boar sperm.

Evolution of CatSper channel

Distinct evolution of CatSper orthologues in eukaryotes

Although CatSper channel has been intensively understood in mammalian sperm, previous studies reported the homologues for CatSper subunits in early eukaryotes. Comparative genomic studies identified orthologues for the CatSper proteins from early metazoans such as sea squirt [79] and sea anemone [22]. Surprisingly, the sequence homologues were discovered from even unicellular organisms, bikonta and unikonta [21]. In addition, functional study demonstrated that CatSper

can introduce Ca^{2+} into sperm from marine invertebrates, sea urchin [80] and an ascidian - *Ciona intestinalis* [81], to regulate their motility pattern. These results suggest that CatSper channel could evolve from ancient eukaryotes to modulate flagellar movement by entering extracellular Ca^{2+} . Interestingly, CatSper subunits seem to undergo distinct evolution in eukaryotes [21]. Comparative genomic studies further revealed that orthologues for all CatSper subunits are not detected from certain lineages, like insects, worms, and mollusks [16,17,22]. By contrast, CatSper components are preserved and functionally conserved as a key Ca^{2+} channel in earlier metazoans, sea urchin and *C. intestinalis* [80,81]. This discrepancy highlights lineage-specific evolution of the CatSper channel by all-in or all-out manner. How the CatSper channel is specifically preserved in certain lineages remains to be further understood.

Species-specific mechanisms for CatSper activation

Lineage-specific preservation of the CatSper channel suggests that CatSper components would be evolved rapidly under the strong selective pressure [57]. Indeed, certain CatSper components are newly appeared in specific lineage and their orthologues are highly sequence-variable in eukaryotes [16,17,57]. This variation could result in species-specific mechanisms for CatSper activation in eukaryotes.

Despite its conserved role to introduce Ca^{2+} to modulate flagellar movement, specific chemoattractants, resact and sperm-activating and -attracting factor (SAAF), are required to activate CatSper channel in marine invertebrates [80,81]. By contrast, activation of mammalian CatSper mainly relies on intracellular alkalinization and/or steroid hormone [23,25,26]. This discrepancy within mammals and marine invertebrates could be due to different molecular organization of the CatSper channel. A previous studies identified mammalian-specific CatSper components, CatSper ζ [16]. CatSper ζ forms a binary complex with another non-TM CatSper subunit, EFCAB9, and the machinery serves as pH-dependent Ca^{2+} sensor to regulate CatSper activity in mouse sperm [17]. In addition, the interaction between EFCAB9 and CatSper ζ are conserved even in ancient mammals [57]. Thus, the newly evolved CatSper ζ might determine mammalian-specific mechanisms for CatSper activation in responding to pH-dependent Ca^{2+} sensitivity.

Of note, activation mechanisms of the CatSper channel are even different within mammals. Especially, CatSper activation by progesterone and prostaglandin E1 is critical in human sperm, but not in mouse sperm [41]. Although what causes the difference between two species is still unclear, recent cryo-EM studies to resolve CatSper structure would provide a clue for the question [18,19]. Those studies demonstrated that an organic anion transporter, Slco6c1, is a CatSper component to form wing structure in mouse CatSper complex. Although the ligands and/or substrates for Slco6c1 are not defined yet, Lin et al. [18] speculate that Slco6c1 might mediate steroid hormone transportation. Interestingly, the sequence homologue of mouse Slco6c1 is SLCO6A1 in human, which is also exclusively expressed in testis. Interestingly, Slco6c1 orthologues are only annotated in a few rodents, like mouse and rat, but SLCO6A1 is broadly annotated, indicating the rodent orthologues - Slco6c1 - could have distinctive sequences. Indeed, mouse Slco6c1 is the most sequence variable among the examined sequence homologues in other mammals (Fig. 2). One note is that progesterone triggers Ca^{2+} influx in sperm from bull and boar, which have orthologues for SLCO6A1. Thus, presumably, the transporters in CatSper complex might determine steroid-mediated channel activation in a species-specific manner, which await to be studied in future.

CONCLUSION

Studies for last 20 years have expanded our knowledge in CatSper-mediated Ca^{2+} signaling and




Fig. 2. Slco6c1 is a sequence-variable CatSper subunit in mammals. (A) Distribution of the pairwise distances of Slco6c1, CatSper θ , and CatSper η between each species are represented in violin plots. (B) Pairwise distances of the Slco6c1 (left), CatSper θ (middle), and CatSper η (right) orthologues within two species are shown in heatmaps. SLCO6A1 in other species were considered for the orthologue of mouse Slco6c1. Mouse Slco6c1 is the most sequence variable among the sequence homologue examined here. Protein sequence information were obtained from NCBI (<https://www.ncbi.nlm.nih.gov/gene/>).

its physiological significance in male reproduction. Recent structural studies resolved the channel structure in atomic level, which opens the new era for the CatSper channel in translational approaches – developing male contraceptive systems and its application to assistant reproductive technology. However, our knowledge of the CatSper channel has mainly been obtained from mouse models. Despite its great contribution, there are some knowledge gaps between mouse and human CatSper, such as different activation mechanisms. This could be clearly answered by obtaining more information for CatSper channel in other mammals because of its possible species-specificity. Domestic animals could be good models to study CatSper channel because of commercially available semen samples and relatively well-established genome information. Thus, knowledge of the CatSper channel in domestic animals will provide more insights into the CatSper channel with answering remaining questions in future by reducing the current knowledge gaps between human and mouse.

REFERENCES

1. Sironen A, Shoemark A, Patel M, Loebinger MR, Mitchison HM. Sperm defects in primary ciliary dyskinesia and related causes of male infertility. *Cell Mol Life Sci.* 2020;77:2029-48. <https://doi.org/10.1007/s00018-019-03389-7>
2. Inaba K. Sperm flagella: comparative and phylogenetic perspectives of protein components. *Mol Hum Reprod.* 2011;17:524-38. <https://doi.org/10.1093/molehr/gar034>
3. Wachten D, Jikeli JF, Kaupp UB. Sperm sensory signaling. *Cold Spring Harb Perspect Biol.* 2016;9:a028225. <https://doi.org/10.1101/cshperspect.a028225>
4. Vyklicka L, Lishko PV. Dissecting the signaling pathways involved in the function of sperm flagellum. *Curr Opin Cell Biol.* 2020;63:154-61. <https://doi.org/10.1016/j.ceb.2020.01.015>
5. Wang H, McGoldrick LL, Chung JJ. Sperm ion channels and transporters in male fertility and infertility. *Nat Rev Urol.* 2021;18:46-66. <https://doi.org/10.1038/s41585-020-00390-9>
6. Chang MC. Fertilizing capacity of spermatozoa deposited into the fallopian tubes. *Nature.* 1951;168:697-8. <https://doi.org/10.1038/168697b0>
7. Austin CR. Observations on the penetration of the sperm into the mammalian egg. *Aust J Biol*

- Sci. 1951;4:581-96. <https://doi.org/10.1071/BI9510581>
8. Yanagimachi R. The movement of golden hamster spermatozoa before and after capacitation. *J Reprod Fertil.* 1970;23:193-6. <https://doi.org/10.1530/jrf.0.0230193>
 9. Ren D, Navarro B, Perez G, Jackson AC, Hsu S, Shi Q, et al. A sperm ion channel required for sperm motility and male fertility. *Nature.* 2001;413:603-9. <https://doi.org/10.1038/35098027>
 10. Suarez SS, Varosi SM, Dai X. Intracellular calcium increases with hyperactivation in intact, moving hamster sperm and oscillates with the flagellar beat cycle. *Proc Natl Acad Sci USA.* 1993;90:4660-4. <https://doi.org/10.1073/pnas.90.10.4660>
 11. Quill TA, Ren D, Clapham DE, Garbers DL. A voltage-gated ion channel expressed specifically in spermatozoa. *Proc Natl Acad Sci.* 2001;98:12527-31. <https://doi.org/10.1073/pnas.221454998>
 12. Qi H, Moran MM, Navarro B, Chong JA, Krapivinsky G, Krapivinsky L, et al. All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility. *Proc Natl Acad Sci USA.* 2007;104:1219-23. <https://doi.org/10.1073/pnas.0610286104>
 13. Wang H, Liu J, Cho KH, Ren D. A novel, single, transmembrane protein CATSPERG is associated with CATSPER1 channel protein. *Biol Reprod.* 2009;81:539-44. <https://doi.org/10.1095/biolreprod.109.077107>
 14. Liu J, Xia J, Cho KH, Clapham DE, Ren D. CatSper β , a novel transmembrane protein in the CatSper channel complex. *J Biol Chem.* 2007;282:18945-52. <https://doi.org/10.1074/jbc.M701083200>
 15. Chung JJ, Navarro B, Krapivinsky G, Krapivinsky L, Clapham DE. A novel gene required for male fertility and functional CATSPER channel formation in spermatozoa. *Nat Commun.* 2011;2:153. <https://doi.org/10.1038/ncomms1153>
 16. Chung JJ, Miki K, Kim D, Shim SH, Shi HF, Hwang JY, et al. CatSper ζ regulates the structural continuity of sperm Ca²⁺ signaling domains and is required for normal fertility. *elife.* 2017;6:e23082. <https://doi.org/10.7554/eLife.23082>
 17. Hwang JY, Mannowetz N, Zhang Y, Everley RA, Gygi SP, Bewersdorf J, et al. Dual sensing of physiologic pH and calcium by EFCAB9 regulates sperm motility. *Cell.* 2019;177:1480-94. <https://doi.org/10.1016/j.cell.2019.03.047>
 18. Lin S, Ke M, Zhang Y, Yan Z, Wu J. Structure of a mammalian sperm cation channel complex. *Nature.* 2021;595:746-50. <https://doi.org/10.1038/s41586-021-03742-6>
 19. Zhao Y, Wang H, Wieschofer C, Shah NB, Reetz E, Hwang JY, et al. 3D structure and in situ arrangements of CatSper channel in the sperm flagellum. *Nat Commun.* 2022;13:3439. <https://doi.org/10.1038/s41467-022-31050-8>
 20. Chung JJ, Shim SH, Everley RA, Gygi SP, Zhuang X, Clapham DE. Structurally distinct Ca²⁺ signaling domains of sperm flagella orchestrate tyrosine phosphorylation and motility. *Cell.* 2014;157:808-22. <https://doi.org/10.1016/j.cell.2014.02.056>
 21. Cai X, Wang X, Clapham DE. Early evolution of the eukaryotic Ca²⁺ signaling machinery: conservation of the CatSper channel complex. *Mol Biol Evol.* 2014;31:2735-40. <https://doi.org/10.1093/molbev/msu218>
 22. Cai X, Clapham DE. Evolutionary genomics reveals lineage-specific gene loss and rapid evolution of a sperm-specific ion channel complex: CatSper α and CatSper β . *PLOS ONE.* 2008;3:e3569. <https://doi.org/10.1371/journal.pone.0003569>
 23. Kirichok Y, Navarro B, Clapham DE. Whole-cell patch-clamp measurements of spermatozoa reveal an alkaline-activated Ca²⁺ channel. *Nature.* 2006;439:737-40. <https://doi.org/10.1038/nature04417>

24. Lishko PV, Botchkina IL, Fedorenko A, Kirichok Y. Acid extrusion from human spermatozoa is mediated by flagellar voltage-gated proton channel. *Cell*. 2010;140:327-37. <https://doi.org/10.1016/j.cell.2009.12.053>
25. Lishko PV, Botchkina IL, Kirichok Y. Progesterone activates the principal Ca²⁺ channel of human sperm. *Nature*. 2011;471:387-91. <https://doi.org/10.1038/nature09767>
26. Strünker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R, et al. The CatSper channel mediates progesterone-induced Ca²⁺ influx in human sperm. *Nature*. 2011;471:382-6. <https://doi.org/10.1038/nature09769>
27. Yanagimachi R. In vitro capacitation of hamster spermatozoa by follicular fluid. *J Reprod Fertil*. 1969;18:275-86. <https://doi.org/10.1530/jrf.0.0180275>
28. Kay VJ, Robertson L. Hyperactivated motility of human spermatozoa: a review of physiological function and application in assisted reproduction. *Hum Reprod Update*. 1998;4:776-86. <https://doi.org/10.1093/humupd/4.6.776>
29. Suárez SS, Osman RA. Initiation of hyperactivated flagellar bending in mouse sperm within the female reproductive tract. *Biol Reprod*. 1987;36:1191-8. <https://doi.org/10.1095/biolreprod36.5.1191>
30. Suarez SS, Pacey AA. Sperm transport in the female reproductive tract. *Hum Reprod Update*. 2006;12:23-37. <https://doi.org/10.1093/humupd/dmi047>
31. Stauss CR, Votta TJ, Suarez SS. Sperm motility hyperactivation facilitates penetration of the hamster zona pellucida. *Biol Reprod*. 1995;53:1280-5. <https://doi.org/10.1095/biolreprod53.6.1280>
32. Yanagimachi R. Requirement of extracellular calcium ions for various stages of fertilization and fertilization-related phenomena in the hamster. *Gamete Res*. 1982;5:323-44. <https://doi.org/10.1002/mrd.1120050404>
33. Fraser LR. Minimum and maximum extracellular Ca²⁺ requirements during mouse sperm capacitation and fertilization in vitro. *J Reprod Fertil*. 1987;81:77-89. <https://doi.org/10.1530/jrf.0.0810077>
34. Ho HC, Granish KA, Suarez SS. Hyperactivated motility of bull sperm is triggered at the axoneme by Ca²⁺ and not cAMP. *Dev Biol*. 2002;250:208-17. <https://doi.org/10.1006/dbio.2002.0797>
35. Marquez B, Ignatz G, Suarez SS. Contributions of extracellular and intracellular Ca²⁺ to regulation of sperm motility: release of intracellular stores can hyperactivate CatSper1 and CatSper2 null sperm. *Dev Biol*. 2007;303:214-21. <https://doi.org/10.1016/j.ydbio.2006.11.007>
36. Darszon A, Labarca P, Nishigaki T, Espinosa F. Ion channels in sperm physiology. *Physiol Rev*. 1999;79:481-510. <https://doi.org/10.1152/physrev.1999.79.2.481>
37. Wennemuth G, Westenbroek RE, Xu T, Hille B, Babcock DF. Ca_v2.2 and Ca_v2.3 (N- and R-type) Ca²⁺ channels in depolarization-evoked entry of Ca²⁺ into mouse sperm. *J Biol Chem*. 2000;275:21210-7. <https://doi.org/10.1074/jbc.M002068200>
38. Treviño CL, Felix R, Castellano LE, Gutiérrez C, Rodríguez D, Pacheco J, et al. Expression and differential cell distribution of low-threshold Ca²⁺ channels in mammalian male germ cells and sperm. *FEBS Lett*. 2004;563:87-92. [https://doi.org/10.1016/S0014-5793\(04\)00257-1](https://doi.org/10.1016/S0014-5793(04)00257-1)
39. Lishko PV, Kirichok Y, Ren D, Navarro B, Chung JJ, Clapham DE. The control of male fertility by spermatozoan ion channels. *Annu Rev Physiol*. 2012;74:453-75. <https://doi.org/10.1146/annurev-physiol-020911-153258>
40. Lobley A, Pierron V, Reynolds L, Allen L, Michalovich D. Identification of human and mouse CatSper3 and CatSper4 genes: characterisation of a common interaction domain and evidence for expression in testis. *Reprod Biol Endocrinol*. 2003;1:53. <https://doi.org/10.1186/1477->

7827-1-53

41. Hwang JY, Chung JJ. CatSper calcium channels: 20 years on. *Physiology*. 2023;38:125-40. <https://doi.org/10.1152/physiol.00028.2022>
42. Quill TA, Sugden SA, Rossi KL, Doolittle LK, Hammer RE, Garbers DL. Hyperactivated sperm motility driven by CatSper2 is required for fertilization. *Proc Natl Acad Sci USA*. 2003;100:14869-74. <https://doi.org/10.1073/pnas.2136654100>
43. Jin J, Jin N, Zheng H, Ro S, Tafolla D, Sanders KM, et al. Catsper3 and Catsper4 are essential for sperm hyperactivated motility and male fertility in the mouse. *Biol Reprod*. 2007;77:37-44. <https://doi.org/10.1095/biolreprod.107.060186>
44. Huang X, Miyata H, Wang H, Mori G, Iida-Norita R, Ikawa M, et al. A CUG-initiated CATSPER0 functions in the CatSper channel assembly and serves as a checkpoint for flagellar trafficking. *Proc Natl Acad Sci USA*. 2023;120:e2304409120. <https://doi.org/10.1073/pnas.2304409120>
45. Ded L, Hwang JY, Miki K, Shi HF, Chung JJ. 3D in situ imaging of the female reproductive tract reveals molecular signatures of fertilizing spermatozoa in mice. *elife*. 2020;9:e62043. <https://doi.org/10.7554/eLife.62043>
46. Avenarius MR, Hildebrand MS, Zhang Y, Meyer NC, Smith LL, Kahrizi K, et al. Human male infertility caused by mutations in the CATSPER1 channel protein. *Am J Hum Genet*. 2009;84:505-10. <https://doi.org/10.1016/j.ajhg.2009.03.004>
47. Schiffer C, Rieger S, Brenker C, Young S, Hamzeh H, Wachten D, et al. Rotational motion and rheotaxis of human sperm do not require functional CatSper channels and transmembrane Ca²⁺ signaling. *EMBO J*. 2020;39:e102363. <https://doi.org/10.15252/embj.2019102363>
48. Avidan N, Tamary H, Dgany O, Cattani D, Pariente A, Thulliez M, et al. CATSPER2, a human autosomal nonsyndromic male infertility gene. *Eur J Hum Genet*. 2003;11:497-502. <https://doi.org/10.1038/sj.ejhg.5200991>
49. Luo T, Chen HY, Zou QX, Wang T, Cheng YM, Wang HF, et al. A novel copy number variation in CATSPER2 causes idiopathic male infertility with normal semen parameters. *Hum Reprod*. 2019;34:414-23. <https://doi.org/10.1093/humrep/dey377>
50. Smith JF, Syritsyna O, Fellous M, Serres C, Mannowetz N, Kirichok Y, et al. Disruption of the principal, progesterone-activated sperm Ca²⁺ channel in a CatSper2-deficient infertile patient. *Proc Natl Acad Sci USA*. 2013;110:6823-8. <https://doi.org/10.1073/pnas.1216588110>
51. Brown SG, Miller MR, Lishko PV, Lester DH, Publicover SJ, Barratt CLR, et al. Homozygous in-frame deletion in CATSPERE in a man producing spermatozoa with loss of CatSper function and compromised fertilizing capacity. *Hum Reprod*. 2018;33:1812-6. <https://doi.org/10.1093/humrep/dey278>
52. Oud MS, Okutman Ö, Hendricks LAJ, de Vries PF, Houston BJ, Vissers LELM, et al. Exome sequencing reveals novel causes as well as new candidate genes for human globozoospermia. *Hum Reprod*. 2020;35:240-52. <https://doi.org/10.1093/humrep/dez246>
53. Hwang JY, Wang H, Lu Y, Ikawa M, Chung JJ. C2cd6-encoded CatSper² targets sperm calcium channel to Ca²⁺ signaling domains in the flagellar membrane. *Cell Rep*. 2022;38:110226. <https://doi.org/10.1016/j.celrep.2021.110226>
54. Nanou E, Catterall WA. Calcium channels, synaptic plasticity, and neuropsychiatric disease. *Neuron*. 2018;98:466-81. <https://doi.org/10.1016/j.neuron.2018.03.017>
55. Wang D, King SM, Quill TA, Doolittle LK, Garbers DL. A new sperm-specific Na⁺/H⁺ exchanger required for sperm motility and fertility. *Nat Cell Biol*. 2003;5:1117-22. <https://doi.org/10.1038/ncb1072>
56. Chen SR, Chen M, Deng SL, Hao XX, Wang XX, Liu YX. Sodium–hydrogen exchanger

- NHA1 and NHA2 control sperm motility and male fertility. *Cell Death Dis.* 2016;7:e2152. <https://doi.org/10.1038/cddis.2016.65>
57. Hwang JY, Maziarz J, Wagner GP, Chung JJ. Molecular evolution of CatSper in mammals and function of sperm hyperactivation in gray short-tailed opossum. *Cells.* 2021;10:1047. <https://doi.org/10.3390/cells10051047>
 58. Wang T, Young S, Krenz H, Tüttelmann F, Röpke A, Krallmann C, et al. The Ca²⁺ channel CatSper is not activated by cAMP/PKA signaling but directly affected by chemicals used to probe the action of cAMP and PKA. *J Biol Chem.* 2020;295:13181-93. <https://doi.org/10.1074/jbc.RA120.013218>
 59. Berger TK, Fußhöller DM, Goodwin N, Bönigk W, Müller A, Dokani Khesroshahi N, et al. Post-translational cleavage of Hv1 in human sperm tunes pH- and voltage-dependent gating. *J Physiol.* 2017;595:1533-46. <https://doi.org/10.1113/JP273189>
 60. Miller MR, Mannowetz N, Iavarone AT, Safavi R, Gracheva EO, Smith JF, et al. Unconventional endocannabinoid signaling governs sperm activation via the sex hormone progesterone. *Science.* 2016;352:555-9. <https://doi.org/10.1126/science.aad6887>
 61. Sumigama S, Mansell S, Miller M, Lishko PV, Cherr GN, Meyers SA, et al. Progesterone accelerates the completion of sperm capacitation and activates CatSper channel in spermatozoa from the rhesus macaque. *Biol Reprod.* 2015;93:130. <https://doi.org/10.1095/biolreprod.115.129783>
 62. Singh JP, Babcock DF, Lardy HA. Motility activation, respiratory stimulation, and alteration of Ca²⁺ transport in bovine sperm treated with amine local anesthetics and calcium transport antagonists. *Arch Biochem Biophys.* 1983;221:291-303. [https://doi.org/10.1016/0003-9861\(83\)90146-7](https://doi.org/10.1016/0003-9861(83)90146-7)
 63. Lefebvre R, Suarez SS. Effect of capacitation on bull sperm binding to homologous oviductal epithelium. *Biol Reprod.* 1996;54:575-82. <https://doi.org/10.1095/biolreprod54.3.575>
 64. Suarez SS. Formation of a reservoir of sperm in the oviduct. *Reprod Domest Anim.* 2002;37:140-3. <https://doi.org/10.1046/j.1439-0531.2002.00346.x>
 65. Ho HC, Suarez SS. Characterization of the intracellular calcium store at the base of the sperm flagellum that regulates hyperactivated motility. *Biol Reprod.* 2003;68:1590-6. <https://doi.org/10.1095/biolreprod.102.011320>
 66. Marquez B, Suarez SS. Bovine sperm hyperactivation is promoted by alkaline-stimulated Ca²⁺ influx. *Biol Reprod.* 2007;76:660-5. <https://doi.org/10.1095/biolreprod.106.055038>
 67. Johnson GP, English AM, Cronin S, Hoey DA, Meade KG, Fair S. Genomic identification, expression profiling, and functional characterization of CatSper channels in the bovine. *Biol Reprod.* 2017;97:302-12. <https://doi.org/10.1093/biolre/iox082>
 68. Miki K, Clapham DE. Rheotaxis guides mammalian sperm. *Curr Biol.* 2013;23:443-52. <https://doi.org/10.1016/j.cub.2013.02.007>
 69. Romero-Aguirregomez J, Cronin S, Donnellan E, Fair S. Progesterone induces the release of bull spermatozoa from oviductal epithelial cells. *Reprod Fertil Dev.* 2019;31:1463-72. <https://doi.org/10.1071/RD18316>
 70. Nagai T, Niwa K, Iritani A. Effect of sperm concentration during preincubation in a defined medium on fertilization in vitro of pig follicular oocytes. *J Reprod Fertil.* 1984;70:271-5. <https://doi.org/10.1530/jrf.0.0700271>
 71. Suarez SS, Dai XB, DeMott RP, Redfern K, Miranda MA. Movement characteristics of boar sperm obtained from the oviduct or hyperactivated in vitro. *J Androl.* 1992;13:75-80. <https://doi.org/10.1002/j.1939-4640.1992.tb01631.x>
 72. Holt WV, Fazeli A. Sperm storage in the female reproductive tract. *Annu Rev Anim Biosci.*

- 2016;4:291-310. <https://doi.org/10.1146/annurev-animal-021815-111350>
73. Song C, Gao B, Wu H, Xie Y, Wang X, Li B, et al. Molecular cloning, spatial and temporal expression analysis of CatSper genes in the Chinese Meishan pigs. *Reprod Biol Endocrinol*. 2011;9:132. <https://doi.org/10.1186/1477-7827-9-132>
 74. Vicente-Carrillo A, Álvarez-Rodríguez M, Rodríguez-Martínez H. The CatSper channel modulates boar sperm motility during capacitation. *Reprod Biol*. 2017;17:69-78. <https://doi.org/10.1016/j.repbio.2017.01.001>
 75. Machado SA, Sharif M, Wang H, Bovin N, Miller DJ. Release of porcine sperm from oviduct cells is stimulated by progesterone and requires CatSper. *Sci Rep*. 2019;9:19546. <https://doi.org/10.1038/s41598-019-55834-z>
 76. Harayama H. Roles of intracellular cyclic AMP signal transduction in the capacitation and subsequent hyperactivation of mouse and boar spermatozoa. *J Reprod Dev*. 2013;59:421-30. <https://doi.org/10.1262/jrd.2013-056>
 77. Otsuka N, Harayama H. Characterization of extracellular Ca²⁺-dependent full-type hyperactivation in ejaculated boar spermatozoa preincubated with a cAMP analog. *Mol Reprod Dev*. 2017;84:1203-17. <https://doi.org/10.1002/mrd.22921>
 78. Loux SC, Crawford KR, Ing NH, González-Fernández L, Macías-García B, Love CC, et al. CatSper and the relationship of hyperactivated motility to intracellular calcium and pH kinetics in equine sperm. *Biol Reprod*. 2013;89:123. <https://doi.org/10.1095/biolreprod.113.111708>
 79. Okamura Y, Nishino A, Murata Y, Nakajo K, Iwasaki H, Ohtsuka Y, et al. Comprehensive analysis of the ascidian genome reveals novel insights into the molecular evolution of ion channel genes. *Physiol Genomics*. 2005;22:269-82. <https://doi.org/10.1152/physiolgenomics.00229.2004>
 80. Seifert R, Flick M, Bönigk W, Alvarez L, Trötschel C, Poetsch A, et al. The CatSper channel controls chemosensation in sea urchin sperm. *EMBO J*. 2015;34:379-92. <https://doi.org/10.15252/embj.201489376>
 81. Kijima T, Kurokawa D, Sasakura Y, Ogasawara M, Aratake S, Yoshida K, et al. CatSper mediates not only chemotactic behavior but also the motility of ascidian sperm. *Front Cell Dev Biol*. 2023;11:1136537. <https://doi.org/10.3389/fcell.2023.1136537>