


Exploring effects of organic selenium supplementation on pork loin: Se content, meat quality, antioxidant capacity, and metabolomic profiling during storage

Hyun Young Jung¹, Hyun Jung Lee², Hag Ju Lee¹, Yoo Yong Kim^{1,3} and Cheorun Jo^{1,2,3*}

¹Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea

²Center for Food and Bioconvergence, Seoul National University, Seoul 08826, Korea

³Institute of Green Bio Science and Technology, Seoul National University, Pyeongchang 25354, Korea



Received: May 3, 2023
Revised: Jun 25, 2023
Accepted: Jun 26, 2023

*Corresponding author

Cheorun Jo
Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea.
Tel: +82-2-880-4804
E-mail: cheorun@snu.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Hyun Young Jung
<https://orcid.org/0000-0002-4561-9304>
Hyun Jung Lee
<https://orcid.org/0000-0002-6891-8008>
Hag Ju Lee
<https://orcid.org/0000-0003-2906-7666>
Yoo Yong Kim
<https://orcid.org/0000-0001-8121-3291>
Cheorun Jo
<https://orcid.org/0000-0003-2109-3798>

Abstract

This research was conducted to study the effects of organic selenium (Se) supplements at different levels on pork loin quality during storage. Fifteen pork loins were procured randomly from three groups, Con (fed basal diet), Se15 (fed 0.15 ppm organic Se along with 0.10 ppm inorganic Se), and Se45 (fed 0.45 ppm organic Se along with 0.10 ppm inorganic Se). Each sample was analyzed for Se contents, antioxidant properties (glutathione peroxidase [GPx] activity, 2,2'-azino-bis-[3-ethylbenzothiazoline-6-sulfonic acid] [ABTS] and 2,2-diphenyl-1-picrylhydrazyl [DPPH] radical scavenging activities, 2-thiobarbituric acid reactive substances), physicochemical properties (water holding capacity, pH, color), and metabolomic analysis during 14-day storage period. Se45-supplemented group showed significantly higher Se contents and GPx activity than the other groups throughout the storage period. However, other antioxidant properties were not significantly affected by Se supplementation. Selenium supplementation did not have an adverse impact on physicochemical properties. Nuclear Magnetic Resonance-based metabolomic analysis indicated that the selenium supply conditions were insufficient to induce metabolic change. These results suggest that organic Se (0.15 and 0.45 ppm) can accumulate high Se content in pork loins without compromising quality.

Keywords: Pork loin, Selenium supplementation, Meat quality, Antioxidant properties, Metabolites

INTRODUCTION

Pork feed primarily consists of soybean meal and corn, supplemented with various additives such as vitamins and minerals to control the growth rate of pigs [1,2]. The composition of pig feed can also influence pork quality [3]. Many studies have been conducted to improve both pork production and quality by supplementing pig feed with various additives, including antioxidants [4]. Vitamin C, vitamin

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This research was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through the Useful Agricultural Life Resources Industry Technology Development Project, funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (120051022SB010).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Lee Hyun Jung, Kim YY, Jo C.

Data curation: Lee Hyun Jung.

Formal analysis: Jung HY, Lee Hag Ju.

Methodology: Jung HY, Lee Hyun Jung.

Investigation: Jung HY, Lee Hag Ju.

Writing - original draft: Jung HY, Lee Hyun Jung.

Writing - review & editing: Jung HY, Lee Hyun Jung, Lee Hag Ju, Kim YY, Jo C.

Ethics approval and consent to participate

This article does not require IRB/ACUC approval because there are no human and animal participants.

E, and selenium (Se) have been used as antioxidants in feed, and previous research has shown that their use can modulate the antioxidant capacity, nutritional quality, and fatty acid composition of pork [1,5].

Se is a commonly used in pork farming due to its regulatory and immune system function [6,7]. It can also improve pork quality and nutritional value as it is an essential components of glutathione peroxidase (GPx) [8,9]. GPx is one of the antioxidant enzymes that can reduce lipid hydroperoxides and free hydrogen peroxide in body tissues [10]. Therefore, Se supplementation can increase GPx activity, potentially improving antioxidant capacity of pork [11].

Se exists in two chemical forms in nature, organic and inorganic [12]. Inorganic Se, mainly in the form of selenite and selenium salts, is commonly used in pork feed due to its easy supply and cost-effectiveness [13]. However, the use of inorganic Se has limitations such as low accumulation rate in the body despite high digestion and absorption rate [14], lower absorption rate compared to organic Se [15], and potential toxic effects at high levels [16].

On the other hand, organic Se, in the form of selenomethionine and selenium-yeast, has a higher accumulation efficiency and antioxidant activity when fed to livestock [17,18]. It can also prevent Se deficiency, which frequently occurs in weaning piglets when fed to sows [19]. In addition, organic Se has been reported to delay the post-oxidative reaction of the muscle, improving the nutritional value, flavor, and shelf life of meat, as well as meat color and water holding capacity (WHC) [20–22]. Despite being expensive, organic Se has been considered for pig feeding [23].

Recently, there has been emphasis on converting feed supplements from inorganic Se to organic Se due to the limitation of Se and the potential benefits of organic Se [24]. However, economic feasibility is an important factor in livestock industry, and the conversion rate must be considered. Several studies are currently underway to replace and/or combine inorganic Se with organic Se, and some have reported improved antioxidant performance and health levels [25]. While we have confirmed the combined effect of inorganic and organic Se on the growth performance of pigs at different levels (data not shown), their effect on antioxidant capacity and quality has not been studied for our market consumers. Therefore, we evaluated the combined effect of inorganic and organic Se on the quality of pork loin during refrigerated storage.

MATERIALS AND METHODS

Sample preparation

A total of 105 growing pigs ([Yorkshire × Landrace] × Duroc) with an average body weight of 39.85 ± 0.01 kg were divided into 15 pens with 7 pigs in a randomized complete block design. The pigs were kept in climate-controlled facility that had a fully concrete floor measuring 2.4 by 2.9 m². A feeder and a nipple drinker were provided in each pen to ensure that the pigs had unrestricted access to food and water. The experimental period was 14 weeks during with three types of experimental treatments were implemented. Each of the 5 pens was assigned to one of 3 treatment groups, resulting 5 pens per group. The experimental treatments were as follows: Con (fed basal diet), Se15 (fed 0.15 ppm organic Se along with 0.10 ppm inorganic Se), and Se45 (fed 0.45 ppm organic Se along with 0.10 ppm inorganic Se). Each treatment group was fed with 0.10 ppm of inorganic Se (Genebiotech, Gongju, Korea), while the addition of organic Se (Sel-Plex™, Alltechm, Nicholasville, KY, USA) was adjusted to induce Se accumulation in pork. The transformation from inorganic to organic Se was accomplished by partially modifying the feeding quantity of inorganic Se. From each group, 5 pigs were randomly selected and their loins (*M. longissimus*) were obtained. The samples were cut into 3 pieces (330 ± 20 g) and packaged in air permeable bags. They were then stored at 4 °C, and the following experiments were conducted on

days 0, 7 and 14. On each storage day, WHC, pH, and meat color were analyzed immediately, and the samples were frozen at -70°C until further analyses.

Se content

The Se concentration in pork loins was determined using the fluorometric method. To perform the analysis, 0.5 g of the sample was added to a screw cap culture tube containing 5 mL of a mixed solution of HClO_4 (perchloric acid 70%) and HNO_3 (nitric acid 70%) in 1:4 ratio. The culture tube was digested for 4 h in a digestion block at 210°C , then cooled down in room temperature. After cooling, add 0.5 mL HCl was added to the tube and the tube was heated at 150°C for 30 min. Then, the tube was cooled again, and 15 mL of 0.1M EDTA solution and 2 mL of 0.1% 2,3-diaminonaphthalene solution were added. The tube was vortexed for 5 sec and incubate in a water bath at 60°C for 30 min. Following incubation, a 10-second vortexing of the tube was done after adding 5 mL of cyclohexane. The extracted cyclohexane layer was transferred to a cuvette, and the absorbance was measured using 369 nm excitation and 525 nm emission settings.

Glutathione peroxidase (GPx) activity

The activity of GPx activity was measured through the utilization of Glutathione Peroxidase Assay Kit (353919, Sigma-Aldrich, Burlington, USA). Briefly, minced meat sample (5 g) was homogenized with 25 mL of cold homogenization buffer (50 mM Tris-HCl, pH 7.5, 5 mM EDTA, 1 mM DTT) at 12,000 rpm for 1 min (T25 digital ULTRA-TURRAX®, IKA, Staufen, Germany). The homogenized sample was centrifuged (Continent 512 R, Hanil, Incheon, Korea) at $10,000\times g$ for 15 min, and the supernatant was taken. The Assay Buffer, Co-Substrate Mixture, and NADPH included in the kit were mixed with the supernatant. Then, the reaction was initiated by adding hydroperoxide. Thereafter, the absorbance was measured at 340 nm every min for 10 min to confirm the GPx activity.

Antioxidant activity

Ground sample (3 g) was homogenized with 12 mL of deionized distilled water at 9,600 rpm for 30 s (T25 digital ULTRA-TURRAX®, IKA). The homogenized samples were centrifuged (Continent 512 R, Hanil) at $2,265\times g$ for 10 min, and filtered using filter paper (No. 1, Whatman PLC, Maidstone, UK). For the meat extract, after centrifuging at $2,265\times g$ for 10 min, 10 mL of chloroform was added to the filtrate.

For the 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assay, a solution of 14 mM ABTS and 4.9 mM potassium persulfate was prepared and left in the dark for 16 minutes after vigorous vortexing. The subsequent steps were performed following the protocol described by Choe et al. [26].

For the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay, 1 mL of the diluted meat extract was mixed with 1 ml of 0.2mM DPPH in methanol, vortexed, and placed in the dark for 30 min at room temperature. The subsequent steps were performed following the protocol described by Choe et al. [26].

For the 2-thiobarbituric acid reactive substances (TBARS) assay, the meat sample (5 g) was homogenized with 15 mL of deionized distilled water and 50 μL of 7.2% butylated hydroxy toluene solution at 9,600 rpm for 30 s (T25 digital ULTRA-TURRAX®, IKA). Then, the subsequent steps were followed by Rupasinghe et al. [27].

Physicochemical analysis

Minced meat sample (5 g) was placed on a filter paper and centrifuged at $252\times g$ for 10 min

(Continent 512R, Hanil). The WHC was measured as described by Kwon et al. [28] and pH by Rupasinghe et al. [27], respectively. The meat color of pork loin was measured using a colorimeter (CM-5, Konica Minolta, Osaka, Japan). Prior to measurement, the colorimeter was calibrated with a standard black plate. The meat color was measured at three different locations on the top and the bottom of each sample [22]. The color value was expressed as CIE L*, CIE a*, CIE b* and delta E was calculated as $\sqrt{(L^*)^2 + (a^*)^2 + (b^*)^2}$.

Nuclear Magnetic Resonance (NMR)-based metabolic analysis

The NMR analysis was performed according to Kim et al. [29]. In brief, each minced sample (5 g) was homogenized with 20 mL of 0.6 M perchloric acid at 12,000 rpm for 1 min (T25 digital ULTRA-TURRAX®, IKA). The homogenized samples were centrifuged at 2,265×g for 20 min (Continent 512R, Hanil), and the supernatant was transferred in another test tube and adjusted to 7.0 with sodium hydroxide. Then, the subsequent steps were performed following the method [29].

Statistical analysis

The data were analyzed using two-way analysis of variance (SAS 9.4, SAS Institute, Cary, NC, USA). The mean values and standard errors of the means were presented as the results. Differences with a significance level of 0.05 were determined by the Student-Newman-Keuls multiple range test. Partial least squares-discriminant analysis was conducted using MetaboAnalyst 5.0 (<https://www.metaboanalyst.ca>).

RESULTS AND DISCUSSION

Se content

Throughout all storage days, the pork loin supplemented with Se45 showed the highest Se contents followed by Se15 and Con (Fig. 1; $p = 0.0009$). This indicates that the higher organic Se supplementation leads to higher residual Se contents in pork loins, as organic Se sources are highly bioavailable [15,29]. When Se-yeast was supplied as organic Se source, the amount of Se in the loin increased with increasing Se concentration in the feed [30]. Zhan et al. [22] also confirmed that pig muscle Se content increased more than double when fed with organic Se compared to inorganic Se. According to the findings of Zhang et al. [31], intramuscular Se content increased significantly when SeMet was used as a feed source, in comparison to inorganic Se sources such as SeNa or basic feeding treatment groups. Furthermore, organic Se has low toxicity, high transfer efficiency, and the ability to build and maintain Se reserve in muscle [30].

Meanwhile, Se contents were slightly decreased in Se15 and Se45 on day 7 and remained constant thereafter (Fig. 1; $p < 0.0001$). This reduction in Se content in pork during the refrigerated storage is likely due to microbial activity, temperature, etc. [32]. Despite this decrease, Se15 and Se45 still had higher Se contents than Con, indicating that the effect of Se supplementation can be maintained in pork during storage. We found no further impact from the interaction between the treatment and storage period ($p = 0.6826$).

The increased Se content in pork can have various impacts, as Se may have prevented oxidative damage from live animals to meat storage [33]. Therefore, high productivity can be promoted for pigs, consumers who lack selenium can be relieved, and several beneficial effects can be provided to consumers. Se supplementation in live animals can improve reproductive physiological characteristics, such as semen volume and semen concentration [34]. Furthermore, Se supplementation in live animals can improve reproductive physiological characteristics, such as semen volume and semen concentration [34]. Furthermore, Se content in milk from sow increases,




Fig. 1. Selenium contents of pork loin raised under different selenium supplementation conditions and storage period. Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm. ^{A-C}Different letters in the same storage days indicate significant differences among selenium feeding conditions ($p < 0.05$). ^{a,b}Different letters within the same selenium feeding conditions indicate significant differences during storage ($p < 0.05$).

which has the advantage of solving Se deficiency that can easily occur in piglets [19]. With regards to meat quality, the supplementation of organic Se can enhance meat color stability by protecting myoglobin from oxidation with its antioxidant ability [22]. Calvo et al. [35] confirmed that Se-fed pork has high lipid stability during storage. In addition, consumption of Se-enriched pork may result in a reduction in toxic factors, as Se in pork has the ability to bind with heavy metals (such as cadmium, mercury, zinc, etc.) and facilitate their excretion from the body [36,37]. Moreover, Se content in pork exhibits antioxidant effects by interacting with various antioxidant enzymes in the body, which can prevent DNA damage by averting several harmful effects of free radicals [38]. Therefore, when higher organic Se is fed to pigs, pork with the higher Se content can be served to consumers, providing additional health benefits at the point of their consumption.

Antioxidant properties

Glutathione peroxidase (GPx) activity

GPx is an antioxidant enzyme that contains Se [8,9] and can be increased by Se supplementation in pigs [39,40]. As a result of confirming GPx activity in this study, organic Se supplementation had a significant effect on GPx activity (Fig. 2; $p = 0.0179$), but the effect of interaction between organic Se supplementation and storage period was not confirmed ($p = 0.7874$). Previous research has indicated that selenium can be absorbed through the digestive system and subsequently accumulated in various organs [6]. The accumulated Se undergoes various metabolic processes and plays a key role in the synthesis of GPx. As GPx contains Se in its active center, increased uptake and accumulation of Se in the body can promote its activity [41].

The increased activity of antioxidant enzymes may improve the storage stability of meat. Although the Se content in muscle decreased as the storage days increased in the experimental groups fed Se, Se45 had the highest Se content on all storage days. The increased activity of antioxidant enzymes can increase the antioxidant capacity of meat, which can have a positive effect




Fig. 2. Glutathione peroxidase (GPx) activity of pork loin raised under different selenium supplementation conditions and storage period. Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.

on improving meat quality such as storage stability.

2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and 2,2-diphenyl-1-picrylhydrazyl (DPPH) scavenging activities

To investigate antioxidant capacity of Se-supplemented pork, ABTS/DPPH scavenging activities were conducted (Table 1). Organic Se supplementation did not significantly change the ABTS

Table 1. Antioxidant properties of pork loin as raised under different selenium supplementation conditions and storage period

Item	Treatment	Storage period (days)			SEM ¹⁾
		0	7	14	
TBARS (mg MDA/kg)	Con ²⁾	0.18	0.15	0.18	0.016
	Se15	0.18 ^a	0.13 ^b	0.18 ^a	0.011
	Se45	0.16 ^{ab}	0.12 ^b	0.18 ^a	0.015
	SEM	0.021	0.009	0.010	
ABTS scavenging rate (%)	Con	32.59 ^b	39.79 ^a	39.63 ^a	1.815
	Se15	31.28 ^c	36.70 ^b	42.31 ^a	1.233
	Se45	33.11 ^b	42.46 ^a	44.79 ^a	1.495
	SEM	0.948	1.987	1.484	
DPPH scavenging rate (%)	Con	82.42 ^a	60.56 ^c	68.26 ^b	2.267
	Se15	81.06 ^a	59.89 ^c	68.46 ^b	1.473
	Se45	83.72 ^a	61.87 ^c	71.36 ^b	1.180
	SEM	1.530	1.901	1.657	

¹⁾n = 15.

²⁾Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.

^{a-c}Different letters within the same row differ significantly (p < 0.05).

TBARS, 2-thiobarbituric acid reactive substances; ABTS, 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid); DPPH, 2,2-diphenyl-1-picrylhydrazyl.

scavenging activity, the DPPH scavenging activity showed a similar trend in each treatment, possibly due to their strong correlation ($r = 0.906$). These unexpected results could be attributed to the fact that the change in GPx activity was not sufficient to affect the antioxidant activity of meat (Figs. 1 and 2). Although GPx plays a role in reducing lipid peroxide to alcohol and free hydrogen peroxide to water [10], ABTS/DPPH scavenging activities confirm the antioxidant effect through scavenging of free radicals, not hydrogen peroxide, and may not directly related to the high activity of GPx.

During 14 days of storage period, the tendencies in DPPH and ABTS scavenging activities were different (Table 1). ABTS scavenging activity was gradually increased, possibly due to the increased functional peptides from protein degradation during post-mortem ($p < 0.05$) [42]. However, in the case of DPPH assay, its activity was significantly decreased on day 7 and increased thereafter. The different results in ABTS and DPPH scavenging activities may be attributed to different mechanisms and subjects of both analytical methods. The ABTS assay is for both hydrophilic and lipophilic antioxidants, whereas DPPH assay is more applicable to hydrophobic system. It seems that post-mortem changes in pork induced stronger impact on ABTS and DPPH scavenging activities than that from organic Se supplementation.

2-thiobarbituric acid reactive substances (TBARS)

Lipid oxidation is a major concern in pork quality, as it can negatively affect acceptability of the meat. The oxidation of lipids can occur due to the inadequate scavenging capacity of antioxidants against the release of free radicals [43]. The extent of lipid oxidation during storage was assessed by conducting TBARS analysis as shown in Table 1. In the present study, organic Se supplementation did not exhibit a significant impact on lipid oxidation compared to the control group. This was unexpected as meat GPx activity can counteract free radicals, thereby influencing lipid oxidation [44]. Several factors may have contributed to this finding. Firstly, slow lipid oxidation rate by low-fat content in pork loin may have made it difficult to observe the differences from the enhanced GPx activity in the Se-supplemented groups (Fig. 2), as fat content is one of the main factors affecting lipid oxidation [43]. Additionally, the progress of lipid oxidation may have been delayed as the samples were stored at low temperatures. Consequently, we found that the lipid oxidation barely occurred in all groups after 14 days of storage, regardless of different Se feedings (Table 1). On day 7, a slight but significant decrease in TBARS value was found only in the Se-supplemented groups. Secondly, the increase in GPx may not have been enough to inhibit further lipid oxidation in pork loin. Hoac et al. [45] reported a certain decrease in lipid oxidation by GPx activity when 4 U/g GPx was added to chickens and ducks.

Taken the results from antioxidant properties together, although Se supplementation improved the activity of GPx, these changes did not affect the antioxidant activity and the lipid stability of pork loin during storage.

Physicochemical properties

Water holding capacity (WHC) and pH

During the storage period, no significant difference was observed in WHC and pH between the control and groups supplemented with organic Se (Table 2; $p = 0.5897$ and $p = 0.2557$, respectively). However, the changes in these properties varied depending on the levels of organic Se supplementation. During 14 days of storage, the WHC changed by 13.59%, 18.79%, and 18.89% in the control, Se15, and Se45 groups, respectively. It can be attributed to the decrease in water content over time (data not shown), as its decrease may limit free water release [46]. Similarly, the pH decreased at different rates in each group, with the control group having a decrease of 0.39,

Table 2. Water holding capacity (WHC) and pH of pork loin raised under different selenium supplementation conditions and storage period

Item	Treatment	Storage period (days)			SEM ¹⁾
		0	7	14	
WHC (%)	Con ²⁾	59.35 ^b	61.50 ^b	72.94 ^a	2.414
	Se15	57.80 ^b	65.27 ^b	76.59 ^a	2.681
	Se45	55.37 ^c	61.06 ^b	74.26 ^a	1.289
	SEM	2.319	2.336	1.960	
pH	Con	5.90 ^a	5.53 ^b	5.51 ^b	0.058
	Se15	5.79 ^a	5.50 ^b	5.53 ^b	0.050
	Se45	5.81 ^a	5.54 ^b	5.57 ^b	0.048
	SEM	0.067	0.047	0.038	

¹⁾n = 15.²⁾Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.^{a-c}Different letters within the same row differ significantly ($p < 0.05$).

while Se15 and Se45 had reduction of 0.26 and 0.24, respectively. Even though several studies have reported that organic Se supplementation can increase WHC and reduce the decrease in pH in pork after slaughter [33,47], in this study, organic Se supplementation (15 or 45 ppm) with 10 ppm inorganic Se did not affect WHC and pH in pork during 14 days of storage.

Meat color

In regards to meat color, there was no significant difference in the CIE L^{*}-, CIE a^{*}-, and CIE b^{*}-values among different organic Se supplementation, except for CIE a^{*}-value on day 7 (Table 3). While previous studies have reported that organic Se supplementation at 0.3 ppm can increase CIE a^{*} and CIE b^{*}-values [35], this study did not observe any changes in meat color due to the lack of pH change in pork. The pH plays an important role in the mechanism by which oxymyoglobin is oxidized to metmyoglobin. In the case of Se-yeast, a type of organic Se fed in this experiment, it was absorbed through the methionine transporter and incorporated into the protein constituting the body, suggesting that it may not have affected meat quality, including its color. Nevertheless, previous research has indicated that consumption of organic Se may enhance muscle antioxidant capacity, protecting myoglobin from oxidation and thereby improving color stability [22]. Conversely, inorganic Se has been reported to induce lighter color than pigs fed with organic Se, mainly due to water droplet loss that occurred when fed with inorganic Se [21].

During storage, different atmospheres can cause variation in the meat color of pork can [46]. The total color difference (ΔE) was calculated to confirm the changes in color (Table 3). Overall, no distinct color changes were observed in this study, indicating that the organic Se supplementation did not affect meat color in pork loin. The CIE L^{*}-value tended to decrease, possibly due to an increase in WHC (Table 2), regardless of the type of organic Se supplementation. The CIE a^{*}-value in each group was also affected by post-mortem changes. Its increases on day 7 is possibly due to the oxygenation of myoglobin and the value decreased due to oxidation to metmyoglobin [48]. No previous study has investigated the effect of mixed feeding of organic and inorganic Se on the meat color of pork. Based on the results of this study, the organic Se supplementation treatment did not affect meat color.

Nuclear Magnetic Resonance (NMR)-based metabolic analysis

We performed NMR-based metabolic analysis to investigate the effects of different Se supplementation on the metabolic profiles of pork loin during 14 days of storage. Table 4 presents

Table 3. Meat color of pork loin raised under different selenium supplementation conditions and storage period

Item	Treatment	Storage period (days)			SEM ¹⁾
		0	7	14	
CIE L*	Con ²⁾	55.56	54.47	50.93	1.353
	Se15	54.77 ^a	55.14 ^a	48.85 ^b	0.833
	Se45	54.63 ^{ab}	57.94 ^a	51.17 ^b	1.120
	SEM	0.816	1.320	1.247	
CIE a*	Con	6.70 ^b	11.26 ^{ABa}	10.41 ^a	0.636
	Se15	6.78 ^c	12.04 ^{Aa}	10.28 ^b	0.565
	Se45	6.76 ^b	10.03 ^{Ba}	9.05 ^a	0.585
	SEM	0.431	0.529	0.775	
CIE b*	Con	13.10 ^c	17.11 ^a	15.49 ^b	0.509
	Se15	13.05 ^c	17.87 ^a	14.89 ^b	0.317
	Se45	12.09 ^c	16.31 ^a	14.15 ^b	0.593
	SEM	0.285	0.676	0.415	
Chroma	Con	14.74 ^b	20.52 ^a	18.73 ^a	0.631
	Se15	14.75 ^c	21.60 ^a	18.14 ^b	0.525
	Se45	13.86 ^c	19.19 ^a	16.85 ^b	0.747
	SEM	0.371	0.792	0.683	
Hue angle	Con	62.97 ^a	56.73 ^b	56.48 ^b	1.649
	Se15	62.56 ^a	56.32 ^b	55.66 ^b	1.269
	Se45	60.87	58.57	57.67	1.290
	SEM	1.386	0.965	1.773	
ΔE	Con	-	7.50	6.69	1.437
	Se15	-	7.21	7.30	0.985
	Se45	-	6.57	5.47	1.221
	SEM	-	1.075	1.362	

¹⁾n = 15.²⁾Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.^{A,B}Different letters within the same column indicate significant differences ($p < 0.05$).^{a-c}Different letters within the same row differ significantly ($p < 0.05$).

a total of 31 metabolites that were identified across all groups, including 15 free amino acids, 4 nucleotide-related products, and 3 organic acids. To assess the metabolomic differences among treatment groups and storage periods, multivariate analysis was performed, as shown in Figs. 3 and 4, respectively. The metabolic profiles of Con, Se15, and Se45 were not distinctly different from each other on each storage day, as indicated in Fig. 3. This suggests that the accumulated Se content in Se15 and Se45 did not have an impact on the metabolic differences during the storage period. No significant changes in metabolites, except for a few such as tyrosine, inosine, and betaine on day 0 and glutamate on day 14, were observed with different Se supplementation. Furthermore, lactate content was not significantly different between Con and both Se-supplemented groups (Table 4), but its content increased during storage, leading to a pH decrease (Table 2). Although slight changes in the metabolites in each group were observed during storage period, in overall, these changes were not distinct (Fig. 4). Each group exhibited different changes in the levels of amino acids (alanine, asparagine, creatine, glutamate, glutamine, glycine, isoleucine, leucine, methionine, phenylalanine, threonine, tyrosine, and valine) and nucleotide-related compounds (hypoxanthine and inosine), as shown in Table 4. These changes can be attributed to the degradation of proteins

Table 4. Metabolites profiles (mg/100 g) of pork loin raised under different selenium supplementation conditions and storage period

Item	Treatment	Storage period (days)			SEM ¹⁾
		0	7	14	
Free amino acids					
Alanine	Con ²⁾	29.67	22.10	28.83	2.752
	Se15	24.89 ^{ab}	22.15 ^b	31.60 ^a	2.291
	Se45	28.67	26.81	33.32	2.033
	SEM	3.125	1.922	1.870	
Asparagine	Con	3.54 ^b	3.77 ^b	6.67 ^a	0.770
	Se15	2.93 ^b	4.11 ^{ab}	4.95 ^a	0.451
	Se45	3.21 ^b	4.64 ^{ab}	6.08 ^a	0.606
	SEM	0.506	0.497	0.813	
Creatine	Con	391.88 ^b	431.87 ^b	509.16 ^a	16.008
	Se15	406.25 ^c	453.50 ^b	488.20 ^a	9.882
	Se45	410.58	476.93	504.61	30.326
	SEM	10.728	30.223	15.654	
Glutamate	Con	7.08 ^b	8.34 ^b	12.49 ^{Ba}	1.179
	Se15	8.87 ^b	10.99 ^{ab}	14.211 ^{ABa}	1.284
	Se45	9.05 ^b	11.28 ^b	16.74 ^{Aa}	0.996
	SEM	1.161	1.194	1.121	
Glutamine	Con	27.97 ^a	16.53 ^b	18.89 ^b	2.882
	Se15	25.37	17.02	18.51	3.409
	Se45	26.81	22.90	19.65	2.703
	SEM	4.629	2.143	1.105	
Glycine	Con	26.71	34.43	41.64	7.313
	Se15	28.90 ^b	36.20 ^{ab}	43.24 ^a	3.604
	Se45	27.88	36.75	36.57	4.295
	SEM	4.004	5.113	6.537	
Isoleucine	Con	2.47 ^b	4.25 ^b	7.40 ^a	0.868
	Se15	2.96 ^c	5.80 ^b	8.93 ^a	0.430
	Se45	3.48 ^b	5.24 ^b	8.43 ^a	0.786
	SEM	0.269	0.728	0.977	
Leucine	Con	2.80 ^b	5.28 ^b	9.22 ^a	1.197
	Se15	4.18 ^c	7.12 ^b	11.27 ^a	0.667
	Se45	4.12 ^b	6.65 ^b	10.95 ^a	0.975
	SEM	0.568	0.983	1.241	
Methionine	Con	5.59 ^b	6.82 ^b	11.39 ^a	1.250
	Se15	5.12 ^c	8.79 ^b	11.96 ^a	0.636
	Se45	5.97 ^b	8.97 ^b	12.71 ^a	1.107
	SEM	0.345	1.18 ⁶	1.290	
Phenylalanine	Con	2.68 ^c	5.13 ^b	7.96 ^a	0.743
	Se15	3.35 ^c	6.34 ^b	9.54 ^a	0.304
	Se45	3.83 ^b	5.92 ^b	9.25 ^a	0.722
	SEM	0.192	0.653	0.838	
Taurine	Con	38.23	35.77	40.20	4.510
	Se15	36.53	40.09	43.24	3.619
	Se45	46.18	42.41	38.74	2.928
	SEM	3.399	4.239	3.534	
Threonine	Con	6.30 ^c	9.82 ^b	12.72 ^a	0.920
	Se15	7.28	13.99	13.14	1.950
	Se45	7.71 ^b	11.35 ^a	13.83 ^a	1.073
	SEM	0.429	2.217	0.839	
Tyrosine	Con	3.69 ^{Bb}	8.54 ^b	14.82 ^a	1.632
	Se15	4.45 ^{Bc}	10.18 ^b	16.50 ^a	0.708
	Se45	5.60 ^A	10.19	16.78	1.408
	SEM	0.307	1.288	1.842	
Valine	Con	4.16 ^b	6.10 ^b	9.68 ^a	1.124
	Se15	4.78 ^c	7.94 ^b	11.87 ^a	0.627
	Se45	5.70 ^d	7.58 ^b	11.60 ^a	0.992
	SEM	0.439	0.941	1.250	
β-alanine	Con	7.49	7.33	8.40 ^{AB}	0.591
	Se15	7.72	7.96	7.99 ^B	0.381
	Se45	7.84	9.05	9.55 ^A	0.498
	SEM	0.393	0.641	0.420	

Table 4. Continued

Item	Treatment	Storage period (days)			SEM ¹⁾
		0	7	14	
Nucleotide-related products					
Hypoxanthine	Con	11.43	9.47	12.92	1.168
	Se15	11.74 ^{ab}	10.15 ^b	13.40 ^a	0.709
	Se45	12.24	11.69	13.47	1.051
	SEM	1.373	0.745	0.731	
IMP	Con	79.80	92.02	76.74	5.137
	Se15	89.49	90.69	73.91	5.777
	Se45	90.51	100.20	82.46	7.372
	SEM	7.590	7.072	2.549	
Inosine	Con	37.95 ^{Bb}	54.53 ^b	75.34 ^a	6.024
	Se15	37.73 ^{Bc}	57.22 ^b	77.24 ^a	2.165
	Se45	42.24 ^{Ac}	60.48 ^b	74.93 ^a	4.201
	SEM	0.820	4.769	5.934	
UMP	Con	2.94	3.68	2.95	0.212
	Se15	3.52	3.65	3.54	0.173
	Se45	3.20	3.48	3.16	0.232
	SEM	0.215	0.217	0.188	
Organic acids					
Acetate	Con	3.41 ^b	4.73 ^b	6.55 ^a	0.434
	Se15	3.37 ^c	5.33 ^b	7.16 ^a	0.269
	Se45	3.99	5.33	5.95	0.529
	SEM	0.223	0.467	0.522	
Lactate	Con	266.39 ^b	345.02 ^a	389.90 ^a	18.649
	Se15	284.39 ^b	360.10 ^a	384.63 ^a	13.010
	Se45	277.13 ^b	362.30 ^a	371.84 ^a	24.988
	SEM	16.370	24.980	15.794	
Methylmalonate	Con	5.59 ^b	7.15 ^b	8.96 ^a	0.529
	Se15	6.15 ^b	8.04 ^a	8.79 ^a	0.297
	Se45	5.68 ^b	7.82 ^a	8.60 ^a	0.590
	SEM	0.302	0.560	0.557	
Others					
Betaine	Con	34.96 ^B	30.79	30.13	2.225
	Se15	34.86 ^B	35.28	28.78	4.596
	Se45	46.01 ^A	44.94	38.50	3.970
	SEM	2.740	3.920	4.355	
Carnosine	Con	224.98 ^b	313.86 ^a	357.50 ^a	15.515
	Se15	284.85	323.75	347.96	27.245
	Se45	221.90 ^b	315.65 ^a	337.96 ^a	28.333
	SEM	28.347	27.062	15.810	
Ethanol	Con	0.88	1.78	2.28	0.391
	Se15	1.04 ^b	2.47 ^a	2.40 ^a	0.141
	Se45	1.04 ^b	2.19 ^a	2.25 ^a	0.285
	SEM	0.107	0.253	0.423	
Glucose	Con	42.92	72.18	81.68	19.945
	Se15	46.56 ^b	74.86 ^{ab}	87.63 ^a	9.465
	Se45	67.56	89.19	77.10	25.165
	SEM	17.404	18.664	21.666	
Glycerol	Con	9.06	9.29	10.76	1.339
	Se15	9.65	9.24	12.96	1.112
	Se45	11.32	10.64	11.31	0.627
	SEM	0.829	0.892	1.393	
Methanol	Con	0.74 ^a	0.30 ^b	0.33 ^b	0.105
	Se15	0.61	0.38	0.35	0.082
	Se45	0.75	0.49	0.41	0.100
	SEM	0.136	0.091	0.033	
N,N-Dimethylglycine	Con	1.93 ^b	2.27 ^b	2.81 ^a	0.158
	Se15	1.90 ^c	2.45 ^b	2.71 ^a	0.055
	Se45	2.01 ^b	2.58 ^{ab}	2.83 ^a	0.190
	SEM	0.058	0.189	0.158	
Niacinamide	Con	4.55 ^b	6.70 ^a	7.69 ^a	0.471
	Se15	5.05 ^c	6.99 ^b	7.86 ^a	0.244
	Se45	5.16 ^b	6.91 ^a	7.60 ^a	0.537
	SEM	0.351	0.564	0.358	
O-Acetylcarnitine	Con	7.56 ^a	2.60 ^b	3.46 ^b	0.783
	Se15	7.51	3.64	3.96	1.300
	Se45	7.66 ^a	4.56 ^b	4.14 ^b	0.764
	SEM	1.541	0.658	0.280	

¹⁾n = 15.²⁾Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.^{A,B}Different letters within the same column indicate significant differences ($p < 0.05$).^{a-c}Different letters within the same row differ significantly ($p < 0.05$).




Fig. 3. Partial least squares-discriminant analysis of metabolites by storage period from pork loin raised under different selenium supplementation conditions and storage period. Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.




Fig. 4. Partial least squares-discriminant analysis of metabolites by treatment group from pork loin raised under different selenium supplementation conditions and storage period. Con, fed basal diet; Se15, pork loin from feeding organic Se 0.15 ppm + inorganic Se 0.10 ppm; Se45, pork loin from feeding organic Se 0.45 ppm + inorganic Se 0.10 ppm.

and nucleic acids during storage, leading to an increase in the content of degradation products [49]. Additionally, lactate, which was previously mentioned, the other metabolites (acetate, carnosine, ethanol, glucose, N,N-dimethylglycine, niacinamide, and O-acetylcarnitine) also showed significant changes during 14 days of storage, but not due to Se supplementation. These results suggest that the Se feeding conditions used in this experiment were not sufficient to induce metabolomic changes in pork loin.

CONCLUSION

This study found that different levels of organic Se (0.15 and 0.45 ppm) combined with inorganic Se did not significantly affect pork quality during 14 days of storage, despite an increase in

tissue Se content and GPx activity. Therefore, high Se content in the organic Se-fed group may have a positive effect on Se accumulation in pig muscle, but organic Se supplementation up to 45 ppm does not affect pork quality during storage periods of up to 14 days. In the results of supplementation with Se, the same phenomenon as the control group was confirmed on all days of storage. Therefore, through this study, it was confirmed that Se, a trace mineral used for pig breeding management, does not adversely affect pork quality.

REFERENCES

- Schwarz T, Przybyło M, Zapletal P, Turek A, Pabiańczyk M, Bartlewski PM. Effects of using corn dried distillers' grains with solubles (cDDGS) as a partial replacement for soybean meal on the outcomes of pig fattening, pork slaughter value and quality. *Animals*. 2021;11:2956. <https://doi.org/10.3390/ani11102956>
- Lebret B. Effects of feeding and rearing systems on growth, carcass composition and meat quality in pigs. *Animal*. 2008;2:1548-58. <https://doi.org/10.1017/S1751731108002796>
- Kouba M, Sellier P. A review of the factors influencing the development of intermuscular adipose tissue in the growing pig. *Meat Sci*. 2011;88:213-20. <https://doi.org/10.1016/j.meatsci.2011.01.003>
- Jiang J, Xiong YL. Natural antioxidants as food and feed additives to promote health benefits and quality of meat products: a review. *Meat Sci*. 2016;120:107-17. <https://doi.org/10.1016/j.meatsci.2016.04.005>
- Rosenvold K, Andersen HJ. Factors of significance for pork quality: a review. *Meat Sci*. 2003;64:219-37. [https://doi.org/10.1016/S0309-1740\(02\)00186-9](https://doi.org/10.1016/S0309-1740(02)00186-9)
- Dalgaard TS, Briens M, Engberg RM, Lauridsen C. The influence of selenium and selenoproteins on immune responses of poultry and pigs. *Anim Feed Sci Technol*. 2018;238:73-83. <https://doi.org/10.1016/j.anifeedsci.2018.01.020>
- Surai PF, Fisinin VI. Selenium in pig nutrition and reproduction: boars and semen quality: a review. *Asian-Australas J Anim Sci*. 2015;28:730-46. <https://doi.org/10.5713/ajas.14.0593>
- Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG, Hoekstra WG. Selenium: biochemical role as a component of glutathione peroxidase. *Science*. 1973;179:588-90. <https://doi.org/10.1126/science.179.4073.588>
- Surai PF. Selenium in nutrition and health. 1st ed. Nottingham: Nottingham University Press; 2006.
- Muthukumar K, Rajakumar S, Sarkar MN, Nachiappan V. Glutathione peroxidase3 of *Saccharomyces cerevisiae* protects phospholipids during cadmium-induced oxidative stress. *Antonie van Leeuwenhoek*. 2011;99:761-71. <https://doi.org/10.1007/s10482-011-9550-9>
- Chen J, Tian M, Guan W, Wen T, Yang F, Chen F, et al. Increasing selenium supplementation to a moderately-reduced energy and protein diet improves antioxidant status and meat quality without affecting growth performance in finishing pigs. *J Trace Elem Med Biol*. 2019;56:38-45. <https://doi.org/10.1016/j.jtemb.2019.07.004>
- Tinggi U. Selenium: its role as antioxidant in human health. *Environ Health Prev Med*. 2008;13:102-8. <https://doi.org/10.1007/s12199-007-0019-4>
- Wei C, Lin X, Zhang Y, Wan X, Wu H, He T, et al. Effects of inorganic and organic selenium sources on the growth performance of broilers in China: a meta-analysis. *Open Life Sci*. 2021;16:31-8. <https://doi.org/10.1515/biol-2021-0007>
- Kim YY. Differences in biological activity and metabolism of selenium due to its chemical form. *J Anim Sci Technol*. 2000;42:835-48.

15. Mahan DC, Parrett NA. Evaluating the efficacy of selenium-enriched yeast and sodium selenite on tissue selenium retention and serum glutathione peroxidase activity in grower and finisher swine. *J Anim Sci.* 1996;74:2967-74. <https://doi.org/10.2527/1996.74122967x>
16. Kim YY, Mahan DC. Comparative effects of high dietary levels of organic and inorganic selenium on selenium toxicity of growing-finishing pigs. *J Anim Sci.* 2001;79:942-8. <https://doi.org/10.2527/2001.794942x>
17. Lawler TL, Taylor JB, Finley JW, Caton JS. Effect of supranutritional and organically bound selenium on performance, carcass characteristics, and selenium distribution in finishing beef steers. *J Anim Sci.* 2004;82:1488-93. <https://doi.org/10.2527/2004.8251488x>
18. Mahan DC, Azain M, Crenshaw TD, Cromwell GL, Dove CR, Kim SW, et al. Supplementation of organic and inorganic selenium to diets using grains grown in various regions of the United States with differing natural Se concentrations and fed to grower-finisher swine. *J Anim Sci.* 2014;92:4991-7. <https://doi.org/10.2527/jas.2014-7735>
19. Mahan DC, Kim YY. Effect of inorganic or organic selenium at two dietary levels on reproductive performance and tissue selenium concentrations in first-parity gilts and their progeny. *J Anim Sci.* 1996;74:2711-8. <https://doi.org/10.2527/1996.74112711x>
20. Morrissey PA, Sheehy PJ, Galvin K, Kerry JP, Buckley DJ. Lipid stability in meat and meat products. *Meat Sci.* 1998;49:S73-86. [https://doi.org/10.1016/S0309-1740\(98\)90039-0](https://doi.org/10.1016/S0309-1740(98)90039-0)
21. Mahan DC, Cline TR, Richert B. Effects of dietary levels of selenium-enriched yeast and sodium selenite as selenium sources fed to growing-finishing pigs on performance, tissue selenium, serum glutathione peroxidase activity, carcass characteristics, and loin quality. *J Anim Sci.* 1999;77:2172-9. <https://doi.org/10.2527/1999.7782172x>
22. Zhan X, Wang M, Zhao R, Li W, Xu Z. Effects of different selenium source on selenium distribution, loin quality and antioxidant status in finishing pigs. *Anim Feed Sci Technol.* 2007;132:202-11. <https://doi.org/10.1016/j.anifeedsci.2006.03.020>
23. Gjerlaug-Enger E, Haug A, Gaarder M, Ljøkjel K, Stenseth RS, Sigfridson K, et al. Pig feeds rich in rapeseed products and organic selenium increased omega-3 fatty acids and selenium in pork meat and backfat. *Food Sci Nutr.* 2015;3:120-8. <https://doi.org/10.1002/fsn3.182>
24. Khalili M, Chamani M, Amanlou H, Nikkhah A, Sadeghi AA, Dehkordi FK, et al. The effect of feeding inorganic and organic selenium sources on the hematological blood parameters, reproduction and health of dairy cows in the transition period. *Acta Sci Anim Sci.* 2019;42:e45371.
25. Wang Z, Kong L, Zhu L, Hu X, Su P, Song Z. The mixed application of organic and inorganic selenium shows better effects on incubation and progeny parameters. *Poult Sci.* 2021;100:1132-41. <https://doi.org/10.1016/j.psj.2020.10.037>
26. Choe J, Park B, Lee HJ, Jo C. Potential antioxidant and angiotensin I-converting enzyme inhibitory activity in crust of dry-aged beef. *Sci Rep.* 2020;10:7883. <https://doi.org/10.1038/s41598-020-64861-0>
27. Rupasinghe RA, Alahakoon AU, Alakolanga AW, Jayasena DD, Jo C. Oxidative stability of vacuum-packed chicken wings marinated with fruit juices during frozen storage. *Food Sci Anim Resour.* 2022;42:61-72. <https://doi.org/10.5851/kosfa.2021.e62>
28. Kwon JA, Yim DG, Kim HJ, Ismail A, Kim SS, Lee HJ, et al. Effect of temperature abuse on quality and metabolites of frozen/thawed beef loins. *Food Sci Anim Resour.* 2022;42:341-9. <https://doi.org/10.5851/kosfa.2022.e9>
29. Kim HC, Yim DG, Kim JW, Lee D, Jo C. Nuclear magnetic resonance (NMR)-based quantification on flavor-active and bioactive compounds and application for distinguishment of chicken breeds. *Food Sci Anim Resour.* 2021;41:312-23. <https://doi.org/10.5851/kosfa.2020>

e102

30. Burk RF, Hill KE. Regulation of selenium metabolism and transport. *Annu Rev Nutr.* 2015;35:109-34. <https://doi.org/10.1146/annurev-nutr-071714-034250>
31. Zhang K, Zhao Q, Zhan T, Han Y, Tang C, Zhang J. Effect of different selenium sources on growth performance, tissue selenium content, meat quality, and selenoprotein gene expression in finishing pigs. *Biol Trace Elem Res.* 2020;196:463-71. <https://doi.org/10.1007/s12011-019-01949-3>
32. Sentkowska A, Pyrzyńska K. Stability of selenium compounds in aqueous extracts of dietary supplements during storage. *J Pharm Biomed Anal.* 2022;214:114714. <https://doi.org/10.1016/j.jpba.2022.114714>
33. Calvo L, Toldrá F, Aristoy MC, López-Bote CJ, Rey AI. Effect of dietary organic selenium on muscle proteolytic activity and water-holding capacity in pork. *Meat Sci.* 2016;121:1-11. <https://doi.org/10.1016/j.meatsci.2016.05.006>
34. Marin-Guzman J, Mahan DC, Chung YK, Pate JL, Pope WF. Effects of dietary selenium and vitamin E on boar performance and tissue responses, semen quality, and subsequent fertilization rates in mature gilts. *J Anim Sci.* 1997;75:2994-3003. <https://doi.org/10.2527/1997.75112994x>
35. Calvo L, Toldrá F, Rodríguez AI, López-Bote C, Rey AI. Effect of dietary selenium source (organic vs. mineral) and muscle pH on meat quality characteristics of pigs. *Food Sci Nutr.* 2017;5:94-102. <https://doi.org/10.1002/fsn3.368>
36. Feroci G, Badiello R, Fini A. Interactions between different selenium compounds and zinc, cadmium and mercury. *J Trace Elem Med Biol.* 2005;18:227-34. <https://doi.org/10.1016/j.jtemb.2004.09.005>
37. Kieliszek M, Błażej S. Current knowledge on the importance of selenium in food for living organisms: a review. *Molecules.* 2016;21:609. <https://doi.org/10.3390/molecules21050609>
38. Karag E, Németh I, Ferke A, Hajdú J, Pintér S. A vörösvértest szelén és antagonistá nyomelemek, valamint a plazma antioxidánsok koncentrációja és összefüggése érett újszülöttek köldökzsinór vérében. In: Cser MÁ, Sziklai-László I., (editors): *A szelén Szerepe a környezetben és egészségvédelemben.* Budapest: Frag Bt.; 1998. p. 112-4.
39. Jiang J, Tang X, Xue Y, Lin G, Xiong YL. Dietary linseed oil supplemented with organic selenium improved the fatty acid nutritional profile, muscular selenium deposition, water retention, and tenderness of fresh pork. *Meat Sci.* 2017;131:99-106. <https://doi.org/10.1016/j.meatsci.2017.03.014>
40. Marković R, Ćirić J, Drljačić A, Šefer D, Jovanović I, Jovanović D, et al. The effects of dietary Selenium-yeast level on glutathione peroxidase activity, tissue selenium content, growth performance, and carcass and meat quality of broilers. *Poult Sci.* 2018;97:2861-70. <https://doi.org/10.3382/ps/pey117>
41. Navarro-Alarcón M, López-Martínez MC. Essentiality of selenium in the human body: relationship with different diseases. *Sci Total Environ.* 2000;249:347-71. [https://doi.org/10.1016/S0048-9697\(99\)00526-4](https://doi.org/10.1016/S0048-9697(99)00526-4)
42. Sohaib M, Anjum FM, Sahar A, Arshad MS, Rahman UU, Imran A, et al. Antioxidant proteins and peptides to enhance the oxidative stability of meat and meat products: a comprehensive review. *Int J Food Prop.* 2017;20:2581-93. <https://doi.org/10.1080/10942912.2016.1246456>
43. Domínguez R, Pateiro M, Gagaoua M, Barba FJ, Zhang W, Lorenzo JM. A comprehensive review on lipid oxidation in meat and meat products. *Antioxidants.* 2019;8:429. <https://doi.org/10.3390/antiox8100429>
44. Surai PF, Kochish II, Fisinin VI, Juniper DT. Revisiting oxidative stress and the use of organic

- selenium in dairy cow nutrition. *Animals*. 2019;9:462. <https://doi.org/10.3390/ani9070462>
45. Hoac T, Daun C, Trafikowska U, Zackrisson J, Åkesson B. Influence of heat treatment on lipid oxidation and glutathione peroxidase activity in chicken and duck meat. *Innov Food Sci Emerg Technol*. 2006;7:88-93. <https://doi.org/10.1016/j.ifset.2005.10.001>
 46. Szymańko T, Lesiów T, Górecka J. The water-holding capacity of meat: a reference analytical method. *Food Chem*. 2021;357:129727. <https://doi.org/10.1016/j.foodchem.2021.129727>
 47. Li JG, Zhou JC, Zhao H, Lei XG, Xia XJ, Gao G, et al. Enhanced water-holding capacity of meat was associated with increased Sepw1 gene expression in pigs fed selenium-enriched yeast. *Meat Science*. 2011;87:95-100. <https://doi.org/10.1016/j.meatsci.2010.05.019>
 48. Viana ES, Gomide LAM, Vanetti MCD. Effect of modified atmospheres on microbiological, color and sensory properties of refrigerated pork. *Meat Sci*. 2005;71:696-705. <https://doi.org/10.1016/j.meatsci.2005.05.013>
 49. Tamura Y, Iwatoh S, Miyaura K, Asikin Y, Kusano M. Metabolomic profiling reveals the relationship between taste-related metabolites and roasted aroma in aged pork. *LWT-Food Sci Technol*. 2022;155:112928. <https://doi.org/10.1016/j.lwt.2021.112928>