

Complete genome sequence of candidate probiotic *Limosilactobacillus fermentum* KUFM407

Bogun Kim^{1#}, Ji yu Heo^{2#}, Xiaoyue Xu¹, Hyunju Lee¹, Duleepa Pathiraja¹, Jae-Young Kim^{2,3}, Yi Hyun Choi², In-Geol Choi^{1*}, Sae Hun Kim^{2,3*}

¹Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Korea

²Department of Food Bioscience and Technology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Korea

³Institute of Life Science and Natural Resources, Korea University, Seoul 02841, Korea



Received: Aug 21, 2023

Revised: Oct 17, 2023

Accepted: Oct 31, 2023

#These authors contributed equally to this work.

*Corresponding author

In-Geol Choi

Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Korea.

Tel: +82-2-3290-3152

E-mail: igchoi@korea.ac.kr

Sae Hun Kim

Department of Food Bioscience and Technology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Korea.

Tel: +82-2-3290-3055

E-mail: saehkim@korea.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Bogun Kim

<https://orcid.org/0000-0002-5493-1674>

Ji yu Heo

<https://orcid.org/0009-0009-5334-0125>

Abstract

It has been reported that the administration of *Limosilactobacillus fermentum* alleviates diseases such as osteoporosis and colitis. In this study, we report the complete genome sequence of *Limosilactobacillus fermentum* KUFM407, a probiotic strain of LAB isolated from Korean traditional fermented food, Kimchi. Whole genome sequencing of *L. fermentum* KUFM407 was performed on the Illumina MiSeq and Oxford Nanopore MinION platform. The genome consisted of one circular chromosome (2,077,616 base pair [bp]) with a guanine cytosine (GC) content of 51.5% and one circular plasmid sequence (13,931 bp). Genome annotation identified 1,932 protein-coding genes, 15 rRNAs, and 58 tRNAs in the assembly. The function annotation of the predicted proteins revealed genes involved in the biosynthesis of bacteriocin and fatty acids. The complete genome of *L. fermentum* KUFM407 could provide valuable information for the development of new probiotic food and health supplements.

Keywords: *Limosilactobacillus fermentum*, KUFM407, Complete genome sequence, Probiotics

Limosilactobacillus fermentum has been widely used in the fermentation of various foods and is considered a strain with high probiotic potential. Probiotics are “live microorganisms that, when administered in adequate amounts, confer a health benefit on the host” [1]. Strains of *L. fermentum* have high survival rates in the gastrointestinal tract. They strongly attach to enterocytes and produce antimicrobial compounds. In addition, *L. fermentum* has been shown to benefit host and human health by regulating immune responses and improving intestinal health.

For lactic acid bacteria (LAB) to act as functional probiotic strains, properties such as the ability to adhere to mucosal surfaces and resistance to low pH and high bile concentrations are required [2]. For acid tolerance confirmation, 0.1 mL aliquots of each active culture were inoculated in 10 mL De Man–Rogosa–Sharpe (MRS) broth (BD, Franklin Lakes, NJ, USA) broth acidified to pH 2.5 and supplemented with 1,000 U mL⁻¹ of porcine pepsin (Sigma-Aldrich, St. Louis, MO, USA). The samples were then incubated at 37°C for 3 h. To determine bile salt tolerance, 0.1 mL aliquots of each

Xiaoyue Xu
<https://orcid.org/0000-0001-9654-8676>
 Hyunju Lee
<https://orcid.org/0009-0000-5423-4812>
 Duleepa Pathiraja
<https://orcid.org/0000-0001-6239-5958>
 Jae-Young Kim
<https://orcid.org/0000-0003-1937-9535>
 Yi Hyun Choi
<https://orcid.org/0009-0006-0513-1503>
 In-Geol Choi
<https://orcid.org/0000-0001-7403-6274>
 Sae Hun Kim
<https://orcid.org/0000-0002-0990-2268>

Competing interests

No potential conflict of interest to report.

Funding sources

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through High Value-added Food Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (321034053HD020, 1545027002) and supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry, and Fisheries, funded by the Ministry of Agriculture, Food, and Rural Affairs (32136-05-1-SB010).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Choi IG, Kim SH.
 Data curation: Kim B, Heo JY, Xu X, Pathiraja D, Choi IG, Kim SH.
 Formal analysis: Kim B, Heo JY, Xu X, Pathiraja D.
 Methodology: Kim B, Heo JY, Xu X, Pathiraja D.
 Software: Kim B, Xu X, Pathiraja D.
 Validation: Kim B, Heo JY, Xu X, Lee H, Pathiraja D, Kim JY, Choi YH.
 Investigation: Heo JY, Xu X, Lee H, Pathiraja D, Kim JY, Choi YH.
 Writing - original draft: Kim B, Heo JY, Kim JY, Choi YH.
 Writing - review & editing: Kim B, Heo JY, Xu X, Lee H, Pathiraja D, Kim JY, Choi YH, Choi IG, Kim SH.

Ethics approval and consent to participate

This article does not require IRB/IACUC approval because there are no human and animal participants.

active culture were inoculated in 10 mL MRS broth containing 0.3% oxgall bile salt (Sigma-Aldrich) and incubated at 37°C for 24h. Following incubation, cell suspensions were spread on MRS agar plates, and viable cell counts were determined through plate counting methods. *L. fermentum* KUFM407 (KUFM407) showed high stability against acid and bile salts (Table 1).

KUFM407 obtained from the Food Microbiology Laboratory, Division of Food Bioscience and Technology, Korea University (Seoul, Korea) was cultivated in MRS broth for 24h at 37°C and sub-cultured three times before the extraction of genomic DNA (gDNA).

Subcultured strains were washed three times with PBS buffer, and 1 mL aliquots of the washed strains were adjusted to the OD600 range of 1.0 to 2.0. Exgene™ Cell SV (Geneall, Seoul, Korea) was used to extract gDNA after gram-positive bacteria-specific pretreatment. The presence of a single strain of KUFM407 was confirmed by gel electrophoresis and 16S rRNA sequencing.

The extracted gDNA was prepared for short-read sequencing using an Illumina® DNA Prep Kit (Illumina, San Diego, CA, USA). Short-read sequencing was performed on an Illumina MiSeq sequencer using the Illumina MiSeq® Reagent Kit v3 (Illumina), resulting in paired-end reads of 300 base pairs (bp) in length. A long-read sequencing library was prepared using an Oxford Nanopore Ligation Sequencing Kit (Oxford Nanopore, Oxford, UK). Long-read sequencing was performed on a MinION sequencing device (Oxford Nanopore) using an R9.4.1 flow cell (Oxford Nanopore). Illumina short-read sequencing yielded 1,699,990 paired-end reads (419,571,925 bp), and Oxford Nanopore long-read sequencing produced 53,365 reads totaling 298,111,808 bp.

The draft genome sequence was constructed from the long reads using Flye assembler (v. 2.9.2) [3] after two polishing iterations. Adapter sequences were removed, and short reads were quality controlled using TrimGalore (v. 0.6.7) [4] in paired-end mode. The quality of the draft genome assembly was improved by error correction with PolyPolish (v. 0.5.0) [5] using quality controlled short reads. Genome and functional annotations of predicted genes were performed using the Prokaryotic Genome Annotation Pipeline (v. 6.4) [6]. Genome completeness was assessed with BUSCO (v. 5.4.6) [7] using the Lactobacillales_odb10 dataset. Default parameters were used for all software unless otherwise noted.

The complete genome sequence of KUFM407 consisted of a circular chromosome (2,077,616 bp) with a guanine + cytosine (G+C) ratio of 51.5% and a circular plasmid sequence of 13,931 bp (Table 2). The genome was 99.7% complete. A total of 2,143 genes, including 1,932 protein-coding,

Table 1. Acid and bile tolerance of *Limosilactobacillus fermentum* KUFM407 (Log CFU/mL)¹

Variable	Initial mean counts (0 h)	Resistant to gastric juice (3 h)	Bile tolerance (24 h)
KUFM407	7.71 ± 0.10	7.45 ± 0.06*	8.52 ± 0.08*
Strain A	7.30 ± 0.08	7.24 ± 0.14	4.55 ± 0.04*
<i>L. rhamnosus</i> GG	7.02 ± 0.06	6.97 ± 0.07	8.26 ± 0.23*

¹Each value represents mean ± SD from three trials (log CFU/mL).

* $p < 0.05$ (Student's t-test, two tailed).

Table 2. Genome features of *Limosilactobacillus fermentum* KUFM407

	Length (bp)	GC (%)	Depth	CDSs	tRNA	rRNA
Chromosome	2,077,616	51.5	122.0	1,920	58	15
Plasmid	13,931	40.5	36.0	12	0	0
Total	2,091,547	51.4	121.4	1,932	58	15

bp, base pair; G, guanine; C, cytosine; tRNA, transfer RNA; rRNA, ribosomal RNA.




Fig. 1. Circular chromosome and plasmid maps of *Limosilactobacillus fermentum* KUFM407. (A) Chromosome, (B) Plasmid. Marked features are shown from the periphery to the center; protein-coding genes (forward strand), protein-coding genes (reverse strand), pseudogenes, tRNA, rRNA, GC content, and GC skew. bp: base pair; G, guanine; C, cytosine.

Table 3. Genes with biosynthetic functions in *Limosilactobacillus fermentum* KUFM407

Predicted function	Gene name	Functions	Gene position	Length (aa)
Bacteriocin production	<i>garQ</i>	Garvicin Q family class II bacteriocin	p(7,514–7,720)	68
	<i>garI</i>	Bacteriocin immunity protein	p(7,720–8,031)	103
	<i>garC</i>	Peptide cleavage/export ABC transporter	p(8,421–10,580)	719
	<i>garD</i>	Bacteriocin secretion accessory protein	p(10,59111,967)	458
Fatty acid biosynthetic gene cluster	<i>fabZ1</i>	3-hydroxyacyl-ACP dehydratase	183,265–183,711	148
	<i>marR</i>	MarR family transcriptional regulator	183,796–184,236	146
	<i>fabH</i>	Ketoacyl-ACP synthase III	184,260–185,219	319
	<i>accP</i>	Acyl carrier protein	185,248–185,496	82
	<i>fabD</i>	ACP S-malonyltransferase	185,496–186,443	315
	<i>fabG</i>	3-oxoacyl-ACP reductase FabG	186,427–187,158	243
	<i>fabF</i>	Beta-ketoacyl-ACP synthase II	187,171–188,409	412
	<i>accB</i>	Acetyl-CoA carboxylase biotin carboxyl carrier protein	188,412–188,858	148
	<i>fabZ2</i>	3-hydroxyacyl-ACP dehydratase	188,861–189,295	144
	<i>accC</i>	Acetyl-CoA carboxylase biotin carboxylase subunit	189,316–190,713	465
	<i>accD</i>	Acetyl-CoA carboxylase carboxyltransferase subunit beta	190,682–191,530	282
	<i>accA</i>	Acetyl-CoA carboxylase carboxyltransferase subunit alpha	191,523–192,296	257
	<i>fabI</i>	Enoyl-ACP reductase	192,314–193,078	254

aa, amino acid; p, plasmid.

15 rRNA, and 58 tRNA genes, and 135 pseudogenes were predicted in the genome sequence (Fig. 1). Biological functions were assigned to 1,729 (89.5%) of the protein-coding genes. The most assigned proteins were associated with replication, recombination and repair; amino acid transport and metabolism; translation, ribosomal structure and biogenesis; transcription; and carbohydrate transport and metabolism (207, 170, 155, 137, 123 genes, respectively).

In the plasmid sequence of KUFM407, four genes (*garQ*, *garI*, *garC*, *garD*) known to be involved in the production of the garvicin Q family class II bacteriocin were found [8]. Also, fatty acid biosynthetic gene cluster was identified in the chromosome and short-chain fatty acids such as

acetate, propionate, and butyrate produced by gut microbes are known to have anti-inflammatory effects [9] (Table 3). The genomic information of *L. fermentum* KUFM407 could provide insights for future research on the characteristics of this strain as a functional food and health supplement.

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The complete genome sequence has been deposited in NCBI GenBank under accession number GCA_030290995.1. The BioProject accession number is PRJNA981335 and the BioSample accession number is SAMN35673550.

REFERENCES

1. Hill C, Guarner F, Reid G, Gibson GR, Merenstein DJ, Pot B, et al. The International Scientific Association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol*. 2014;11:506-14. <https://doi.org/10.1038/nrgastro.2014.66>
2. Bao Y, Zhang Y, Zhang Y, Liu Y, Wang S, Dong X, et al. Screening of potential probiotic properties of *Lactobacillus fermentum* isolated from traditional dairy products. *Food Control*. 2010;21:695-701. <https://doi.org/10.1016/j.foodcont.2009.10.010>
3. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol*. 2019;37:540-6. <https://doi.org/10.1038/s41587-019-0072-8>
4. Krueger F, James F, Ewels P, Ayoufian E, Schuster-Boeckler B. FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo [Internet]. Zenodo. 2021 [cited 2023 Aug 7]. <https://doi.org/10.5281/zenodo.5127899>
5. Wick RR, Holt KE. Polypolish: short-read polishing of long-read bacterial genome assemblies. *PLOS Comput Biol*. 2022;18:e1009802. <https://doi.org/10.1371/journal.pcbi.1009802>
6. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res*. 2016;44:6614-24. <https://doi.org/10.1093/nar/gkw569>
7. Manni M, Berkeley MR, Seppely M, Simão FA, Zdobnov EM. BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol*. 2021;38:4647-54. <https://doi.org/10.1093/molbev/msab199>
8. Desiderato CK, Hasenauer KM, Reich SJ, Goldbeck O, Holivololona L, Ovchinnikov KV, et al. Garvicin Q: characterization of biosynthesis and mode of action. *Microb Cell Fact*. 2022;21:236. <https://doi.org/10.1186/s12934-022-01952-9>
9. Rivière A, Selak M, Lantin D, Leroy F, De Vuyst L. Bifidobacteria and butyrate-producing colon bacteria: importance and strategies for their stimulation in the human gut. *Front Microbiol*. 2016;7:979. <https://doi.org/10.3389/fmicb.2016.00979>