

Evaluation of the nutrient digestibility at each age in dogs diet by *in vitro* and *in vivo* methods

Kyeongho Jeon^{1#}, Jihwan Lee^{2#}, Minho Song^{3#}, Kihyun Kim⁴, Minseok Jo⁵, Seyeon Chang¹, Dongcheol Song¹, Sehyun Park¹, Hyuck Kim¹, Hyeun Bum Kim^{6*}, Jinho Cho^{1*}

¹Department of Animal Science, Chungbuk National University, Cheongju 28644, Korea

²Swine Science Division, National Institute of Animal Science, Rural Development Administration, Cheonan 31000, Korea

³Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Korea

⁴National Institute of Animal Science, Rural Development Administration, Wanju 55365, Korea

⁵Central Research Institute, Woosung Feed Co., Ltd, Daejeon 34379, Korea

⁶Department of Animal Biotechnology, Dankook University, Cheonan 31116, Korea



Received: Jun 2, 2024

Revised: Jul 2, 2024

Accepted: Jul 2, 2024

#These authors contributed equally to this work.

*Corresponding author

Hyeun Bum Kim

Department of Animal Biotechnology, Dankook University, Cheonan 31116, Korea.

Tel: +82-41-550-3653

E-mail: hbkim@dankook.ac.kr

Jinho Cho

Department of Animal Science, Chungbuk National University, Cheongju 28644, Korea.

Tel: +82-43-261-2544

E-mail: jinhcho@chungbuk.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Kyeongho Jeon

<https://orcid.org/0000-0003-2321-3319>

Jihwan Lee

<https://orcid.org/0000-0001-8161-4853>

Abstract

The objective of this study was to evaluate *in vitro* predictions of digestibility at each age (puppy, adult, and senior) in dogs of dry matter (DM), organic matter (OM), crude protein (CP), gross energy (GE), crude fiber (CF), and ether extract (EE) using dog diets. First, to determine the digestibility of dog diets using pepsin and pancreatin incubations, conduct the *in vitro* method. Later, 18 mixed-sex beagles were used in this experiment to compare *in vivo* digestibility. Beagles are divided into 3 groups according to their age and body weight: six puppies (under 1-year-old; 6.21 ± 0.56 kg), six adult dogs (2 to 7 years old; 8.16 ± 0.64 kg), and six senior dogs (over 8 years old; 6.95 ± 1.39 kg). Except for DM in puppies and adult dogs, in all cases, *in vitro* digestibility values were higher than *in vivo* digestibility values ($p < 0.05$). In puppies, there were strong relationships for DM and GE with r^2 values of 0.95 and 0.84, respectively, between *in vitro* and *in vivo* digestibility. Also, in adult dogs, there were strong relationships for DM and GE with r^2 values of 0.97 and 0.84, respectively, between *in vitro* and *in vivo* digestibility. However, in senior dogs, there was a lower relationship for DM, OM, CP, GE, CF, and EE with r^2 values of 0.18, 0.42, 0.01, 0.02, 0.11, and 0.04, respectively, between *in vitro* and *in vivo* digestibility. In conclusion, *in vitro*, the prediction of nutrient digestibility of DM and GE in puppies and adult dogs seems to have significant potential for practical application. However, additional research is needed to compare senior dogs with the *in vitro* method.

Keywords: *In vitro* digestibility, *In vivo* digestibility, Dog, Age

INTRODUCTION

Pets positively affect people's physical health and emotional stability [1]. These effects improve their quality of life and increase people's preference for pet ownership [2]. Pets are raised in about 66%, 69%,

Minho Song
<https://orcid.org/0000-0002-4515-5212>
Kihyun Kim
<https://orcid.org/0000-0002-9834-2126>
Minseok Jo
<https://orcid.org/0000-0001-6742-4316>
Seyeon Chang
<https://orcid.org/0000-0002-5238-2982>
Dongcheol Song
<https://orcid.org/0000-0002-5704-603X>
Sehyun Park
<https://orcid.org/0000-0002-6253-9496>
Hyuck Kim
<https://orcid.org/0000-0002-5280-0734>
Hyeun Bum Kim
<https://orcid.org/0000-0003-1366-6090>
Jinho Cho
<https://orcid.org/0000-0001-7151-0778>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was carried out with the support of 'Cooperative Research Program for Agriculture Science and Technology Development (Project No. RS-2023-00230754)' Rural Development Administration, Korea.

Acknowledgements

Not applicable.

Availability of data and material

All data generated or analyzed during this study are included in this published article.

Authors' contributions

Conceptualization: Jeon K, Lee J, Song M, Kim HB, Cho J.
Data curation: Jeon K, Park S, Kim H.
Formal analysis: Chang S, Song D.
Methodology: Song D, Park S.
Software: Jo M, Chang S, Kim H.
Validation: Song D, Park S.
Investigation: Lee J, Song M, Jo M, Kim H.
Writing - original draft: Jeon K, Lee J, Song M.
Writing - review & editing: Jeon K, Lee J, Song M, Kim K, Jo M, Chang S, Song D, Park S, Kim H, Kim HB, Cho J.

Ethics approval and consent to participate

This experiment was examined and approved (approval # 202310A-CNU-179) by the Institutional Animal Care and Use Committee of Chungnam National University, Daejeon, Korea. In experiment, dogs were collected and managed by the procedures.

and 60% of households in the United States, Australia, and the United Kingdom, respectively [3–5]. Interest in pets has increased as the majority of the population is raising them, which raises nutritional and health anxiety about their diets [6]. Because dogs are normally provided nutrients from complete and balanced diets, the nutrient content of diets and nutrient digestibility are important [7]. Pet food companies routinely perform digestibility testing to provide important information on the nutrient content of their diets [8]. Several nations have recognized the importance of the nutrient digestibility of dogs and offered related information [9–11].

In the Republic of Korea, pets have become a fundamental component of daily life, and the number of households with dogs has increased dramatically in recent years [12]. According to Joo et al. [13], dogs represent 77.4% of the total household pets. However, research on domestic dog diets is insufficient in the Republic of Korea compared to the increasing number of dogs being raised. Most domestic dog diets developed in the Republic of Korea consult overseas nutritional requirements, such as NRC [9] and AAFCO [10]. Few nutritional studies have been conducted on dog diets, so it is necessary to investigate and establish nutrient digestibility standards.

Both *in vitro* and *in vivo* methods are used to evaluate the nutrient digestibility of diets [14]. Among them, *in vitro* methods have positive features, such as being cheaper, ethical, and more time-saving, and can be utilized as an alternative to *in vivo* methods [15]. Numerous studies have used two-step *in vitro* methods to simulate digestion in the stomach and small intestines of dogs [16,17]. Most *in vitro* studies have compared feedstuff digestibility to *in vivo* studies and generated predictive equations for their relationships [18]. However, few studies based in the Republic of Korea have used dog diets to study *in vitro* digestibility and compared them with *in vivo* digestibility. Therefore, this study was conducted to evaluate *in vitro* prediction of digestibility at each age (puppy, adult, and senior) of dry matter (DM), organic matter (OM), crude protein (CP), gross energy (GE), crude fiber (CF), and ether extract (EE) using dog diets.

MATERIALS AND METHODS

Experimental diet

The experimental diet using *in vitro* and *in vivo* methods based on hydrolyzed chicken powder, soy protein, and brown rice was manufactured in extruded form. The diet was formulated to meet or exceed the nutrient requirements according to the AAFCO guideline (Table 1).

In vitro method

The *in vitro* method described by Hervera et al. [19] method was conducted in two steps with 6 replicates of dog diet.

Step 1: The samples were prepared in finely ground (< 1.0 mm) form. In stomach simulation, weigh (1.000 ± 0.001 g) of each sample in 250 mL Erlenmeyer flasks, then add 25 mL of phosphate buffer (0.1 M, pH 6.0) and 10 mL of HCl solution (0.2 M, pH 0.7) to each flask. The pH was adjusted to 2.0 using 1 M HCl and 1 M NaOH solution, and 1 mL pepsin solution (10 mg/mL; ≥ 250 units/mg solid, P7000, pepsin from porcine gastric mucosa; Sigma-Aldrich, St. Louis, MO, USA) was added to the flask to simulate stomach digestion in the dog. In addition, 1 mL of chloramphenicol solution (C0378, chloramphenicol; Sigma-Aldrich with 5 g/L ethanol) was also added to avoid bacterial fermentation. The flasks were closed with a Parafilm M® film and incubated in a shaking incubator (SWB-35, Hanyang Science Lab, Seoul, Korea) at 39 °C for 2 h.

Step 2: 5 mL of NaOH solution (0.6 M) and 10 mL of phosphate buffer (0.2 M, pH 6.8) were added to the flask after cooling at room temperature. The pH was adjusted to 6.8 using 1 M HCl and 1 M NaOH solution, and 1 mL of pancreatin solution (100 mg/mL; 4 × USP, P1750,

Table 1. Compositions of experimental dog diet

Items	Contents
Ingredient (%)	100
Hydrolyzed chicken powder	35.00
Brown rice	32.65
Tapioca starch	5.00
Soy protein	15.00
Carrot	1.00
Sweet pumpkin	2.00
Cabbage	2.00
Salt	0.40
Canola oil	3.00
Monocalcium phosphate	1.80
Calcium carbonate	1.60
Vitamin-mineral premix ¹⁾	0.50
Tocopherol	0.05
Chemical composition	
Dry matter (%)	91.09
Crude protein (%)	40.84
Ether extract (%)	6.65
Crude fiber (%)	0.27
Calcium (%)	0.78
Phosphorus (%)	0.65
Crude ash (%)	6.55
Nitrogen free extract (%)	38.81
Metabolic energy ²⁾ (kcal/kg)	3,707.00

¹⁾Vitamin and mineral premix supplied per kg of diets: 3,500 IU vitamin A; 250 IU vitamin D₃; 25 mg vitamin E; 0.052 mg vitamin K; 2.8 mg vitamin B₁ (thiamine); 2.6 mg vitamin B₂ (riboflavin); 2 mg vitamin B₆ (pyridoxine); 0.014 mg vitamin B₁₂; 6 mg Cal-d-pantothenate; 30 mg niacin; 0.4 mg folic acid; 0.036 mg biotin; 1,000 mg taurine; 44 mg FeSO₄; 3.8 mg MnSO₄; 50 mg ZnSO₄; 7.5 mg CuSO₄; 0.18 mg Na₂SeO₃; 0.9 mg Ca (IO₃)₂.

²⁾Metabolizable energy (ME) was calculated follow equation; ME (kcal/kg) = [(CP × 3.5) + [EE × 8.5] + [NFE × 3.5)] × 10.

pancreatin from the porcine pancreas; Sigma-Aldrich) was added in the flask to simulate digestion conditions in the small intestine of the dog. Then, the flasks were closed with a Parafilm M[®] film and incubated in a shaking incubator (SWB-35; Hanyang Science Lab) at 39 °C for 4 h.

Then, the collected undigested samples were filtered through pre-dried and pre-weighed glass filter crucibles (Gooch Type Filter Crucibles, PYREX[®], Sunderland, UK). During filtering, the flasks were rinsed three times with distilled water. Additionally, 10 mL of 95% ethanol and 10 mL of 99.5% acetone were added twice to the glass filter crucibles.

Chemical analyses and calculation

At the end of the *in vitro* procedure, the filter crucibles containing undigested residues were dried at 70 °C for 24 h to calculate DM. Then, they were burned at 550 °C for 4 h to calculate OM. After being dried and combusted, it was cooled to room temperature and then weighed. The methods utilized for the determination of DM (method 930.15), OM (method 942.05), CF (method 978.10) and EE (method 920.39) were conducted with the methods of AOAC [20]. The CP and GE content were analyzed by using the dumas (Rapid MAX N-Exceed, Elementar, Langensfeld, Germany) and bomb calorimeter (Parr 6400 Bomb Calorimeter, Parr Instrument, Moline, IL,

USA), respectively.

Calculating the *in vitro* digestibility of DM using the following formula:

$$\text{Digestibility (\%)} = 100 - \{(\text{residue weight/sample weight}) \times 100\}$$

Calculating the *in vitro* digestibility of OM, CP, GE, CF and EE used the following formula:

$$\text{Digestibility (\%)} = 100 - \{Nr \times (100 - \text{IDDM})/Nd\}$$

Nr = nutrient concentration in residues (DM %), Nd = nutrient concentration in diet (DM %), and IDDM = *in vitro* digestibility (DM %)

In vivo method

Animal ethics

This experiment was examined and approved (approval # 202310A-CNU-179) by the Institutional Animal Care and Use Committee of Chungnam National University, Daejeon, Korea. In experiment, dogs were collected and managed by the procedures.

Animals and experiment design

A total of 18 mixed-sex beagles were used in this experiment. Beagles were divided into 3 groups according to their age: six puppies (under 1 year old), six adult dogs (2 to 7 years old), and six senior dogs (over 8 years old). Total experimental period was 17 days which included 7 days adaptation period. Each dog was managed in individual cage (0.9 m × 0.9 m × 0.9 m), and the temperature was maintained at 23°C. The maintenance energy requirements (MER) for each growth stage were calculated using metabolic body weight (mBW).

Calculating the MER used the following formula:

$$\begin{aligned} \text{Puppies} &= 132 \times \text{mBW} (\text{BW}^{0.75}) \times 1.5; \text{Adult dogs} = 132 \times \text{mBW} (\text{BW}^{0.75}); \\ \text{Senior dogs} &= 105 \times \text{mBW} (\text{BW}^{0.75}) \end{aligned}$$

Daily feed requirements were calculated in accordance with MER applied to each dog and fed twice a day at 9:00 and 17:00.

Nutrient digestibility

At the bottom of each kennel, dense mesh was attached to separate urine and feces for collecting pure fecal samples. Pee pads absorbed urine through the mesh, and the fecal samples remained on the mesh. Fecal samples for calculating digestibility by the total fecal collection method were collected during 8 days of experimental periods. Fresh fecal and feed samples were stored in a freezer at -20°C after collection immediately. The stored fecal samples were dried at 103°C for 12 h and then finely ground (< 1 mm) for chemical analysis at the end of the experiment. The total fecal collection digestibility of DM, OM, CP, GE, CF and EE were analyzed using samples. The methods utilized for the determination of DM (method 930.15), OM (method 942.05), and EE (method 920.39) were conducted with the methods of AOAC [20]. The CP and GE content were analyzed by using the dumas (Rapid MAX N-Exceed, Elementar, Langensfeld, Germany) and bomb calorimeter (Parr 6400 Bomb Calorimeter, Parr Instrument), respectively. The equation for the total fecal collection method described by Donadelli and Aldrich [21].

Total fecal collection digestibility was determined by the following formula:

$$\text{Digestibility (\%)} = \left[\frac{\{\% \text{Nutrient in Diet} \times \text{Feed Intake (g)}\} - \{\% \text{Nutrient in Fecal} \times \text{Fecal Output (g)}\}}{\{\% \text{Nutrient in Diet} \times \text{Feed Intake}\}} \right]$$

Statistical analysis

Dog means served as the experimental unit. The means of the treatments were also compared by using orthogonal contrasts: *in vitro* digestibility vs. other treatments. Variability in the data was expressed as the SEM. The relationship between *in vitro* and *in vivo* digestibility measured in dogs was determined by regression analyses using a general linear model (GLM) in a JMP (JMP® Pro version 16.0.0, SAS Institute, Cary, NC, USA). The model was $y = ax + b$, where $y =$ *in vivo* digestibility, $a =$ slope, $x =$ *in vitro* digestibility and $b =$ intercept. Statistical differences were determined to be significant at $p < 0.05$.

RESULTS

In vitro and *in vivo* digestibility

The *in vitro* and *in vivo* digestibility of DM, OM, CP, GE, CF and EE of puppies, adult dogs, and senior dogs are presented in Table 2. The *in vivo* digestibility of DM in senior dogs was significantly higher ($p = 0.027$) than *in vitro* digestibility. Also, the *in vivo* digestibility of CP, GE, CF, and EE in all ages was significantly higher ($p < 0.001$) than *in vitro* digestibility. However, there was no significant difference in the *in vitro* digestibility compared to the *in vivo* digestibility of DM in adults and senior groups and OM in all age groups, respectively.

The relationships between *in vitro* and *in vivo* digestibility

The statistical relationships between *in vitro* and *in vivo* digestibility as linear regression equations are shown in Table 3. There was a strong relationship between DM and GE ($r^2 = 0.95$ and 0.84 , respectively) in puppies. In adult dogs, there was a strong relationship between DM and GE ($r^2 = 0.97$ and 0.84 , respectively). However, in senior dogs, there was a low relationship between whole contents (DM, $r^2 = 0.18$; OM, $r^2 = 0.42$; CP, $r^2 = 0.01$; GE, $r^2 = 0.02$; CF, $r^2 = 0.11$; EE, $r^2 = 0.04$).

DISCUSSION

This study evaluated the digestibility of a dog diet using *in vivo* and *in vitro* methods and generated predictive equations for the relationships between *in vivo* and *in vitro* digestibility. Previous studies reported that *in vitro* digestibility was higher than *in vivo* digestibility due to endogenous losses

Table 2. Comparison of *in vitro* and *in vivo* digestibility using developed dog diet¹⁾

Items (%)	IVT	IVVP	IVVA	IVVS	SE	Contrasts (p-value)		
						IVT vs IVVP	IVT vs IVVA	IVT vs IVVS
DM	95.87	95.92	96.14	95.30	0.17	0.809	0.266	0.027
OM	92.05	92.06	92.88	92.32	0.48	0.983	0.241	0.695
CP	96.10	92.25	92.01	89.65	0.54	< 0.001	< 0.001	< 0.001
GE	95.22	92.99	93.82	92.63	0.28	< 0.001	< 0.001	< 0.001
CF	94.59	79.47	84.11	83.40	0.73	< 0.001	< 0.001	< 0.001
EE	93.60	82.86	86.23	85.63	0.49	< 0.001	< 0.001	< 0.001

¹⁾Each mean represents 6 observations for *in vivo* and *in vitro*, respectively.

IVT, *in vitro* digestibility; IVVP, *in vivo* digestibility of puppies; IVVA, *in vivo* digestibility of adult dogs; IVVS, *in vivo* digestibility of senior dogs; DM, dry matter; OM, organic matter; CP, crude protein; GE, gross energy; CF, crude fiber; EE, ether extract.

Table 3. Linear regression analysis between *in vivo* (y) and *in vitro* digestibility (x) in dog diets¹⁾

Items	Equation	r ²	RMSE
Puppies			
DM	y = 0.85x+14.11	0.95	0.08
OM	y = -0.19x +109.83	0.43	0.50
CP	y = 0.12x+80.52	0.01	1.03
GE	y = 0.66x+30.47	0.84	0.12
CF	y = 1.48x-60.63	0.20	2.12
EE	y = -0.08x+90.74	0.01	1.43
Adult dogs			
DM	y = 1.17x-16.13	0.97	0.08
OM	y = 0.11x+82.85	0.25	0.43
CP	y = 0.05x+87.14	0.00	1.34
GE	y = 1.07x-7.66	0.84	0.19
CF	y = 1.39x-47.51	0.29	1.57
EE	y = -0.06x+91.54	0.02	0.64
Senior dogs			
DM	y = 0.65x+32.55	0.18	0.54
OM	y = -0.31x+120.63	0.42	0.82
CP	y = 0.30x+61.13	0.01	2.24
GE	y = 0.40x+54.56	0.02	1.34
CF	y = 1.20x-29.68	0.11	2.39
EE	y = 0.18x+69.19	0.04	1.40

¹⁾Each mean represents 6 observations for *in vivo* and *in vitro*, respectively.

RMSE, root mean squared error; DM, dry matter; OM, organic matter; CP, crude protein; GE, gross energy; CF, crude fiber; EE, ether extract; IVT, *in vitro* digestibility; IVVP, *in vivo* digestibility of puppies; IVVA, *in vivo* digestibility of adult dogs; IVVS, *in vivo* digestibility of senior dogs.

in the body [18,21,22]. In this study, the *in vitro* digestibility of CP, GE, CF, and EE was higher than the *in vivo* digestibility at all ages. Consistent with our results, Penazzi et al. [23] suggested that *in vitro* digestibility overestimated *in vivo* digestibility. Endogenous losses in the body have a significant influence on *in vivo* digestibility [18]. In the *in vitro* method, chloramphenicol was added to avoid bacterial fermentation, and the method was conducted under strictly controlled temperature, digestion time, pH, and enzyme content conditions [24], which explains why *in vitro* digestibility was higher than *in vivo* digestibility. Le Bon et al. [25] reported that senior dogs had less inflammation and attributed it to gut microbial diversity decreases in aging dogs. Decreases in gut microbial diversity affect gut health, leading to low digestibility [26]. In this study, a significant difference between DM *in vivo* and *in vitro* digestibility was seen due to the low digestibility of senior dogs.

The *in vitro* method can assist in identifying nutritional availability in non-ruminant animals [27]. Prior studies were conducted on the *in vitro* digestibility of dog diets compared to *in vivo* digestibility [17,28,29]. This study adopted a modified two-step *in vitro* procedure for dogs, which involved reducing the doses of exogenous digestive enzymes to account for the shorter gastrointestinal tract and faster digestion rate in dogs compared to pigs [17].

The wide range of nutrient contents in dog diets may affect the accuracy of *in vitro* equations for predicting nutrient availability [26]. Endogenous losses, enzymatic secretion, and microbial activity were reported to be other influencing factors [30]. In this study, a predictive equation was

generated by comparing *in vivo* and *in vitro* digestibility in each age group. A strong relationship between DM and GE was found in puppy and adult-aged dogs. Satterlee et al. [31] reported that the analysis of animal protein-based diets resulted in lower accuracy, leading to differences in the digestibility relationship. Burrows et al. [32] suggested that the presence of dietary fiber also affects the digestibility of diets. Consistent with previous studies, Biagi et al. [29] assumed that the low relationship between *in vitro* and *in vivo* digestibility could be attributed to the fact that feces include bacteria and other endogenous protein sources, as well as to proteins derived from diets, which causes protein digestibility to be underestimated. In this study, the low relationship between the *in vitro* and *in vivo* digestibility of CP, CF, and EE was assumed to be caused by endogenous losses. In senior dogs, a low relationship between *in vitro* and *in vivo* digestibility was found for all dietary components analyzed. The low level of adjustment may have been affected by the limited number of samples and the consistent *in vivo* values recorded across samples [33]. Weber et al. [34] reported that growth affected digestibility by altering the transit time of the digestive system. Consistent with previous studies, our findings were likely due to differences in *in vivo* digestibility due to age differences, resulting in a low correlation with *in vitro* digestibility values.

Based on these results, we can use equations to predict age-specific digestibility through *in vitro* experiments. However, additional research is needed to investigate the relationship between *in vitro* and *in vivo* methods in senior dogs.

CONCLUSION

There were strong linear relationships between *in vivo* and *in vitro* digestibility (DM and GE) in puppies, (DM and GE) in adult dogs. *In vitro*, prediction of digestibility (DM and GE) in puppies and adult dogs seem to have significant potential for practical application. However, additional research investigating the *in vitro* method in senior dogs is needed.

REFERENCES

1. Jadhav S, Gaonkar T, Joshi M, Rathi A. Modulation of digestibility of canine food using enzyme supplement: an *in vitro* simulated semi-dynamic digestion study. *Front Vet Sci*. 2023;10:1220198. <https://doi.org/10.3389/fvets.2023.1220198>
2. Martins CF, Silva L, Soares J, Pinto GS, Abrantes C, Cardoso L, et al. Walk or be walked by the dog? The attachment role. *BMC Public Health*. 2024;24:684. <https://doi.org/10.1186/s12889-024-18037-4>
3. APPA [American Pet Products Association]. 2024 APPA National Pet Owners Survey Stats [Internet]. APPA. 2024 [cited 2024 May 15]. <https://americanpetproducts.org/industry-trends-and-stats>
4. AMA [Animal Medicines Australia]. Pet in Australia [Internet]. AMA. 2022 [cited 2024 May 15]. https://animalmedicinesaustralia.org.au/wp-content/uploads/2022/11/AMAU008-Pet-Ownership22-Report_v1.6_WEB.pdf
5. PFMA [Pet food manufacturing association]. Annual report [Internet]. PFMA. 2024 [cited 2024 May 15]. <https://www.pfma.org.uk/annual-reports>
6. Bos E, Hendriks W, Beerda B, Bosch G. Determining the protocol requirements of in-home dog food digestibility testing. *Br J Nutr*. 2023;130:164-73. <https://doi.org/10.1017/S0007114522003191>
7. Cline MG, Burns KM, Coe JB, Downing R, Durzi T, Murphy M, et al. 2021 AAHA nutrition and weight management guidelines for dogs and cats. *J Am Anim Hosp Assoc*. 2021;57:153-

78. <https://doi.org/10.5326/JAAHA-MS-7232>
8. Alvarenga IC, Aldrich CG, Shi YC. Factors affecting digestibility of starches and their implications on adult dog health. *Anim Feed Sci Technol.* 2021;282:115134. <https://doi.org/10.1016/j.anifeedsci.2021.115134>
 9. NRC [National Research Council]. Nutrient requirements of dogs and cats. Washington, DC: The National Academies Press; 2006.
 10. AAFCO [Association of American Feed Control Officials]. Official publication. Champaign, IL: AAFCO; 2021. p. 111-236.
 11. FEDIAF [The European Pet Food Industry]. Nutritional guidelines: for complete and complementary pet food for cats and dogs. Bruxelles: FEDIAF; 2021.
 12. Cho J, Seo G, Kim H, Kim W, Ji I. The estimation of current and future market size of pet related industries. *Korean J Agric Manag Policy.* 2018;45:611-29. <https://doi.org/10.30805/KJAMP.2018.45.3.611>
 13. Joo Y, Jo Y, Jo H, Choi HE, Yoon YS. How much are you willing to pay when you travel with a pet? Evidence from a choice experiment. *Curr Issues Tour* 2023;27:2118-33. <https://doi.org/10.1080/13683500.2023.2223912>
 14. German AJ, Holden SL, Serisier S, Queau Y, Biourge V. Assessing the adequacy of essential nutrient intake in obese dogs undergoing energy restriction for weight loss: a cohort study. *BMC Vet Res.* 2015;11:253. <https://doi.org/10.1186/s12917-015-0570-y>
 15. Pujol S, Torrallardona D. Evaluation of in vitro methods to estimate the in vivo nutrient digestibility of barley in pigs. *Livest Sci.* 2007;109:186-8. <https://doi.org/10.1016/j.livsci.2007.01.143>
 16. Adner M, Canning BJ, Meurs H, Ford W, Ramos Ramírez P, van den Berg MPM, et al. Back to the future: re-establishing guinea pig in vivo asthma models. *Clin Sci (Lond).* 2020;134:1219-42. <https://doi.org/10.1042/CS20200394>
 17. Smeets-Peters M. Feeding FIDO: development, validation and application of a dynamic, in vitro model of the gastrointestinal tract of the dog [Ph.D. dissertation]. Wageningen, Netherlands: Wageningen University and Research; 2000.
 18. Cho JH, Kim IH. Evaluation of the apparent ileal digestibility (AID) of protein and amino acids in nursery diets by in vitro and in vivo methods. *Asian-Australas J Anim Sci.* 2011;24:1007-10. <https://doi.org/10.5713/ajas.2011.10435>
 19. Hervera M, Baucells MD, González G, Pérez E, Castrillo C. Prediction of digestible protein content of dry extruded dog foods: comparison of methods. *J Anim Physiol Anim Nutr.* 2009;93:366-72. <https://doi.org/10.1111/j.1439-0396.2008.00870.x>
 20. AOAC [Association of Official Analytical Chemists] International. Official methods of analysis of the AOAC International. 18th ed. Gaithersburg, MD: AOAC International; 2006.
 21. Donadelli RA, Aldrich CG. The effects on nutrient utilization and stool quality of beagle dogs fed diets with beet pulp, cellulose, and Miscanthus grass. *J Anim Sci.* 2019;97:4134-9. <https://doi.org/10.1093/jas/skz265>
 22. Boisen S, Fernández JA. Prediction of the apparent ileal digestibility of protein and amino acids in feedstuffs and feed mixtures for pigs by in vitro analyses. *Anim Feed Sci Technol.* 1995;51:29-43. [https://doi.org/10.1016/0377-8401\(94\)00686-4](https://doi.org/10.1016/0377-8401(94)00686-4)
 23. Penazzi L, Schiavone A, Russo N, Nery J, Valle E, Madrid J, et al. In vivo and in vitro digestibility of an extruded complete dog food containing black soldier fly (*Hermetia illucens*) larvae meal as protein source. *Front Vet Sci.* 2021;8:653411. <https://doi.org/10.3389/fvets.2021.653411>
 24. Seo K, Cho HW, Lee MY, Kim CH, Kim KH, Chun JL. Prediction of apparent total tract

- digestion of crude protein in adult dogs. *J Anim Sci Technol.* 2024;66:374-86. <https://doi.org/10.5187/jast.2024.e20>
25. Le Bon M, Carvell-Miller L, Marshall-Jones Z, Watson P, Amos G. A novel prebiotic fibre blend supports the gastrointestinal health of senior dogs. *Animals.* 2023;13:3291. <https://doi.org/10.3390/ani13203291>
 26. Mizukami K, Uchiyama J, Igarashi H, Murakami H, Osumi T, Shima A, et al. Age-related analysis of the gut microbiome in a purebred dog colony. *FEMS Microbiol Lett.* 2019;366:fnz095. <https://doi.org/10.1093/femsle/fnz095>
 27. Song YS, Kim BG. Prediction equations for in vitro ileal disappearance of dry matter and crude protein based on chemical composition in dog diets. *Animals.* 2023;13:1937. <https://doi.org/10.3390/ani13121937>
 28. Hervera M, Baucells MD, Blanch F, Castrillo C. Prediction of digestible energy content of extruded dog food by in vitro analyses. *J Anim Physiol Anim Nutr.* 2007;91:205-9. <https://doi.org/10.1111/j.1439-0396.2007.00693.x>
 29. Biagi G, Cipollini I, Grandi M, Pinna C, Vecchiato CG, Zaghini G. A new in vitro method to evaluate digestibility of commercial diets for dogs. *Ital J Anim Sci.* 2016;15:617-25. <https://doi.org/10.1080/1828051X.2016.1222242>
 30. Hendriks WH, Thomas DG, Bosch G, Fahey GC Jr. Comparison of ileal and total tract nutrient digestibility of dry dog foods. *J Anim Sci.* 2013;91:3807-14. <https://doi.org/10.2527/jas.2012-5864>
 31. Satterlee LD, Marshall HF, Tennyson JM. Measuring protein quality. *J Am Oil Chem Soc.* 1979;56:103. <https://doi.org/10.1007/BF02671431>
 32. Burrows CF, Kronfeld DS, Banta CA, Merritt AM. Effects of fiber on digestibility and transit time in dogs. *J Nutr.* 1982;112:1726-32. <https://doi.org/10.1093/jn/112.9.1726>
 33. Kawauchi IM, Sakomura NK, Pontieri CFF, Rebelato A, Putarov TC, Malheiros EB, et al. Prediction of crude protein digestibility of animal by-product meals for dogs by the protein solubility in pepsin method. *J Nutr Sci.* 2014;3:e36. <https://doi.org/10.1017/jns.2014.32>
 34. Weber M, Martin L, Biourge V, Nguyen P, Dumon H. Influence of age and body size on the digestibility of a dry expanded diet in dogs. *J Anim Physiol Anim Nutr.* 2003;87:21-31. <https://doi.org/10.1046/j.1439-0396.2003.00410.x>